

## Correction to: Incidence and Risk Factors for Mortality Following Bariatric Surgery: a Nationwide Registry Study

Nasser Sakran<sup>1,2</sup> · Shiri Sherf-Dagan<sup>3</sup> · Orit Blumenfeld<sup>4</sup> · Orly Romano-Zelekha<sup>4</sup> · Asnat Raziel<sup>5</sup> · Dean Keren<sup>2,6</sup> · Itamar Raz<sup>7</sup> · Dan Hershko<sup>1,2</sup> · Ian M. Gralnek<sup>2,8</sup> · Tamy Shohat<sup>4,9</sup> · David Goitein<sup>5,9,10</sup>

Published online: 25 April 2018  
© Springer Science+Business Media, LLC, part of Springer Nature 2018

### Correction to: Obesity Surgery

<https://doi.org/10.1007/s11695-018-3212-1>

In Table 5 the *P* value for the parameter “More than one chronic disease” is incorrect. The correct value is 0.387, not 0.0387.

---

The online version of the original article can be found at <https://doi.org/10.1007/s11695-018-3212-1>

---

✉ Nasser Sakran  
sakranas@gmail.com

Shiri Sherf-Dagan  
shirisherf@gmail.com

Orit Blumenfeld  
orit.blumenfeld@moh.health.gov.il

Orly Romano-Zelekha  
orly.romano@moh.health.gov.il

Asnat Raziel  
asnatraziel@aol.com

Dean Keren  
dr.deankeren@gmail.com

Itamar Raz  
ntv502@netvision.net.il

Dan Hershko  
dan\_he@clalit.org.il

Ian M. Gralnek  
ian\_gr@clalit.org.il

Tamy Shohat  
tamar.shohat@moh.health.gov.il

David Goitein  
david.goitein@sheba.health.gov.il

Extended author information available on the last page of the article

**Table 5** Adjusted hazard ratios for mortality among patients following bariatric surgery (Cox model)

Parameter	HR of death (95% CI)	<i>P</i> value
Age	1.06 (1.04–1.09)	<0.001
Gender (male)	1.94 (1.16–3.25)	0.011
Marital status (married)	0.43 (0.26–0.71)	0.001
BMI	1.08 (1.05–1.11)	<0.001
Hypertension	1.42 (0.76–2.66)	0.275
Type 2 diabetes	1.74 (0.96–3.14)	0.065
Sleep apnea	1.08 (0.59–1.97)	0.802
Depression	2.38 (1.25–4.52)	0.008
Previous CVA	1.57 (0.37–6.53)	0.535
Public hospital (compared to private)	1.71 (0.98–2.98)	0.059
Laparoscopy (compared to open access <sup>a</sup> )	0.30 (0.09–1.03)	0.056
More than one chronic disease	0.68 (0.29–1.61)	0.387
Recurrent surgery	0.72 (0.31–1.65)	0.445
Different types of surgery (SG compared to other types)	0.71 (0.40–1.25)	0.243

Data were available for  $N = 22,777$  (of them, 65 events of death)

*HR* hazard ratio, *BMI* body mass index, *SG* sleeve gastrectomy

<sup>a</sup> Included open and laparoscopic conversion to open procedures

## Affiliations

Nasser Sakran<sup>1,2</sup> · Shiri Sherf-Dagan<sup>3</sup> · Orit Blumenfeld<sup>4</sup> · Orly Romano-Zelekha<sup>4</sup> · Asnat Raziel<sup>5</sup> · Dean Keren<sup>2,6</sup> · Itamar Raz<sup>7</sup> · Dan Hershko<sup>1,2</sup> · Ian M. Gralnek<sup>2,8</sup> · Tamy Shohat<sup>4,9</sup> · David Goitein<sup>5,9,10</sup>

<sup>1</sup> Department of Surgery A, Emek Medical Center, 21 Yitzhak Rabin Blvd, 1834111 Afula, Israel

<sup>2</sup> Rappaport Faculty of Medicine, Technion Israel Institute of Technology, Haifa, Israel

<sup>3</sup> Department of Nutrition, Assuta Medical Center, Tel-Aviv, Israel

<sup>4</sup> Israel Center for Disease Control, Ministry of Health, Tel Hashomer, Israel

<sup>5</sup> Assia Medical Group, Assuta Medical Center, Tel-Aviv, Israel

<sup>6</sup> Department of Gastroenterology, Bnai-Zion Medical Center, Haifa, Israel

<sup>7</sup> Diabetes Unit, Hadassah University Hospital, Jerusalem, Israel

<sup>8</sup> Ellen and Pinchas Mamber Institute of Gastroenterology and Hepatology, Emek Medical Center, Afula, Israel

<sup>9</sup> Sackler School of Medicine, Tel Aviv University, Tel-Aviv, Israel

<sup>10</sup> Department of Surgery C, Sheba Medical Center, Tel Hashomer, Israel