Correction to: Nutrition, Physical Activity, and Prescription of Supplements in Pre- and Post-bariatric Surgery Patients: a Practical Guideline

Mastaneh Rajabian Tabesh¹ · Faezeh Maleklou¹ · Fatemeh Ejtehad¹ · Zahra Alizadeh^{1,2}

Published online: 5 December 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to: Obesity Surgery (2019) 29:3385–3400

<https://doi.org/10.1007/s11695-019-04112-y>

Post-surgery treatment: 1000 mg of folate daily: 1000 microgram of folate daily)

In Table 7 the following corrections are required:

Vitamin B12 (cobalamin):

Post-surgery recommendation: mg must be changed to microgram (350–500 mg: 350–500 microgram/ 1000 mg monthly: 1000 microgram monthly)

Folate (folic acid):

Post-surgery recommendation: mg must be changed to microgram (400–800 mg: 400–800 microgram/ 800–1000 mg: 800–1000 microgram)

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11695-019-04112-y>

✉ Zahra Alizadeh
z_alizadeh@tums.ac.ir

Mastaneh Rajabian Tabesh
mastaneh.tabesh@ut.ac.ir

Faezeh Maleklou
faezeh.maleklou88@gmail.com

Fatemeh Ejtehad
ftmhejtehad@gmail.com

¹ Sports Medicine Research Center, Tehran University of Medical Sciences, Tehran, Iran

² Department of Sports and Exercise Medicine, Tehran University of Medical Sciences, No. 7, Ale-ahmad Highway, Opposite of the Shariati Hospital, Tehran 14395-578, Iran