



Correction to: Comparison of Linear Versus Circular-Stapled Gastroenterostomy in Roux-en-Y Gastric Bypass: a Nationwide Population-Based Cohort Study

Marleen M. Romeijn^{1,2} · Stijn van Hoef¹ · Loes Janssen¹ · Kelly G. H. van de Pas^{1,2} · François M. H. van Dielen¹ · Arijan A. P. M. Luijten¹ · Kevin W. A. Göttgens¹ · Jan Willem M. Greve^{2,3} · Wouter K. G. Leclercq¹

© The Author(s) 2021

Correction to: Obesity Surgery.

<https://doi.org/10.1007/s11695-021-05436-4>

In **Table 1**, the headings in the top row are missing. The names in this heading represent the two groups of the study: CSA (left column) and LSA (right column).

The corrected table is shown below.

The online version of the original article can be found at <https://doi.org/10.1007/s11695-021-05436-4>

✉ Marleen M. Romeijn
m.romeijn@maastrichtuniversity.nl; Bariatric.resurge@mmc.nl

Stijn van Hoef
stijnvanhoef@gmail.com

Loes Janssen
Loes.Janssen@mmc.nl

Kelly G. H. van de Pas
kelly.van.de.pas@mmc.nl

François M. H. van Dielen
f.vandienen@mmc.nl

Arijan A. P. M. Luijten
A.Luijten@mmc.nl

Kevin W. A. Göttgens
Kevin.Goettgens@mmc.nl

Jan Willem M. Greve
j.greve@zuyderland.nl

Wouter K. G. Leclercq
w.leclercq@mmc.nl

¹ Department of Surgery, Máxima Medical Center, Veldhoven, The Netherlands

² Research School NUTRIM, Department of Surgery, Maastricht University Medical Center, Maastricht, the Netherlands

³ Department of Surgery, Zuyderland Medical Center, Heerlen, the Netherlands

Table 1 Baseline characteristics of the study population

	CSA, n =881	LSA, n =11,587	p value
Gender, no. (%)			
Female	713 (80.9)	9541 (82.3)	.291
Age (years)	45.2±10.5	45.1±10.6	.885
Preoperative comorbidities, no. (%)			
Hypertension	333 (37.8)	4269 (36.8)	.571
Type II diabetes mellitus	143 (16.2)	2595 (22.4)	<.001*
Hyperlipidaemia	169 (19.2)	2523 (21.8)	.071
Gastroesophageal reflux disease	89 (10.1)	1667 (14.4)	<.001*
OSAS	265 (30.1)	2231 (19.3)	<.001*
Osteoarthritis	344 (39.0)	5699 (49.2)	<.001*
Preoperative weight (kg, ±SD)	122.7±18.6	122.7±17.9	.994
Preoperative BMI (kg/m ² ±SD)	42.7±4.9	42.8±4.7	.265
Laparoscopic, no. (%)	879 (99.8)	11,573 (99.9)	.396
Length of biliopancreatic limb (cm±SD)	72.9±15.3	90.5±38.6	<.001*
Length of alimentary limb (cm±SD)	145.8±9.7	133.6±33.7	<.001*
Length of hospital stay (days ±SD)	2.3±1.7	1.5±2.7	<.001*
Number of readmission (<30 days), no. (%)	25 (2.8)	283 (2.4)	.466
Postoperative complication <30 days, no. (%)			
CD grade I	19 (2.2)	89 (.8)	<.001*
CD grade II	25 (2.8)	99 (.9)	<.001*
CD grade IIIa	3 (.3)	29 (.3)	.610
CD grade IIIb	41 (4.7)	382 (3.3)	.032*
CD grade IVa	4 (.5)	15 (.1)	.017*
CD grade IVb	-	9 (.1)	.408
CD grade V	-	1 (.0)	.783
Type of complication, no. (%)			
Major bleeding	21 (2.4)	136 (1.2)	.002*
Anastomotic leakage	1 (.1)	43 (.4)	.214
Intra-abdominal abscess	2 (.2)	13 (.1)	.343
Wound infection	1 (.1)	14 (.1)	.952
Intestinal obstruction	1 (.1)	28 (.2)	.447
Anastomotic stricture	0 (.0)	1 (.0)	.783

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not

permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

>