

# Astonishing Diversity of Natural Surfactants: 5. Biologically Active Glycosides of Aromatic Metabolites

Valery M. Dembitsky\*

Department of Organic Chemistry and School of Pharmacy, Hebrew University, Jerusalem, Israel

**ABSTRACT:** This review article presents 342 aromatic glycosides, isolated from and identified in plants and microorganisms, that demonstrate different biological activities. They are of great interest, especially for the medicinal and/or pharmaceutical industries. These biologically active natural surfactants are good prospects for the future chemical preparation of compounds useful as antioxidant, anticancer, antimicrobial, and antibacterial agents. These glycosidic compounds have been classified into several groups, including simple aromatic compounds, stilbenes, phenylethanoids, phenylpropanoids, naphthalene derivatives, and anthracene derivatives.

Paper no. L9814 in *Lipids* 40, 869–900 (September 2005).

The father of modern medicine was Hippocrates, who lived sometime between 460 and 377 BC. Hippocrates left historical records of pain-relief treatments, including the use of a powder made from the bark and leaves of the willow tree to help heal headaches, pains, and fevers. Since the time of Hippocrates, people have chewed on willow leaves and bark to alleviate pain. About 2,000 years later, aspirin began being manufactured from a compound extracted from the tree. As it turns out, that family of compounds, called phenolic glycosides, also plays a key role in plant health, making many plants tough and disease resistant.

Aromatic compounds, including phenolic compounds, are widely distributed in the plant kingdom (2). These compounds are essential for the growth and reproduction of plants and are produced as a response to defend injured plants against pathogens. In recent years, interest has been growing in aromatic compounds and their presumed role in the prevention of various

degenerative diseases, such as cancer and cardiovascular diseases (3–5). The antioxidant activities of aromatic compounds and their possible use in processed foods as natural antioxidants have gained considerable recognition in recent years (3,6,7). Terrestrial vascular plants and their aquatic counterparts contain several types of phenylpropanoid metabolites. In vascular plants, this pathway provides major classes of organic compounds of specialized function and structure. These include neolignans and lignans, structural cell wall polymers (lignins and the aromatic portion of suberins), proanthocyanidins (condensed tannins, i.e., the phenylalanine-derived portion), and related compounds. Without this pathway, the normal growth, development, and survival of both woody and nonwoody plants would not be possible. Indeed, since approximately 10–45% of all vascular plant material is of phenylpropanoid origin, these metabolites constitute a major sink of organic carbon. Many of these molecules can occur in plants in the form of glycosides, which means they have one or more simple sugars, such as glucose, galactose, mannose, xylose, or another sugar attached to them. Not only are thousands of aromatic compounds known, but some of the smaller molecules also can polymerize, or join together to form larger and more complex molecules. Different kinds of molecules might be found in one polymer (8–11).

Phenolic compounds (a molecule containing an aromatic ring that bears one or more hydroxyl groups is referred to as “phenolic”) can also be bound to certain proteins, polysaccharides, or lipids (12–18), and as if this were not complex enough, some molecules can have multiple isomers, forms in which their atoms are arranged slightly differently.

Simple and polyphenolic compounds and/or their glycosides of fruits and vegetables may play an important role in physiological functions related to human health (19–26). Different polyphenolics may have varied biological activities, such as having anticancer, anti-inflammatory, cytostatic, cytotoxic, antimetabolic, neurotoxic, antiplatelet, antibacterial, antifungal, aggregational, or antiradical activities, as well as antioxidant or cardioprotective activities (3,20,22,24,27–29).

Aromatic glycosides, representative of a large group of natural compounds mainly originating from plant species that can be separated on several groups, including simple aromatic glycosides, stilbene glycosides, phenylethanoid and phenylpropanoid glycosides, naphthalene glycosides, and glycosides of anthracene derivatives (anthraquinones, angucyclines, oxantrones, and anthrones), as considered in this review article.

\*Address correspondence at Department of Organic Chemistry, P.O. Box 39231, Hebrew University, Jerusalem 91391, Israel.  
E-mail: dvalery@cc.huji.ac.il

For the previous article in this series, see Reference 1.

Abbreviations: D-GalN, D-galactosamine; EC<sub>50</sub>, the median effect concentration, being a statistically derived concentration of a substance that can be expected to cause (i) an adverse reaction in 50% of organisms or (ii) a 50% reduction in growth or in the growth rate of organisms; GI<sub>50</sub>, the concentration needed to reduce the growth of treated cells to half that of untreated (i.e., control) cells; HUVEC, human umbilical vein endothelial cells; IC<sub>50</sub>, concentration at which growth or activity is inhibited by 50% (applies to ligand and growth inhibition); K<sub>i</sub>, prolyl hydroxylase activity, expressed as mol/mg protein; LLC, Lewis lung carcinoma; LPS, lipopolysaccharide; MIC, minimum inhibitory concentration of an antibiotic that inhibits a bacterium; MPP<sup>+</sup>, 1-methyl-4-phenylpyridinium ion; OLE, olive leaf extract; SDG, secoisolaricresinol diglycoside **245**; TNF- $\alpha$ , tumor necrosis factor  $\alpha$ .

## SIMPLE AROMATIC GLYCOSIDES


Simple aromatic (including phenolic) glycosides have been isolated mainly from plants, although a few compounds have been isolated from microorganisms. Compounds in this group often also possess alcoholic, aldehydic, and carboxylic acid groups. Medicinal herbs constitute an important source of biologically active compounds, and they have given us a number of important drugs that are mainstays of treatment in synthetic systems of medicine (30–32).

The Egyptians (*ca.* 1500 BC) recorded a collection of recipes for medicines that included a recipe using an infusion of dried myrtle leaves (which contain salicylic acid) to relieve back pain. The effects of aspirin-like substances have been known since the ancient Romans recorded the use of willow bark as a fever fighter. The leaves and bark of the willow tree contain a substance called salicin, **1**, a naturally occurring compound similar to acetylsalicylic acid (aspirin) (33). Aspirin is probably the most successful medicine of all time, and new uses for it continue to be discovered (34–37).

In 1828, Johann Buchner, a professor at Munich University, isolated pure salicin from willow bark (*Salix alba*). Two Italians, Luigi Brugnatelli and Carlo Fontana, had in fact already obtained salicin in 1826, but in a highly impure form. By 1829, Henri Leroux, a French chemist, had improved the extraction procedure to obtain about 30 g from *S. alba* bark. In 1838, Raffaele Piria, an Italian chemist then working at the Sorbonne in Paris, split salicin into a sugar and an aromatic component (salicylaldehyde) and converted the latter, by hydrolysis and oxidation, to an acid of crystallized colorless needles, which he named salicylic acid (38–43). The name salicin was derived from *salix*, which is the Latin word for willow tree. Many well-known drugs have come from studying natural plant compounds such as salicin, the active constituent found in white willow bark. This herb has been used for centuries to ease discomfort associated with the joints and muscles. Occurring as white crystals or powder, salicin has analgesic, antipyretic, disinfectant, and antiseptic properties, and it has been found in a number of plant species: *S. alba*, *Salix tetresperma*, *Salix fragilis*, *Populus nigra*, *Populus alba*, *Populus tremula*, and *Filipendula ulmaria* (39–44).


Salicin-related compounds, such as gaultherin **2** (formerly known as monotropitoid or monotropitin), monotropin, and spiraein are some of the known compounds present in plant flora having activity similar to salicin. Gaultherin **2** is a glycoside of methyl salicylate that is present in *Gaultheria procumbens* (wintergreen); it is also known as monotropitoid and is

used as a source for the manufacture of salicylic acid (44). Salicin is found in most Salicaceae species, together with other much less abundant glycosides such as populin, tremuloidin, fragilin, glycosmin, picein, salireposide, grandidentatin, triandrin, salicylpopulin (salicyloylpopulin), and salidroside (salicyloylpopulin) (45). Salicylates have long been known as water-soluble compounds derived from a number of plants, particularly willow (*Salix* spp.), meadowsweet (*Spirea* spp. and *F. ulmaria*), sweet birch (*Betula lenta*), white birch (*Betula pendula*), wintergreen (*G. procumbens*), balsam poplar (*Populus balsamifera*), black poplar (*P. nigra*), balm of Gilead (*Populus candicans*), black haw (*Viburnum prunifolium*), and sweet violet (*Viola odorata*), with analgesic, antipyretic and anti-inflammatory properties (46–48). In 1953, Paris and Pointet (49) reported the presence of gaultherin **2** in the bark of *Ostryopsis davidiana*. More recently, Towers *et al.* (50) studied leaf extracts of 22 species of *Gaultheria* after acid or alkali hydrolysis, yielding *p*-hydroxybenzoic, *o*-pyrocatechuic, protocatechuic, gentisic, vanillic, *p*-coumaric, caffeic, and ferulic acids, and found that 13 species contained derivatives of salicylic acid as well; the glycoside gaultherin **2** was isolated from *G. procumbens* and *Gaultheria hispida*. Gaultheria oil (or wintergreen oil), which consists almost solely of methyl salicylate, was formerly distilled on a commercial scale from the leaves of this species. *Gaultheria procumbens*, also known as teaberry, wintergreen, Canada tea, partridge berry, checkerberry, boxberry, wax cluster, spice berry, mountain tea, deerberry, spicy wintergreen, aromatic wintergreen, chink, ground berry, grouse berry, red pollom, redberry tea, hillberry, and ivory plum, is related to the Northwest native evergreen groundcover *Gaultheria shallon*, which was used by Quinault and Klallum Indians for its medicinal properties as a stimulant, antiseptic, astringent, diuretic, emmenagogue, and aromatic, and was also useful in chronic inflammatory rheumatism, rheumatic fever, sciatica, diabetes, all bladder troubles, scrofula, and skin diseases (51). Spiraein **3**, which contains salicylaldehyde, xylose, and glucose, was first isolated from *F. ulmaria* (52) and was later found in other plant species, i.e., *Populus tremuloides*, *Salix purpurea*, *S. fragilis*, *Salix nigricans*, and *Salix rosmarinifolia* (53). Leonuride A **4** and erige-side C **5** were isolated from the aerial parts of *Acanthus ilicifolius* (54). Two new salicin derivative glucosides, 7-*O*-methylsalicin **6** and 7-*O*-(2,6-dihydroxy-benzoyl)salicin **7**, were isolated from the leaves of *Alangium platanifolium* var. *trilobum* (55). Salicin **1** and populin **8** were isolated from hot-water bark extracts of *Populus tomentosa*, *Populus davidiana*, *Populus euphratica*, and *Populus simonii* (56).



HPLC analysis of the bark of *Salix petiolaris* (*S. gracilis* var. *textoris*), which was collected in November, was shown to contain salicin **1**, populin **8**, picein **9**, salicyloisalicin **11**, salireposide **12**, grandidentatin, vimalin, tremulacin **13**, and salicyloisalicin-2-*O*-benzoate **14** (57). Golden root (*Rhodiola rosea*) has been used for many years as an adaptogen in Chinese traditional medicine and was reported by Tolonen *et al.* (58) to have many pharmacological properties; its known phenylpropanoid metabolites, picein **9** and benzyl-*O*- $\beta$ -glucopyranoside **10**, were recently isolated. Three new glycosides, benzyl alcohol  $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2)-[ $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 6)]- $\beta$ -D-glucopyranoside **15**, 2'-*O*- $\beta$ -D-glucopyranosylsalicin **17**, and 2'-*O*- $\beta$ -D-glucopyranosyl-6'-*O*- $\beta$ -D-xylopyranosylsalicin **18**, along with the known benzyl alcohol  $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside **16** and 6'-*O*- $\beta$ -D-xylopyranosyl-salicin **19**, were isolated from the water-soluble fraction of the dried leaves of *Alangium chinense* (59).

Dill (*Anethum graveolens*) has been cultivated in Europe since antiquity and is popular aromatic herb and spice. Its seed also has been used for medicinal purposes to relieve digestive problems and to stimulate milk for nursing mothers. Dill water was believed to have a soothing effect on the digestive system and was given to babies to relieve hiccups and colic, and essential oils from dill have antimicrobial (60,61), antihyperlipidemic, and antihypercholesterolemic activities (62). From the water-soluble fraction of an ethanol extract of dill seed, which has been used as a spice and medicine, 33 compounds were obtained, including the known compound **10** and the new aromatic glucosides **20–25** (63).



Important crude drugs from the *Atractylodes* plant (Japanese name *jutsu*), *Atractylodes* rhizome [*Atractylodes japonica* and *Atractylodes ovata* (Japanese name *byaku-jutsu*)], and *Atractylodes lancea* rhizome [*A. lancea* and *Atractylodes chinensis* (Japanese name *so-jutsu*)] are used as diuretics and stomachics (64). The new aromatic glycosides **26–29** were isolated together with 10 known compounds, including **25**, from the water-soluble fraction of the fresh rhizome of *A. japonica* (65).

Coriander (*Coriandrum sativum*) has one of the longest recorded histories of any of the spices. There is evidence that this native of the Mediterranean and Near East has been in use since 5000 BC. Its seeds have been found in Bronze Age ruins on the Aegean Islands and in the tombs of the pharaohs. Coriander is mentioned in the Ebers Papyrus and in the books of Moses. It was grown in Assyria and Babylon and was used in minced meat dishes, sausages, and stews. The seeds have been used as a drug for indigestion, against worms, and as a component of embrocations for rheumatism and pains in the joints. From the water-soluble portion of a methanol extract of coriander fruits, two new aromatic glycosides, **30** and **31**, were isolated together with the known compounds **4**, **24**, and **25** (66).

The seed of the *Amomum* plant is one of the most ancient and highly valued spices in the world. The seed of *Amomum xanthioides* (Zingiberaceae) also has been used for medicinal purposes; it is prescribed in traditional medicines for stomachic and digestive disorders and contains the simple aromatic glycosides **24** and **32** (67).


The Aceraceae plant *Acer nikoense* is indigenous to Japan (Japanese name, *megusurinoki*), and its stem bark has been used as a folk medicine for the treatment of hepatic disorders and eye diseases. The new aromatic glycoside nikoenoside **33** and also kelampayoside A **34**, **35**, and **36** were isolated from the stem bark of *A. nikoense* (68).

Plants of the genus *Eugenia* showed antidiabetic (69), hypoglycemic (70), and also antibacterial and antifungal activities against gram-positive bacteria (*Bacillus subtilis*, *Staphylococcus aureus*), gram-negative bacteria (*Salmonella typhimurium*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, and *Escherichia coli*), and fungal strains (*Candida albicans*, *Cryptococcus neoformans*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger*, *Rhizopus* sp., *Trichophyton rubrum*, *Trichophyton mentagrophytes*, and *Microsporium gypseum*) (71). Lambertianoside **37**, a novel phenylglycoside with anti-HIV activity, has been isolated from the aqueous extract of leaves from *Eugenia lambertiana* [HIV EC<sub>50</sub> (i.e., median effect concentration causing an adverse reaction in 50% of organisms or a 50% reduction in the growth or growth rate of an organism), 3 µg/mL; HIV IC<sub>50</sub> (i.e., 50% inhibition concentration), 25 µg/mL] (72).

Pumpkin, the fruit of *Cucurbita moschata* (Cucurbitaceae), has been a popular vegetable in cooking since antiquity, and salted, roasted pumpkin seeds are used in several countries as snacks; pumpkin seeds are also an important traditional Chinese medicine used in the treatment of cestodiasis, ascariasis, and schistosomiasis. Pharmacological studies on the seeds have demonstrated hepatoprotective (73) and antitumor activities (74). The new phenolic glycosides cucurbitosides A–E, **38–42**, were isolated from the seeds of *C. moschata* (75). The dried stem bark of *Albizia julibrissin* (Leguminosae) has been used in Chinese herbal medicine to treat insomnia, diuresis, asthenia, ascaricide, and confusion. From the stem bark of *A.*




*julibrissin*, two new phenolic glycosides were isolated, albibrissinosides A **43** and B **44**. Albibrissinoside B **44** showed a strong 1,1-diphenyl-2-picrylhydrazyl radical-scavenging activity, with an  $IC_{50}$  value of 10.2  $\mu$ M (positive control, L-ascorbic acid  $IC_{50}$  10.4 mM), whereas albibrissinoside A **43** showed a weak scavenging activity, with an  $IC_{50}$  value of 253  $\mu$ M (76).




Three new phenolic compounds, 2-methoxy-5-(1'2'3'-trihydroxypropyl)-phenyl-1-*O*-(6''-galloyl)- $\beta$ -D-glucopyranoside **45**, 2-methoxy-5-hydroxymethyl-phenyl-1-*O*-(6''-galloyl)- $\beta$ -D-glucopyranoside **46**, and 1-galloyl- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-galactopyranoside **47**, were isolated from the leaves of *Baseonema acuminatum* (Asclepiadaceae) (77). The antimicrobial activity of all compounds was evaluated *in vitro* against the bacteria *S. aureus* (two strains), *Bacillus cereus*, *B. subtilis*, *E. coli*, and *Salmonella typhimurium* and against three strains of *C. albicans*. The new compounds **45** and **46** showed antifungal activity against two clinically isolated *C. albicans* strains and against *C. albicans* ATCC 2091 [minimum inhibitory concentration (MIC) values in the range of 25–100  $\mu$ g/mL]. Compound

**47** inhibited only one strain of *C. albicans* at the maximum concentration used. Melilotoside **48** was isolated from *Ephedra nebrodensis* (78). The new phenolic glycosides quercetin  $\beta$ -D-glucopyranosyl 4-*O*- $\beta$ -D-glucopyranosylcaffeate **49**, methyl 3-*O*- $\beta$ -D-glucopyranosyl-5-hydroxycinnamate **50**, and  $\beta$ -D-glucopyranosyl 4-*O*- $\beta$ -D-glucopyranosylbenzoate **51** were isolated from the flowers of *Moricandia arvensis* (79). Compounds **49–51** showed antioxidant activity, and **49** proved to possess the most potent radical-scavenging activity.




*Curculigo orchioides* (Amaryllidaceae) is a tiny herbal plant widely distributed in China, India, Malaysia, Japan, and Australia. Its rhizomes have the properties of warming the kidneys, invigorating yang, expelling cold, and eliminating dampness, and are used as the traditional Chinese medicine *xian mao* to cure impotence, enuresis, cold sperm, cold pain of the back and knee, and numbness of the limbs (80). In India, the tuberous roots of this plant are considered to be a tonic, alterative, demulcent, diuretic, and restorative, and are used as a poultice for itching and skin diseases (81). Three phenolic glycosides, curculigosides **52**, **53**, and **54**; orcinol glycoside **55**; and anacardoside **56** have been isolated from the rhizomes of *C. orchioides* and have been described in some recently published papers (81–88).

Three rare new chlorophenyl glycosides, curculigin A **57**, B **58**, and C **59**, have been isolated from *C. orchioides* (82,86). A new chlorine-containing phenoloid named capitulatin A **60** has been isolated from the rhizomes of *Curculigo capitulata* (89).




Three chlorophenyl glycosides **61–63** were extracted from the tissues of *Lemna minor* exposed to 2,4-dichlorophenol (90). Chlorophenols are an important class of xenobiotics used in a variety of biocides and have been shown to be resistant to microbial degradation. Enzyme-catalyzed hydrolysis with  $\beta$ -glucosidase was found to be ineffective in releasing the  $\beta$ -glucosides with chemical modifications at C-6. The presence of these

glucoconjugates confirmed that *L. minor* was capable of xenobiotic uptake and transformation. The products identified suggested that chlorophenols were incorporated into the vacuoles and cell walls of *L. minor* (90). The chlorophenol glucuronides **64** and **65** were isolated from human urine (91).  $\beta$ -(2-Chloroethyl)-D-glucoside **66** was isolated from the green plant *Taraxacum officinale* (92). Chlorophenol glycosides with the common structure **67** can be synthesized by aquatic plants and have been isolated from aquatic environments (93).



From the rhizomes of *Curculigo pilosa*, two benzylbenzoate diglucosides, piloside A **68** and B **69**; the *O*-methyl derivative **70**; nyasicoside **71**; and a glucosyl-fused norlignan, pilosidine **72**, previously obtained only as the tetra-*O*-methyl derivative **73**, were isolated. Pilosidine **72** showed a facilitating effect on adrenaline-evoked contractions in rabbit aorta (85). Nyasicoside **71** also was found in the rhizomes of *C. capitulata* (94). A monoglucoside, shipamanine **77**, and the related compounds obtuside A **74**, obtuside B **75**, and hypoxoside **76** were isolated from the rhizomes of *Hypoxis obtusa* (95,96). Shipamanine **77** and hypoxoside **76** showed antiviral, antitumoral, and analgesic activities.



The poisonous perennial shrub *Breynia officinalis* (Euphorbiaceae) grows in Okinawa, Taiwan, and southern China. It is administered orally as a remedy for healing wounds and edema, as an ointment, for syphilis, and for intestinal hemorrhage due to overwork (97). The novel phenolic glycosides **78–82** have been isolated from the leaves of *B. officinalis* collected on Okinawa Island (98).


The herbaceous plant *Pyrola japonica* (Pyrolaceae) is widespread in Korea, Japan, and China; the leaves are used as a detoxicant for the bites of snakes, insects, and dogs, and as a drug to cure yellowish and bloody sputum (99). Five new phenolic glycosides, **83–87**, were isolated from the whole plants of *P. japonica* (100), and a new cyclic compound, jujuphenoside **88**, was isolated from the seeds of *Ziziphus jujuba* var. *spinosa* (101).

Three active glycosides (including sinenoside **1** **89**), which afford protection to red blood cell membranes to resist hemolysis induced by the peroxy radical initiator 2,2'-azo-bis-(2-amidinopropane) dihydrochloride, were isolated from *Ligustrum sinense* (102). Dracunculifoside **K** **90** and **N** **91**, and 10

other glycosides were isolated from the aerial part of *Baccharis dracunculifolia* (Compositae); these compounds have an (*E*)-caffeoyl group like dracunculifosides A–J, which were reported previously (103).

Syringic acid 4- $\beta$ -D-glucoside **92** was isolated from the aerial part of *Erigeron breviscapus* (104), and a lignan, glycoside (–)-secoisolaricresinol-*O*- $\alpha$ -L-rhamnopyranoside **93**, was isolated from the aerial parts of *Rubus amabilis* (105). Dictamninside A **94** was isolated from the whole herb of *E. breviscapus* (106).


Four erythro-anethole glycol monoglucosides, **95–98**, and two new glycosides of *p*-hydroxyphenylpropylene glycol were isolated from the methanolic extract of the seed of *Foeniculum vulgare* (Umbelliferae). This is the first report of the isolation of all four *O*- $\beta$ -D-glucopyranosides, which have the glucosyl unit attached to the hydroxy group of the erythro-glycol function of individual stereoisomeric aglycons (107).



Aromatic glycosides, named hovetrichoside **99** and **F 100**, shashenoside **I 101**, arillanin **F 102**, and kelampayoside **A 103**, have been isolated from the bark of *Hovenia trichocarea* (108).

The HIV-inhibitory activity of the extract was traced to polymeric tannins, whereas 2-hydroxy-3-*O*-β-D-glucopyranosyl-benzoic acid **104** was inactive in the National Cancer Institute's primary anti-HIV screening. This glycoside was isolated from *Geniostoma antherotrichum* (109).

The unusual compound (7*S*,8*S*)-3,5,8-trihydroxy-7-(4-hydroxyphenyl)-bicyclo[4.2.0]-octa-1,3,5-trien-2-yl-β-D-glucopyranoside **105** was isolated from an aqueous extract of *Polygonum multiflorum* (110).



Many other simple aromatic glycosides with biological activities have been isolated from plant species, and readers of *Lipids* can find additional interesting information in a number of review articles (111–122).

## STILBENES

Stilbenes—or 1,2-diphenylethylenes, of which more than 300 are known—are biologically active compounds that have shown different activities, such as nitric oxide production inhibition and antibacterial, antifungal, antioxidant, anti-inflammatory, anticancer, and antimalarial activities (123–128). These compounds and/or their glycosides also have attracted much attention for their biological effects, including antioxidant cyclooxygenase-I and -II inhibitory (129), antiplatelet-aggregation (130), antifungal (131), tyrosinase-inhibitory (132), anti-HIV-1, and cytotoxic effects (133), which play an important role in photochemical reactions (134,135). They have been found in many families of higher plants but mainly in Vitaceae, Gnetaceae, Polygonaceae, Liliaceae, Moraceae, and Cyperaceae. According to the National Library of Medicine (National Center for Biotechnology Information, Bethesda, MD), more than 28,000 research papers have been published on the activities of stilbenes and related compounds.

Two novel bibenzyl glucosides, together with their known aglycons batatasin III **106** and 3'-*O*-methylbatatasin III **107**, were isolated from the tubers of *Pleione bulbocodioides* (136); icaricide A6 **108** also was isolated from the aerial parts of *Epimedium grandiflorum* var. *thunbergianum* as a minor constituent (137).


The tree *Acer mono* (*A. truncatum* subsp. *mono*, Aceraceae), known as *gorosoe* in Korean, is widely distributed in Korea, China, and Japan. The leaves of *A. mono* have been used in Korean folk medicine for hemostasis, and the roots have been used for the treatment of arthralgia and cataclasis. The sap of *A. mono* has been used to treat difficulty in urination, constipation, other gastrointestinal disorders, gout, and neuralgia (138). Two new stilbene glycosides, **109** and **110**, were isolated from the leaves of *A. mono* along with seven known compounds. Compounds **109** and **110** showed significant hepatoprotective activities (97.3% protection against toxicity induced by H<sub>2</sub>O<sub>2</sub> at 50 μg/mL) in primary cultures of rat hepatocytes (139). Compounds **109** and **111**, isolated from the stem bark extract of *Boswellia papyrifera* (140), exhibited significant inhibition of phosphodiesterase I and xanthine oxidase (140).

Resveratrol **112**, a well-known stilbene glycoside, is widely distributed in grapes and wines (141). The highest contents of resveratrol and resveratrol glycoside **112** were observed in wines made with the mourvedre, pinot, and cabernet sauvignon varieties; the lowest were in those made with gamay, cabernet franc, or grenache (141). It has also been found in many other plant species (142). Resveratrol **112** exhibited relatively higher cytotoxic activities against human oral squamous cell carcinoma and salivary gland tumor cell lines than against normal human gingival fibroblasts (143). Other pharmacological activities of resveratrol and their glycosides have



recently been reviewed (144). 4'-O- $\beta$ -D-(6''-O-Galloyl)-glucopyranoside **113** was isolated from the Japanese rhubarb, *shinshu daio* (145). *Trans*-piceid **114**, *trans*-astringin **115**, isorhapontigenin 3-O- $\beta$ -D-glucoside **116**, and pinostilbenoside **117** were isolated from the bark of some *Pinaceae* species (146). Compounds **115** and **116** also were found in the Colorado blue spruce (*Picea pungens*) and Engelmann spruce (*Picea engelmannii*) (147). Resveratrolsides **112** and pinostilbenoside **117** were found in the inner bark of *Pinus sibirica* (148). Hydroxystilbene glycoside **111** was isolated from the roots of the Italian tree *Terminalia sericea* (149), and **118** was isolated from the Australian *Eucalyptus sideroxylon* (149). Deoxyrhaponticin **119**, rhaponticoside **120**, piceatannol 3'-O- $\beta$ -D-glucopyranoside **121**, and compound **117** were obtained from an extract of the rhizomes of *Rheum rhaponticum* (150).

*trans*-Piceid **114** and compound **122** were found in grapes and wine (151). Kimura and Okuda (151) studied the effects of stilbene glucosides, which were isolated from medicinal plants and grapes, on tumor growth and lung metastasis in mice bearing highly metastatic Lewis lung carcinoma (LLC) tumors, and also the inhibitory effects of stilbene glucosides on the differentiation of human umbilical vein endothelial cells (HUVEC). Piceid **114** inhibited the DNA synthesis in LLC cells at a concentration of 1000  $\mu$ M, but not at lower concentrations (10–100  $\mu$ M). 2,3,5,4'-Tetrahydroxystilbene-2-O-D-glucoside **122** also inhibited DNA synthesis in LLC cells (IC<sub>50</sub>, 81  $\mu$ M). In addition, both stilbene glucosides inhibited the formation of capillary-like tube networks (angiogenesis) of HUVEC at concentrations of 100–1000  $\mu$ M. It was suggested that the antitumor and antimetastatic activities of the stilbene glucosides piceid and 2,3,5,4'-tetrahydroxystilbene-2-O-D-glucoside **122** might be due to the inhibition of DNA synthesis in LLC cells and angiogenesis of HUVEC. The pharmacological effect of stilbene glycoside **122** on two kinds of cell models of amentia imitation was observed; the protective mechanism of stilbene glycoside, the main effective component of the tuber fleeceflower root on nerve cells, was explored; and a basis was provided for clarification of the efficacy of a Chinese drug compound from the tuber fleeceflower root (152). The results indicated that stilbene glycoside **122** significantly promoted cell viability and reduced cell membrane damage in  $\beta$ -amyloid 25–35 and H<sub>2</sub>O<sub>2</sub>, treating nerve cells. This drug may antagonize the cell damage initially induced by hydrogen peroxide and prevent subsequent toxicity of the amyloid  $\beta$ -protein in nerve cells. This nerve protection of the drug will be the basis for treating Alzheimer's disease and other neurodegenerative diseases.




The tetrahydroxystilbene glycosides **122–125** were isolated from *P. multiflorum* (153). These isolated tetrahydroxystilbene glucosides are used to prepare drug and/or health-care foods for the prevention and treatment of cerebrovascular and cardiovascular diseases.

Stilbenes isolated from the rhizomes of the Korean rhubarb *Rheum undulatum* and related compounds were investigated for their antiallergenic activities (154). Thus, the effects of substitutes on the aromatic rings of stilbenes have indicated that stilbenes substituted with methoxyl groups at the 3-, 5-, and 40-positions exhibit higher activity than those with hydroxyl groups. Moreover, the activity is dramatically decreased when there is no substitute on aromatic rings. Substitution of the glycoside moiety on the ring dramatically decreases the activity, which can be observed from the inhibitory activities of rhaponticin [**126**, IC<sub>50</sub> > 100 (9.4)  $\mu$ M], and rhaponticin 2-O-gallate (**127**, 45.6  $\mu$ M), isorhapontin (**128**, 23.5  $\mu$ M), and piceatannol 30-O- $\beta$ -D-glucopyranoside (**131**, -7.1  $\mu$ M) compared with

nonglycosidic compounds such as rhapontigenin (11  $\mu\text{M}$ ), isorhapontigenin (12  $\mu\text{M}$ ), and piceatannol (24  $\mu\text{M}$ ). The results revealed that 3,5,40-trimethylpiceatannol exhibited the most potent inhibition against  $\beta$ -hexosaminidase release as a marker of degranulation in RBL-2H3 cells ( $\text{IC}_{50}$  2.1  $\mu\text{M}$ ), followed by trimethylresveratrol ( $\text{IC}_{50}$  5.1  $\mu\text{M}$ ). Glycosides **128–130** also were found in extracts from the root bark of *Picea abies* (155). The stilbenes rutinoid **132** and dihydrochalcone glucoside **133** were isolated from *Guibourtia tessmanii* (156).


Species of the Dipterocarpaceae family are a rich source of polyphenols, particularly resveratrol oligomers, and they show various biological activities such as being cytotoxic, antiviral, and anti-inflammatory (157). Resveratrol glycosides **134–136**, together with known compounds, were isolated from the acetone-soluble part of the stem of *Upuna borneensis* (Dipterocarpaceae) (157).



Two new stilbene dimer glucosides, resveratrol (*E*)-dehydrodimer 11-*O*- $\beta$ -D-glucopyranoside **137** and resveratrol (*E*)-dehydrodimer 11'-*O*- $\beta$ -D-glucopyranoside **138**, were isolated from grape *Vitis vinifera* cell cultures (158). The structures and stereochemistry of the new compounds were determined on the basis of spectroscopic data analysis. Compounds **137** and **138** are dimers that belong to a new type of oligostilbene formed from a resveratrol unit and a resveratrol glucoside unit. Compound **137** exhibited nonspecific inhibitory activity against cyclooxygenase-I and -II, with an  $\text{IC}_{50}$  value of 5  $\mu\text{M}$ .

The genus *Gnetum* (Gnetaceae) is known to contain abundant stilbene derivatives that have interesting biological activities, such as inducing apoptosis in colon cancer (159) and reducing blood sugar (160), revealing the importance of plants


containing stilbenoids as resources for the development of new drugs. The new stilbene glucosides gnetonosides F **139**, G **140**, H **141**, I **142**, and J **143** were isolated from the liana stems of *Gnetum gnetonoides* and *Gnetum africanum*, along with nine known stilbenoids (161). Other stilbene derivatives, gnetonols K and L (resveratrol trimers), M (isorhapontigenin dimer), and gnetonoside K (glucoside of resveratrol trimer, *cis*-**144** and *trans*-**145** isomers), together with known stilbenoids and a lignan, were isolated from the methanol-soluble parts of the root of *Gnetum gneton* (Gnetaceae). The antioxidant activity of the nonglycoside stilbenoids on lipid peroxide inhibition and superoxide scavenging activity were reported previously (162).



Two dimeric stilbene glucosides, **146** and **147**, were isolated from the root of *Polygonum cuspidatum* (163). One of these glycosides **147**, possesses a novel four-membered ring. Both compounds exhibited strong inhibition of lipid peroxidation but showed no cytotoxic DNA-cleavage activities and no inhibition of protein tyrosine phosphatase 1B.

New glycosides of the stilbene trimers foeniculosides I **148**, II **149**, III **150**, and IV **151** were isolated from *Foeniculum fructus* (seed of *F. vulgare*), along with the known stilbene trimers miyabenol C and *cis*-miyabenol C (164).

The rare, naturally occurring stilbene glucoside sulfates **152–161** were isolated from an aquatic extract of the root of *P. cuspidatum* (165).



A lowland forest tree, *Upuna borneensis* (Dipterocarpaceae), is a monotypic genus distributed in Malaysia (Borneo, including Sabah, Sarawak, and Brunei) (166). Four new stilbene glucosides, upunosides A **162**, B **165**, C **166**, and D **167**, were isolated from the stem of *U. borneensis* (Dipterocarpaceae), together with three known glucosides, vaticaside B **163**, vaticaside C **164**, and paucifloroside A **168**. Upunuside A **162** is the first natural instance of a glucoside of a resveratrol pentamer, and its aglycone has a dibenzo-fused bicyclo[5.3.0]octadiene and two dihydrobenzofuran moieties (167).


## PHENYLETHANOID GLYCOSIDES

Phenylethanoid glycosides are a group of water-soluble natural compounds widely distributed in medicinal plants but also found in other plant species (168–170). Structurally, these metabolites are characterized by benzoic acid derivatives and phenylethyl moieties attached to a  $\beta$ -glucopyranose (apiose, galactose, rhamnose, or xylose) through ester and glycosidic linkages, respectively. More recently, phenylethanoid derivatives attached by an ether and/or ester bond to iridoid glucosides have been found. Phenylethanoid glycosides have been isolated from the aerial parts, leaves, stems, fruit, flowers, roots, and bark of plants or the whole plant; they have been found in the plant families Acanthaceae, Alangiaceae, Asteraceae, Berberidaceae, Bignoniaceae, Burseraceae, Crassulaceae, Lamiaceae, Magnoliaceae, Oleaceae, Orobanchaceae, Plantaginaceae, Scrophulariaceae, and Verbenaceae (170).

Phenethyl  $\beta$ -D-glucopyranoside **169** and phenethyl  $\beta$ -D-galactopyranoside **170** are simple among the identified phenylethanoid glycosides. Phenethyl  $\beta$ -D-glucopyranoside **169** was found in the flowers of *Rosa gallica* (171), and **170** was obtained by fermentation of phenethyl alcohol and the  $\beta$ -galactosidase of *E. coli* (172). Calceolarioside A **171** and B **172** were isolated from the aerial parts of *Calceolaria hypericina* (173). Calceolarioside B **172** and martynoside seemed to possess antifungal activity; both had high MIC values except against the *C. albicans* strain (174). Plantainosides A–F **174**–**179** were isolated from an aqueous extract of the whole plant *Plantago asiatica*, together with eight known phenylethanoid glycosides (175), and the phenylethanoid glycosides **174** and **175** showed antioxidant activity. Calceolarioside A **171**, plantainoside A **174**, and forsythoside A **180** were isolated from *Forsythia suspense* (176). Compounds **171** and **174** were isolated from the genus for the first time, and compound **180** showed immune-enhancing activity in mice (176). Scroside D **181** was isolated from the roots of *Picrorhiza scrophulariiflora* (Scrophulariaceae) (177) and also showed antioxidant activity.

Acteoside **182** (also known as kusagin, NSC 603831, TJC 160, and verbascoside) is a very active phenylethanoid glycoside that has been isolated from many plant species (170). Acteoside **182** and isoacteoside **183** (isoverbascoside) were isolated from the southwestern Indian paintbrush, *Castilleja linariaefolia* (178). The extracts displayed *in vivo* activity against murine P-388 lymphocytic leukemia (178), and the pure glycosides acteoside **182** (ED<sub>50</sub> 2.6  $\mu$ g/mL) and isoacteoside **183** (ED<sub>50</sub> 10  $\mu$ g/mL) showed high cytotoxic activity against the murine P-388 cell line. Acteoside **182** and angoroside A **184** showed strong anti-inflammatory activity *in vitro* (179). Acteoside **182** reduced the risk of atherosclerosis, not only by protecting LDL from oxidative modification but also by its free radical-scavenging properties (180); it also inhibited

human leukocyte elastase (181) and lipopolysaccharide-inducible nitric oxide synthase *in vitro* (182). Acteoside **182** inhibited the telomerase activity (12.5–27 mg/mL) of the human gastric adenocarcinoma cell MKN45 (183). Acteoside **182** and isoacteoside **183** showed antiproliferative activities against B16F10 cells [acteoside,  $GI_{50}$  (concentration needed to reduce the growth of treated cells to half that of untreated cells) 8  $\mu$ M; isoacteoside,  $GI_{50}$  8  $\mu$ M] and their methanolysis products (methyl caffeate  $GI_{50}$  26  $\mu$ M; 3,4-dihydroxyphenethyl alcohol  $GI_{50}$  8  $\mu$ M; 3,4-dihydroxyphenethyl glucoside  $GI_{50}$  10  $\mu$ M; desrhamnosyl acteoside  $GI_{50}$  6  $\mu$ M; and desrhamnosyl isoacteoside  $GI_{50}$  6  $\mu$ M), suggesting that the 3,4-dihydroxyphenethyl alcohol group might be more responsible for the activities of acteoside and isoacteoside than the caffeoyl group (184). Acteoside showed no apparent effect on the marked elevation of serum tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), but it partially prevented *in vitro* TNF- $\alpha$ -induced cell death (100 ng/mL) in D-galactoseamine (D-GalN)-sensitized hepatocytes (0.5 mM) at concentrations of 50, 100, and 200  $\mu$ M. These results indicated that D-GalN/lipopolysaccharide (LPS)-induced hepatic apoptosis can be blocked by an exogenous antioxidant, suggesting the involvement of reactive oxygen intermediates in TNF- $\alpha$ -dependent hepatic apoptosis (185). Acteosides also showed an inhibitory effect on histamine- and bradykinin-induced contractions of guinea pig ileum (186). Campneoside I **185** and II **186** were isolated from a butanol extract of *Paulownia tomentosa* stems and showed antibacterial activity against *S. aureus* (SG511, 285, and 503), *Streptococcus pyogenes* (A308 and A77), and *Streptococcus faecium* MD8b (187). The most active compound of the extract was identified as campneoside I **185**, which had an MIC value of 150  $\mu$ g/mL against *Streptococcus* and *Staphylococcus* species. From such antibacterial activity, the methoxy group of campneoside I was postulated to be the essential element for the antibacterial activity (187). The protective effect of campneoside II **185**, isolated from *Cistanche tubulosa*, on the apoptosis of neurons induced by the neurotoxin 1-methyl-4-phenylpyridinium ion (MPP<sup>+</sup>) was reported (188).




A series of biologically active phenylethanoid trisaccharides with  $\alpha$ -L-rhamnose attached to the C-6' of glucose **187–197** have been isolated from some *Mussatia* species (189–192). Mussatiosides I **188**, II **190**, and III **193** showed inhibitory action on ADP-induced rat platelet aggregation (192). The order of activity was: **188** > **190** > **193**. This antiplatelet effect is likely related to the reported inhibition of cAMP-phosphodiesterase.


Two new phenylethanoid glycosides with cytotoxic activity, 1-*O*-3,4-dimethoxy-phenylethyl-4-*O*-3,4-dimethoxy cinnamoyl-6-*O*-cinnamoyl- $\beta$ -D-glucopyranose **198** and 1-*O*-3,4-dimethoxyphenylethyl-4-*O*-3,4-dimethoxy cinnamoyl- $\beta$ -D-glucopyranose **199**, have been isolated from the acetone extract of




*Psidium guajava* seeds (193). The biological assay of an acetone extract and two isolated compounds showed that the extract had moderate inhibitory activity against both Ehrlich ascites carcinoma (EAC) cells and leukemia P-388 (180% and ED<sub>50</sub> 1/4 14:6), whereas the two new isolated compounds **198** and **199** showed high inhibitory activity against EAC (220 and 240%) and low activity against leukemia P-388 (ED<sub>50</sub> 1/4 17:3 and 16:1).



- 187 R = H  
 188 R = Cinnamoyl  
 189 R = Coumaroyl  
 190 R = Di-Me-Caffeoyl  
 191 R = Feruloyl  
 192 R = Coumaroyl  
 193 R = Me-Coumaroyl  
 194 R = Vanilloyl




- 195 R = H, R<sub>1</sub> = Api, R<sub>2</sub> = Cf  
 196 R = Me, R<sub>1</sub> = Api, R<sub>2</sub> = Fr  
 197 R = H, R<sub>1</sub> = Glc, R<sub>2</sub> = Cf



- 198 R = Cinnamoyl  
 199 R = H


Types of phenylethanoid glycosides linked with iridoids or secoiridoid glucosides have been isolated from plants. Oleoacetoside **200** was isolated from *Syringa reticulata* (194,195), *Jasminum polyanthum* (196,197), and *Picconia excelsa* (198). Oleoechinacoside **201** was isolated from *S. reticulata* (194), and nuzhenide **202** was isolated from the leaves of *S. reticulata* (199). The esterglucoside **203** was found in *Ligustrum japonicum* (200). The unusual polyglucosidic metabolite **204** was isolated from embryos of the American ash (*Fraxinus americana*) (201).



Phenylethanoid derivatives attached by ether and/or ester bonds with iridoid and/or secoiridoid glucosides have been isolated from the Oleaceae family (202). A new secoiridoid glycoside named hirragilide **205** was isolated from the leaves of *Osmanthus ilicifolius* (203). Oleuroside **206** was isolated from *Olea europaea* (204) and from tissues of the novel olive cultivar Hardy's Mammoth (205,206). The antiviral activity of olive leaf extract (OLE), containing oleuroside **206**, hydroxytyrosol, oleuropein, and verbascoside, was reported recently (207). Also, OLE was shown to inhibit acute infection and cell-to-cell transmission of HIV-1, as assayed by syncytia formation using uninfected MT2 cells co-cultured with HIV-1-infected H9 T-lymphocytes. OLE also inhibited HIV-1 replication, as assayed by the expression of p24 in infected H9 cells. These anti-HIV effects of OLE are dose dependent, with an EC<sub>50</sub> of 0.2 µg/mL (207). 10-Acetoxyligstroside **207** and 10-acetoxy-oleuropein **208** were found in an extract of the bark of *Osmanthus asiaticus* (208). Ligustalosite A **209** and ligustalosite B **210** were isolated from the leaves of *Ligustrum lucidum* (Oleaceae) (209) and *Ligustrum vulgare* (210). Kikuchi and Yamauchi (211) found a corresponding pair of compounds, ligstrosidic acid **211** and oleuropeinic acid **212**, in the fruit of *Ligustrum japonica* and *L. lucidum*. Ligstroside **213** and the cytotoxic compound oleuropein **214** were isolated from the stem bark of *Syringa velutina* (212). Oleuropein **214** showed the most potent cytotoxic effect on several tumor cell lines (P-388, L-1210, SNU-5, and HL-60) among eight isolated compounds. Three new secoiridoid glucosides, jasamplexosides A **215**, B **216**, and C (like B, but with the additional part of a secoiridoid glucoside; structure not shown), were isolated from the crude drug *niu du teng*, the leaves and stems of *Jasminum amplexicaule* (213). An unusual cyclic secoiridoid glucoside, insularoside **217**, was isolated



from *Fraxinus insularis* (214), and fraxuhdoside **218** was isolated from the bark of *Fraxinus uhdei* (215).



## PHENYLPROPANOID GLYCOSIDES


A minor iridoid named austromoside **219** was found in *Osmanthus austrocaledonica* (216), and fraxiformoside **220** was isolated from *Fraxinus formosana* (217) and *Fraxinus malacophylla* (218). Framoside **221**, which has anti-inflammatory activity, was isolated from *Fraxinus* species (219); neooleuropein **222** was isolated from *Fraxinus chinensis* (220); and fraximalacoside **223** was isolated from *F. malacophylla* (218). Lilacocide **224** and fliederoside **225** were isolated from *Syringa vulgaris* (221). In 1970, syringopicroside **226** was isolated from *S. vulgaris* (222) and more recently was found in many *Syringa* species (223). A unique series of seven-membered lactones exemplified by jasmolactone C **227** and D **228** were isolated from the aerial part of *Jasminum multiflorum* (224,225). The structures of these compounds, which contain a novel bicyclic 2-oxo-oxepano[4,5-c]pyran ring system, were tested for pharmacological activity, and **227** and **228** were found to possess coronary vasodilating and cardiotropic activities (224).

Phenylpropanoids and phenylethanoid glycosides are a large class of water-soluble natural metabolites widely distributed in plant species (226–231). Phenylpropanoids can conveniently be treated as a group of aromatic compounds consisting of the derivatives of phenylpropanols and/or benzoic acids (caffeic, cinnamic, coumaric, ferulic, sinapic, and others), including their glycosides (see also the Simple Aromatic Glycosides section). More complex phenylpropanoid glycosides are compounds based on phenylethanes (see the Phenylethanoid Glycosides section). Phenylpropanoid compounds have a wide array of important functions in plants. They serve in the interaction of plants with their biotic and abiotic environments, mediate certain aspects of plant growth and development, and are important structural components of the secondary cell wall. Phenylethanoid glycosides have been isolated from the aerial parts of plants, leaves, stems, fruit, flowers, roots, bark, and/or whole plant; they are found in the plant families Araliaceae,

Bignoniaceae, Crassulaceae, Labiatae, Lamiaceae, Oleaceae, Polygonaceae, Scrophulariaceae, Smilacaceae, Verbenaceae, and several others.


The natural phenylpropanoid glycosides isolated from plants showed activities such as being antitumoral, antiviral, anti-inflammatory, antibacterial, antiatherosclerotic, antiplatelet aggregational, antihypertensive, antifatigue, analgesic, hepatoprotective, immunosuppressive, protective of sex and learning behavior, protective of neurodegeneration, reverse transformative of tumor cells, inhibitive of telomerase and capable of shortening the telomere length in tumor cells, having effects on enzymes and cytokines, antioxidative, free radical scavenging, and quickly repairing oxidatively damaged DNA, among others (227).

Simple phenylpropanoid glycosides such as rosin **229**, rosavin **230**, rosarin **231**, triandrin **232**, and vimalin **233** have been isolated from different plant species belonging to the Crassulacea, Liliaceae, and Salicaceae families (226,227). The biologically active substances rosin **229**, rosavin **230**, rosarin **231**, and tyrosol, which are mainly found in plant rhizomes, have demonstrated therapeutic effects (232). These active components affected the central nervous system by increasing the ability to concentrate and mental and physical power; they were efficient in improving asthenic states and the general resistance of the cells and the organism against harmful outer influences. They also prevented the heart system from stress and arrhythmias, and they possessed some antioxidant activities. Some data confirm that the *Rhodiola rosea* preparations stopped the growth of malignant tumors and metastases in the liver (232). Rosin **229**, rosarin **231**, rosavin **231**, and rosiridin isolated from *R. rosea* showed neurotropic activity in mice (233). Citrusin A **234**, B **235**, C **236**, D **237**, E **238**, and F **239** have been isolated from citrus fruit peels, e.g., from the lemon, unshiu, and kinkan (234,235), and after *in vivo* injection of the compounds indicated above into stroke-prone spontaneously hypotensive rats, they were found to lower the blood pressure (236,237).



A new phenylpropanoid glycoside, crenatoside **240**, as well as the known phenylpropanoid, acteoside **182**, have been isolated from the aerial parts of *Orobancha crenata* (238). Two new phenylpropanoid glycosides, junipercomnoside E **241** and F **242**, were isolated from the aerial parts of *Juniperus communis* var.

*depressa* (239); betuloside **243** was isolated from *Betula alba* (240); and cistanoside F **244** was isolated from the stem bark of *Acanthopanax trifoliatum* (241).



## LIGNAN GLYCOSIDES


Lignans are natural bis-phenylpropanoids having a large spectrum of biological activities, and they are widely distributed in the plant kingdom. They are present in small amounts in most unrefined grains such as barley, buckwheat, millet, and oats; legumes such as soybeans; and some vegetables (242–244). Sometimes lignans are referred to as phytoestrogens. However, this term is misleading, as it suggests that lignans may act just like the hormone estrogen in the body. The richest source of lignans is flaxseed, which contains high levels of the lignan precursor secoisolariciresinol diglycoside **245** (SDG). Plant lignans are polyphenolic substances derived from phenylalanine *via* the dimerization of substituted cinnamic alcohols (242). Mammalian lignans are lignans derived from plant lignans. For example, following ingestion, SDG is converted to the aglycone secoisolariciresinol, which is then metabolized to the mammalian lignans enterolactone and enterodiol (245). Most of the effects of oral SDG are mediated by enterolactone and enterodiol (246). SDG has estrogenic and antioxidant activities, and also has antitumor, antiestrogenic, anticarcinogenic, antiatherogenic, and antidiabetic activities (247–249). Similar structures have the glycosides (–)-secoisolariciresinol-*O*- $\alpha$ -L-rhamnopyranoside **93** (105) and (+)-secoisolariciresinol 9-*O*- $\beta$ -D-xylopyranoside **246** (isolated from the leaves of *Laurus nobilis*) (250). The underground parts of *Glehnia littoralis* (Umbelliferae) contain two new lignan glycosides named glehlinosides A **247** and B **248** (251).

Styraxlignolides C **249**, D **250**, E **251**, and F **252**, butyrolactone lignan glucosides with antioxidant activity, were isolated

from the stem bark of *Styrax japonica* (252). Styraxlignolides C **249**, D **250**, and E **251** exhibited weak radical-scavenging activity in the 1,1-diphenyl-2-picrylhydrazyl assay, with IC<sub>50</sub> values of 380, 278, and 194  $\mu$ M, respectively.


Lignans having a diarylhydroxybutyrolactone skeleton, i.e., nortrachelogenin 8'-*O*- $\beta$ -D-glucopyranoside **253**, nortrachelogenin 5'-*C*- $\beta$ -D-glucopyranoside **254**, and nortracheloside **255**, were found in a water-soluble extract of the leaves and stems of *Trachelospermum jasminoides* (253).

The root of *Strobilanthes cusia* (Acanthaceae), popularly known as *da-ching-yeh*, has commonly been used in traditional Chinese medicine. It is used to treat influenza, epidemic cerebrospinal meningitis, encephalitis B, viral pneumonia, mumps, and severe acute respiratory syndrome. A new lignan glycoside, **256**, and the known (+)-9-*O*- $\beta$ -D-glucopyranosyl lyoni-resinol **257** were found in *S. cusia* roots. The isolated compounds **256** and **257** were examined for anti-herpes simplex virus type-1 activity and showed moderate activity (253a).



Bonaspectin C **258** and bonaspectin D **259**, isolated from the aerial parts of *Bonamia spectabilis* (Convolvulaceae), led to the isolation of four minor THF-type sesquilignans (254). Bonaspectin C 4''-*O*-glucoside **258**, its aglycon, and bonaspectin D 4''-*O*-glucoside **259** revealed the highest antiplasmodial activities against *Plasmodium falciparum* [IC<sub>50</sub> values of 0.98 and 5.1  $\mu$ M/mL (chloroquine-sensitive strain of *P. falciparum*), and 1.3 and 2.9  $\mu$ M/mL (chloroquine-resistant clone of *P. falciparum*), respectively]. The cytotoxicity of **258** was estimated by a proliferation assay using the tetrazolium salt 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay against human endothelial cells (ECV-304). A new lignan glucoside, the first 7,9'-monoepoxytetrahydrofuran-type


lignan, named tibeticoside **260**, was isolated from the medicinal plant (roots, stems, and leaves) *Lancea tibetica* (255). Two lignan diglucosides, **261** and **262**, were identified from extracts of germinated sesame seeds (256). The lignan glycoside arctiin **263** is widely distributed in the Compositae family and is used as a chemotaxonomic marker for this family (257). This lignan showed neuroprotective activity against glutamate-induced toxicity in primary cultures of rat cortical cells (258).




The neolignan glucosides tortosides B **264**, D **265**, and E **266** were isolated from *Pedicularis tortu* (259,260). The new lariciresinol-based lignan bis-glucosides, 7*S*,8*R*,8'*R*-(−)-lariciresinol-4,4'-bis-*O*-β-D-glucopyranoside **267** and 7*S*,8*R*,8'*R*-(−)-5-methoxylariciresinol-4,4'-bis-*O*-β-D-glucopyranoside **268** were isolated from the *n*-butanol extract of *Galium sinaicum* roots (261). The two lignan glucosides **267** and **268** were subjected to a cytotoxicity bioassay against the P-388 leukemia cell line, and the tested compounds exhibited weak *in vitro* cytotoxic activity, with IC<sub>50</sub> values of 100 and 42.0 μg/mL, respectively.

The new neolignan glycosides named dichotomosides A **269**, B **270**, C **271**, and D **272** were isolated from a Chinese natural medicine, the roots of *Stellaria dichotoma* var. *lanceolata* (262). Among them, dichotomoside D **272** inhibited the

release of β-hexosaminidase (IC<sub>50</sub> 64 μM) as well as TNF-α and interleukin-4 (IC<sub>50</sub> 16, 34 μM) in RBL-2H3 cells.



A novel lignan glycoside, hibiscuside **273**, which exhibited good antioxidant activity and strongly inhibited lipid peroxidation in rat liver microsomes, was isolated from the roots of *Hibiscus* species (263). Also, a new lignan named eleutheroside E **274** was extracted from the stem bark of *Acanthopanax trifoliatus* (241). A water-soluble natural antioxidant, pinoresinol glycoside **275**, was isolated from sesame seeds (264). Pinoresinol glycoside **275** prevented the oxidation of linoleic acid *in vitro* in an aqueous solution and that of rabbit red cell ghost membranes by *tert*-butyl hydroperoxide (264). A new lignan, triglucoside **276**, was isolated from sesame seeds; this compound had branched (1→2)- and (1→6)-glucosidic linkages and showed antioxidative activity (265). A new furofuran lignan from *Lactuca indica*, lactucaside **277**, showed significant antidiabetic activity (266). *Coptis japonica* (Ranunculaceae) is known to possess several biological activities, such as anti-inflammatory effects. Pinoresinol glycoside **275** and syringaresinol glycoside **278**, which were isolated from the rhizomes of *C. japonica*, were tested to evaluate their *in vitro* anti-inflammatory effects. Pinoresinol glycoside **275** and isolaricresinol showed higher inhibitory effects on TNF- $\alpha$  production than syringaresinol glycoside **278**, whereas syringaresinol glycoside **278** strongly suppressed lymphocyte proliferation (267). For more information on lignans, neolignans, and related compounds, readers are referred to several review articles (242–244,268).






## GLYCOSIDES OF NAPHTHALENE DERIVATIVES

Naphthalene, the fused-ring aromatic hydrocarbon, was first isolated by crystallization from the naphthalene fraction of coal tar, which still remains its major source (269,270). It was also found in combustion processes, including the combustion of refuse, tobacco smoke, coal tar pitch fumes, and oil spills. Most of the naphthalene derivatives found in nature are naphthoquinones, which may arise through the polyketide, terpenoid, or shikimate pathways or a mixture of these (271,272).


The genus *Rumex* (Polygonaceae) is represented by 25 species in the flora of Turkey (273). The roots of *R. patientia* are used as a purgative and tonic in traditional medicine, and its leaves are commonly used as a green vegetable called *labada* or *develik*. Nine *Rumex* species, including *R. patientia*, have been attributed to the Chinese herbal medicine *yangti*, which has been used as a hemostatic and antifungal agent. Two new chlorinated naphthalene derivatives, patientosides A **279** and B **280**, were isolated from the roots of *R. patientia* (274). Daylily flowers are used as an important ingredient in traditional Asian cuisine and are also valued for their reputed medicinal effects. A new naphthalene glycoside, stelladerol **281**, has been isolated from edible daylily (*Hemerocallis*) flowers (275). This isolated compound was tested for antioxidant and cyclooxygenase inhibitory activities and was found to possess strong antioxidant properties, inhibiting lipid oxidation by 94.6% (10  $\mu$ M) in an *in vitro* assay. Also from this plant, three new naphthalene glycosides, rumexoside **282**, rumexoside **283**, and labadoside **284** were identified (276). A new naphthalene glycoside, cassitoroside **285**, was isolated from the seeds of *Cassia tora* (277). One new naphthalene glycoside, 5-hydroxy-dianellin **286**, and a known naphthalene glycoside, dianellin **287**, were isolated from roots of the daylily *Hemerocallis fulva* 'Kwanzo' Kaempfer (278). Diospyronaphthoside **288** was isolated from an ethanolic extract of the stem bark of *Diospyros angustifolia* (279), and plicataloside **289**, an *O,O*-diglycosylated naphthalene derivative, was isolated from *Aloe plicatilis* (280).



The twigs of *Diospyros lycioides*, a plant commonly known as muthala, are frequently used as chewing sticks by the rural and urban people in Namibia for cleaning the teeth. Studies showed that a methanol extract of *D. lycioides* inhibited the growth of selected oral pathogens. Subsequent bioassay-guided

fractionation led to the isolation of four novel bioactive naphthalene glycosides, diospyrosides A **290**, B **291**, C **292**, and D **293** (281). These compounds inhibited the growth of oral cariogenic bacteria (*Streptococcus mutans* and *Streptococcus sanguis*) and periodontal pathogens (*Porphyromonas gingivalis* and *Prevotella intermedia*) at MIC values ranging from 0.019 to 1.25 mg/mL.


Lactonic 6-methoxysorigenin 8-*O*- $\beta$ -D-glucopyranoside **294** was isolated from the cortex of *Rhamnus catharticus* (282), and **295** was isolated from *Rhamnus nakaharai* (283). From the leaves and pods of *Cassia senna* and *C. angustifolia*, the new tinnevellin glucoside **296** was isolated (284). The monoglucoside **297** and diglucoside **298** of 2-naphthalenecarboxylic acid were isolated and characterized from the roots of *Rubia ustulata* (285).



## GLYCOSIDES OF ANTHRACENE DERIVATIVES


Glycosides of anthracene derivatives are widely distributed in the plant kingdom, and many of them have shown anticancer, antiviral, antimicrobial, and other biological activities (286–290). Different anthracene glycosides have been isolated from an aromatic bark called cassia (also known as cinnamon, bastard cinnamon, Chinese cinnamon, *Cassia lignea*, cassia bark, *Cassia aromaticum*, or Canton cassia). Over 400 species of cassia are known. Most are indigenous to North, Central, and South America and Africa, but they are now also found in the tropical and subtropical regions of all continents except Europe. Some are native to South Asia, particularly India and Ceylon, and are now widely cultivated in the tropics; others are considered ornamental trees in southern Florida, the West Indies, and Central and South America. The dried unripe fruit, or Chinese cassia buds, have the odor and taste of the bark and are rather like small cloves in appearance. They have been known as a spice in Europe since the Middle Ages and were probably used in preparing a spiced wine called hippocras. Now they are used in confectionery and in making potpourris.

Emodin is perhaps the most nearly ubiquitous natural anthraquinone, occurring in several higher plants, in fungi, and in lichens. In higher plants, it is chiefly present in glycoconjugates. Emodin and chrysophanol frequently occur together in plants such as *Cassia tora*, *Rumex japonicus*, *Rhamnus purshiana*, *Polygonum multiflorum*, and *P. cuspidatum* (291). Emodins 1-*O*-**299** and 3-*O*-**300** were isolated from the Chinese cassia species (292); 8-*O*- $\beta$ -glucopyranoside **301** was isolated from *C. tora*, *R. japonicus*, *R. purshiana*, *P. multiflorum*, and *P. cuspidatum*; and chrysophanol-8-*O*- $\beta$ -glucoside **302** was found in *R. japonicus* (291). Emodin (IC<sub>50</sub> 3.3  $\mu$ M) is an inhibitor of the growth factor-signaling enzyme phosphatidylinositol-3-kinase (293). A rare sulfated emodin glucoside, emodin 8-*O*- $\beta$ -D-glucopyranosyl-6-*O*-sulfate **303**, was isolated from the roots of the Nepalese medicinal plant *Rheum emodi* (294). Aloe-emodin diglucoside **304** and aloe-emodin  $\omega$ -*O*- $\beta$ -D-glucopyranoside **305** were isolated from Chinese cassia species (292). Emodin 8-*O*- $\beta$ -glucopyranoside **301**, chrysophanol-8-*O*- $\beta$ -glucoside **302**, and rhein-8- $\beta$ -D-glucoside **306**, as well as sennoside A **307**, were isolated from *Rheum palmatum* and *R. tanguticum* (295). Sennoside A **307**, sennoside C **308**, and sennoside E **309** were isolated from *R. palmatum* and *R. tanguticum* species (295).



Cascara sagrada is the common name for *Rhamnus purshiana* (family Rhamnaceae), a short shrubby tree native to the Pacific Northwest in the United States. The dried bark of this tree, also called cascara sagrada, is the source of several anthraquinone laxative drugs, a class of laxatives used by many people throughout the world (296). Dried cascara sagrada bark contains approximately 7–10% hydroxyanthraquinone glycosides (297,298). Hydroxyanthraquinone glycosides are hydroxyanthracene derivatives with hydroxyl groups at the C-1 and C-8 positions and sugar groups at the hydroxyl groups (*O*-glycosides) or at the C-10 position (*C*-glycosides) (296,299). Not less than 60% of this 7–10% consists of cascariosides, expressed as cascarioside A **310**. There are four forms of cascariosides: A **310**, B **312**, C **311**, and D **313**. Cascariosides A **310** and B **312** are diastereoisomers, as are cascariosides C **311** and D **313**. Additional constituents of the dried cascara sagrada bark are the diastereoisomers of barbaloin **314** and **316** and chrysaloin **315** and **317**. Barbaloin stimulates the growth of *Eubacterium* species strain BAR (300). The anthraquinones emodin


**299** and barbaloin **314** showed antiviral activity, but only emodin was a virucidal agent (301). A new cytotoxic 10-epi-veoside, **318**, was isolated from the root bark of *Picramnia antidesma* (302), along with picramnioside E **319** and mayoside B **320**. One new anthraquinone glycoside, *M*-4'-demethylkniphofone-2'-*O*- $\beta$ -D-glucopyranoside **321**, was found in the methanolic extract of the Australian medicinal plant *Bulbine capitata* (303).



Doxorubicin **322**, an anthraquinone glycoside, is currently one of the clinically most important antineoplastic drugs. In pharmacokinetic studies of doxorubicin **322**, capillary blood sampling was recommended, especially for pediatric patients, to avoid the sometimes traumatic venous blood sampling procedure (304). A new anthraquinone glycoside, emodin-6-*O*- $\beta$ -D-glucopyranoside **323**, and the earlier isolated chrysofanol-8-*O*- $\beta$ -glucoside **302**, together with seven known phenolic compounds, have been isolated from the roots of *Rumex patientia*. The cytotoxic effects and radical-scavenging properties of the isolated compounds have been demonstrated (305). The medicinal plant *Bulbine narcissifolia* is used by the Basotho, Griqua, and Whites of southern Africa for wound healing and as a mild purgative. Extraction of the powdered root has yielded acetosyringone, chrysofanol,

knipholone, isoknipholone, 10,7'-bichrysofanol, and chrysalodin in addition to two new anthraquinone glycosides, knipholone-8-*O*- $\beta$ -D-gentiobioside **324** and chrysalodin-10- $\beta$ -D-gentiobioside **325**. NMR spectroscopy was used to elucidate the structures of **324** and **325** and to show that **324** binds weakly to DNA (306).


Frangulin A **326** (also known as emodin-L-rhamnoside, franguloside, or rhamnoxanthin) was discovered in 1907 by Tunman (307) in the bark of *Rhamnus frangula* (alder buckthorn); later, in 1909 the German chemist Schmidt (308), and again in 1913 the Russian scientist Krasovskii (309) found the compound in other *Rhamnus* species. More recently, biologically active frangulin A **326**, frangulin B **327**, glucofrangulin A **328**, and glucofrangulin B **329** have been isolated from *Rhamnus* and *Hypericum* species (310–312). Emodin **299** and frangulin B **326** showed significant antiplatelet effects (313); frangulin B **326** also showed potent inhibitory effects on TNF- $\alpha$  formation in LPS/interferon- $\gamma$ -stimulated murine microglial cell line N9 (314).




An angucycline series compound, P371 A1 **330**, from *Streptomyces* sp. P371, was established as having a novel structure

comprising an ureido group at one of the four sugar units (315). Compound **330** exhibited inhibitory activity against pentagastrin-stimulated acid secretion as well as protective activities against HCl/ethanol- and indomethacin-induced gastric lesions.

The colored urdamycins A **331**, B **332**, C **333**, D **334**, E **335**, and F **336** are six new angucycline antibiotics produced from the *Streptomyces fradiae* strain Tu 2717 (316,317). They are biologically active against gram-positive bacteria and stem cells of murine L1210 leukemia. The urdamycins are glycosides and differ in their aglycones, which can be liberated by acidic hydrolysis in addition to the sugars D-olivose and L-rhodinose.






The antibiotic landomycins A **337**, B **338**, C **339**, D **340**, and E **341** are produced by species of streptomycetes (318–320). Streptomycetes, which are the producers of different polyketide antibiotics, can be divided into four groups based on their sensitivity to landomycins A and E: (i) *Streptomyces glaucescens* Tu49, producer of tetracenomycin, and *Streptomyces aureofaciens* 019, producer of chlortetracycline, belong to the most landomycin-sensitive strains. (ii) *Streptomyces cyanogenus* S136, producer of landomycin A, and *Streptomyces lividans* TK24, producer of actinorhodin, are sensitive to mean and high doses of landomycin E (more than 60–80  $\mu\text{g/mL}$ ). (iii) *Streptomyces globisporus* 1912 (producer of landomycin E), *S. cyanogenus* S136, *S. fradiae* Tu2717 (producer of urdamycins), and *Streptomyces coelicolor* A3(2) (producer of actinorhodin) have moderate sensitivity to all doses of both antibiotics, but strain S136 is sensitive only to low doses. (iv)


Resistance to landomycins A and E was observed in strain *Streptomyces olivaceus* Tu2353, producer of elloramycin. In all these experiments, landomycin E showed lethal activity one to two orders higher than landomycin A, whose molecule is composed of the same aglycon landomycenone A but a longer polysaccharide chain, which probably hampers the penetration of the antibiotic through the cell membrane (319).

A derivative of anthraquinone glycoside, P-1894B **342**, is a potent prolyl hydroxylase inhibitor produced by *Streptomyces albobrisesolus* subspecies no. 1894; it inhibited 50% of the activity of purified chick embryo prolyl hydroxylase at a concentration of  $2.2 \times 10^{-6}$  M (321). The inhibition was noncompetitive with respect to Pro-Pro-Gly, with a  $K_i$  (prolyl hydroxylase activity, expressed in mol/mg protein) of  $1.8 \times 10^{-6}$  M. When excess amounts of  $\text{Fe}^{2+}$  or ascorbate were added to the reaction mixture, the inhibition was slightly reversed. Compound **342**, at a dose of 0.15 mg/kg, reduced the hydroxylation of peptidyl-proline and caused a significant inhibition of collagen biosynthesis in the uterus of the immature rat when stimulated by the administration of 17- $\beta$ -estradiol.

Biological activities, structures, and other information about glycosides of the anthracene derivatives anthraquinones, angucyclines, oxantrones, anthranols, and anthrones can be found in several review articles (322–326).







## SUMMARY

Aromatic natural products have been used for thousands of years, and they are the basis of modern medicines. However, the same yardstick used for modern medicines should not be used to compare natural materials. Modern medicines work very quickly, they are concentrated, they are single components, the results are dramatic, and the risks of side effects are very real. The last couple of decades has seen a tremendous increase in interest in the biological properties of aromatic products as a means of identifying novel small compounds that could have potential in clinical medicine. To that end, stilbenes, lignans, naphthalene-like compounds, and anthracene-like compounds have been of particular interest because of their presence in diet constituents and because of the reported beneficial effects on diverse biological processes and disease conditions. The human diet contains an array of aromatic compounds (phytochemicals) with antioxidant activities capable of coping with reactive oxygen species, thus preventing oxidative stress. Of all the different phenolics, lignans, which are alkylphenolic compounds, constitute the largest class of polyphenols; they have proved to be one of the most powerful antioxidants present in the diet. The latest research has focused on understanding the role played by these dietary metabolites in disease prevention, suggesting a new role for “ancient” food items such as tea, cocoa, grapes (wine), and natural oils as functional foods.

As far as the economic importance of plant aromatic compounds is concerned, different types of medicinal plants are used to treat various ailments pertaining to the nervous system, circulatory system, neuromuscular system, skeletal system, skin, and gastrointestinal system, and are used as immunosuppressives, in oral hygiene, and for the reproductive system. Plant aromatics also find a use as antiviral agents, as metabolic probes, in taxonomy, in genetics, in biotechnology, and in commerce.

## REFERENCES

- Dembitsky, V.M. (2005) Astonishing Diversity of Natural Surfactants. 4. Fatty Acid Amide Glycosides, Their Analogs and Derivatives, *Lipids* 40, 641–660.
- Dey, P.M., and Harborne, J.B. (1997) *Plant Biochemistry*, Academic Press, 554 pp.
- Karakaya, S. (2004) Bioavailability of Phenolic Compounds, *Crit. Rev. Food Sci. Nutr.* 44, 453–464.
- Dell’Agli, M., Busciala, A., and Bosio, E. (2004) Vascular Effects of Wine Polyphenols, *Cardiovasc. Res.* 63, 593–602.
- Kris-Etherton, P.M., Hecker, K.D., Bonanome, A., Coval, S.M., Binkoski, A.E., Hilpert, K.F., Griel, A.E., and Etherton, T.D. (2002) Bioactive Compounds in Foods: Their Role in the Prevention of Cardiovascular Disease and Cancer, *Am. J. Med.* 113 (Suppl. 9B), 71S–88S.
- Cai, Y., Luo, Q., Sun, M., and Corke, H. (2004) Antioxidant Activity and Phenolic Compounds of 112 Traditional Chinese Medicinal Plants Associated with Anticancer, *Life Sci.* 74, 2157–2184.
- Duthie, G., and Crozier, A. (2000) Plant-Derived Phenolic Antioxidants, *Curr. Opin. Clin. Nutr. Metab. Care* 3, 447–451.
- Bravo, L. (1998) Polyphenols: Chemistry, Dietary Sources, Metabolism, and Nutritional Significance, *Nutr. Rev.* 56, 317–333.
- Gubitz, G.M., and Paulo, A.C. (2003) New Substrates for Reliable Enzymes: Enzymatic Modification of Polymers, *Curr. Opin. Biotechnol.* 14, 577–582.
- King, A., and Young, G. (1999) Characteristics and Occurrence of Phenolic Phytochemicals, *J. Am. Diet. Assoc.* 99, 213–218.
- Plomion, C., Leprovost, G., and Stokes, A. (2001) Wood Formation in Trees, *Plant Physiol.* 127, 1513–1523.
- Rabinovich, M.L., Bolobova, A.V., and Vasil’chenko, L.G. (2004) Decomposition of Natural Aromatic Structures and Xenobiotics by Fungi, *Prikl. Biokhim. Mikrobiol.* 40, 5–23.
- Ros Barcelo, A. (1997) Lignification in Plant Cell Walls, *Int. Rev. Cytol.* 176, 87–132.
- Wallace, G., and Fry, S.C. (1994) Phenolic Components of the Plant Cell Wall, *Int. Rev. Cytol.* 151, 229–267.
- Brennan, P.J. (1989) Structure of Mycobacteria: Recent Developments in Defining Cell Wall Carbohydrates and Proteins, *Rev. Infect. Dis.* (Suppl. 2), S420–S430.
- Dembitsky, V.M. (2004) Chemistry and Biodiversity of Biologically Active Natural Glycosides, *Chem. Biodivers.* 1, 673–781.
- Dembitsky, V.M. (2005) Astonishing Diversity of Natural Surfactants. 2. Polyether Glycosidic Ionophores and Macrocyclic Glycosides, *Lipids* 40, 219–248.
- Figuroa-Espinoza, M.C., and Villeneuve, P. (2005) Phenolic Acids Enzymatic Lipophilization, *J. Agric. Food Chem.* 53, 2779–2787.
- Oak, M.H., El Bedoui, J., and Schini-Kerth, V.B. (2005) Antiangiogenic Properties of Natural Polyphenols from Red Wine and Green Tea, *J. Nutr. Biochem.* 16, 1–8.
- De la Lastra, C.A., and Villegas, I. (2005) Resveratrol as an Anti-inflammatory and Anti-aging Agent: Mechanisms and Clinical Implications, *Mol. Nutr. Food Res.* 49, 405–430.
- Pervaiz, S. (2004) Chemotherapeutic Potential of the Chemopreventive Phytoalexin Resveratrol, *Drug Resist. Updat.* 7, 333–344.
- Gescher, A. (2004) Polyphenolic Phytochemicals Versus Nonsteroidal Anti-inflammatory Drugs: Which Are Better Cancer Chemopreventive Agents? *J. Chemother.* (Suppl. 4), 3–6.
- Lambert, J.D., and Yang, C.S. (2003) Cancer Chemopreventive Activity and Bioavailability of Tea and Tea Polyphenols, *Mutat. Res.* 523–524, 201–208.
- Lopez-Velez, M., Martinez-Martinez, F., and Del Valle-Ribes, C. (2003) The Study of Phenolic Compounds as Natural Antioxidants in Wine, *Crit. Rev. Food Sci. Nutr.* 43, 233–244.
- Stark, A.H., and Madar, Z. (2002) Olive Oil as a Functional Food: Epidemiology and Nutritional Approaches, *Nutr. Rev.* 60, 170–176.
- Hollman, P.C., and Katan, M.B. (1999) Health Effects and Bioavailability of Dietary Flavonols, *Free Radic. Res.* 31 (Suppl.), S75–S80.
- Orsini, F., Pelizzoni, F., Bellini, B., and Miglierini, G. (1997) Synthesis of Biologically Active Polyphenolic Glycosides (combreastatin and resveratrol series), *Carbohydr. Res.* 301, 95–109.
- Cheyrier, V. (2005) Polyphenols in Foods Are More Complex Than Often Thought, *Am. J. Clin. Nutr.* 81 (Suppl. 1), 223S–229S.

29. Lee, K.H. (1999) Novel Antitumor Agents from Higher Plants, *Med. Res. Rev.* 19, 569–596.
30. Tan, B.K.H., and Vanitha, J. (2004) Immunomodulatory and Antimicrobial Effects of Some Traditional Chinese Medicinal Herbs: A Review, *Curr. Med. Chem.* 11, 1423–1430.
31. Wojcikowski, K., Johnson, D.W., and Gobe, G. (2004) Medicinal Herbal Extracts—Renal Friend or Foe? Part One: The Toxicities of Medicinal Herbs, *Nephrology* 9, 313–318.
32. Pittler, M.H., and Ernst, E. (2003) Systematic Review: Hepatotoxic Events Associated with Herbal Medicinal Products, *Aliment. Pharmacol. Therapeut.* 18, 451–471.
33. Parojcic, D. (2003) A Historical Overview of the Discovery of Aspirin with Some New Aspects on Its Development, *Arh. Farm. (Belgr.)* 53, 51–70.
34. Weissmann, G. (1991) Aspirin, *Sci. Am.* 264, 84–90.
35. Alstaedter, R. (ed.) (1984) *Aspirin, the Medicine of the Century*, Bayer AG, Leverkusen, Germany.
36. Rainsford, K.D. (1984) *Aspirin and the Salicylates*, Butterworths, London.
37. Gross, M., and Greenburg, L.A. (1948) *The Salicylates: A Critical Bibliographical Review*, Hillhouse, New Haven, CT.
38. Brewer, S. (2000) The Relationship Between Natural Products and Synthetic Chemistry in the Discovery Process, *Spec. Pub. Royal Soc. Chem.* 257, 59–65.
39. Malterud, K.E. (2000) Medicines from Nature, *Kjemi* 60, 10–15.
40. Nahrstedt, A. (1990) Use of Plant Secondary Metabolites by Animals and Humans, *Dtsch. Apoth. Ztg.* 130, 2155–2161.
41. Lukic, P. (1967) Natural Sources of Analgesics, *Arh. Farm. (Belgr.)* 17, 261–270.
42. Bridel, M. (1919) Application of the Biochemical Method to the Branches and Barks of Various Species of the Genus *Populus*, *J. Pharma. Chim. (France)* 19, 429–434.
43. Bridel, M., and Picard, P. (1928) Primeveroside of Salicylic Acid, *Bull. Soc. Chim. Biol.* 10, 381–385.
44. Vane, J.R. (1971) Salicylates, *Nature* 231, 232.
45. Thieme, H. (1963) The Phenolic Glycosides of the Salicaceae. I. General Review (discovery, elucidation of structure, synthesis, occurrence), *Pharmazie* 18, 770–774.
46. Yunes, R.A., Filho, V.C., Ferreira, J., and Calixto, J.B. (2005) The Use of Natural Products as Sources of New Analgesic Drugs, *Stud. Nat. Prod. Chem. Bioact. Nat. Prod.* 30, 191–212.
47. Brubacher, J.R., and Hoffman, R.S. (1996) Salicylism from Topical Salicylates: Review of the Literature, *J. Toxicol. Clin. Toxicol.* 34, 431–436.
48. Smith, M.J.H., and Smith, P.K. (eds.) (1966) *The Salicylates: A Critical Bibliographical Review*, Wiley Interscience, New York, 313 pp.
49. Paris, R.A., and Pointet, M. (1953) Presence of a Monotropitoid in the Bark of *Ostryopsis davidiana*, *Ann. Pharm. Fr.* 11, 346–348.
50. Towers, G.H.N., Tse, A., and Maass, W.S.G. (1966) Phenolic Acids and Phenolic Glycosides of *Gaultheria* Species, *Phytochemistry* 5, 677–681.
51. Boas, F. (1966) *Kwakiutl Ethnobotany*, (Codere, H., ed.) The University of Chicago Press, Chicago.
52. Thieme, H. (1965) Isolation and Structural Clarification of Spiraein, a Phenol Glycoside from *Filipendula ulmaria*, *Pharmazie* 20, 113–114.
53. Meier, B., Lehmann, D., and Sticher, O. (1987) Salicylate in Medicinal Plants. Screening Methods (HPLC, TLC) for Detection, *Dtsch. Apoth. Zeit. (Germany)* 127, 2401–2417.
54. Kanchanapoom, T., Kamel, M.S., Kasai, R., Picheansoonthon, C., Hiraga, Y., and Yamasaki, K. (2001) Benzoxazinoid Glycosides from *Acanthus ilicifolius*, *Phytochemistry* 58, 637–640.
55. Itoh, A., Tanahashi, T., and Nagakura, N. (1993) Two Phenolic Glycosides and an Iridoid Glucoside from *Alangium platani-folium* var. *trilobum*, *Phytochemistry* 33, 161–164.
56. Wei, S., Zhou, Y., and Chang, Y. (1985) Phenolic Glucosides and Phenolic Acids from the Bark of Several Poplar Species, *Linchan Huaxue Yu Gongye* 5, 1–8.
57. Steele, J.W., Weitzel, P.F., and Audette, R.C.S. (1972) Phytochemistry of the Salicaceae. IV. Bark of *Salix petiolaris* (S. *gracilis* var. *textoris*), *J. Chromatogr.* 71, 435–441.
58. Tolonen, A., Pakonen, M., Hohtola, A., and Jalonen, J. (2003) Phenylpropanoid Glycosides from *Rhodiola rosea*, *Chem. Pharm. Bull. (Tokyo)* 51, 467–470.
59. Itoh, A., Tanahashi, T., Nagakura, N., Inoue, K., Kuwajima, H., and Wu, H.X. (2001) Glycosides of Benzyl and Salicyl Alcohols from *Alangium chinense*, *Chem. Pharm. Bull. (Tokyo)* 49, 1343–1345.
60. Delaquis, P.J., Stanich, K., Girard, B., and Mazza, G. (2002) Antimicrobial Activity of Individual and Mixed Fractions of Dill, Cilantro, Coriander and Eucalyptus Essential Oils, *Int. J. Food Microbiol.* 74, 101–109.
61. Singh, G., Kapoor, I.P., Pandey, S.K., Singh, U.K., and Singh, R.K. (2002) Studies on Essential Oils: Part 10: Antibacterial Activity of Volatile Oils of Some Species, *Phytother. Res.* 16, 680–682.
62. Yazdanparast, R., and Alavi, M. (2001) Antihyperlipidaemic and Antihypercholesterolaemic Effects of *Anethum graveolens* Leaves After the Removal of Furocoumarins, *Cytobios* 105, 185–191.
63. Ishikawa, T., Kudo, M., and Kitajima, J. (2002) Water-Soluble Constituents of Dill, *Chem. Pharm. Bull. (Tokyo)* 50, 501–507.
64. Society of Japanese Pharmacopoeia (2001) *Japanese Pharmacopoeia*, 14th edn., pp. 2708–2709, 2869–2871, Hirokawa Publishing, Tokyo.
65. Kitajima, J., Kamoshita, A., Ishikawa, T., Takano, A., Fukuda, T., Isoda, S., and Ida, Y. (2003) Glycosides of *Atractylodes japonica*, *Chem. Pharm. Bull. (Tokyo)* 51, 152–157.
66. Ishikawa, T., Kondo, K., and Kitajima, J. (2003) Water-Soluble Constituents of Coriander, *Chem. Pharm. Bull. (Tokyo)* 51, 32–39.
67. Kitajima, J., and Ishikawa, T. (2003) Water-Soluble Constituents of Amomum Seed, *Chem. Pharm. Bull. (Tokyo)* 51, 890–893.
68. Morikawa, T., Tao, J., Ueda, K., Matsuda, H., and Yoshikawa, M. (2003) Medicinal Foodstuffs. XXXI. 1). Structures of New Aromatic Constituents and Inhibitors of Degranulation in RBL-2H3 Cells from a Japanese Folk Medicine, the Stem Bark of *Acer nikoense*, *Chem. Pharm. Bull. (Tokyo)* 51, 62–67.
69. Ravi, K., Sivagnanam, K., and Subramanian, S. (2004) Antidiabetic Activity of *Eugenia jambolana* Seed Kernels on Streptozotocin-Induced Diabetic Rats, *J. Med. Food* 7, 187–191.
70. Ravi, K., Sekar, D.S., and Subramanian, S. (2004) Hypoglycemic Activity of Inorganic Constituents in *Eugenia jambolana* Seed on Streptozotocin-Induced Diabetes in Rats, *Biol. Trace Elem. Res.* 99, 145–155.
71. Chandrasekaran, M., and Venkatesalu, V. (2004) Antibacterial and Antifungal Activity of *Syzygium jambolanum* Seeds, *J. Ethnopharmacol.* 91, 105–108.
72. Bokesch, H.R., Young, S.M., McKee, T.C., Blunt, J.W., and Boyd, M.R. (1998) Lambertianoside, a Novel Phenylglycoside from *Eugenia lambertiana*, *Nat. Prod. Lett.* 11, 211–216.
73. Hase, K., Kadota, S., Basnet, P., Namba, T., and Takahashi, T. (1996) Hepatoprotective Effects of Traditional Medicines. Isolation of the Active Constituent from Seeds of *Celosia argentea*, *Phytother. Res.* 10, 387–392.
74. Xia, H.C., Li, F., Li, Z., and Zhang, Z.C. (2003) Purification and Characterization of Moschatin, a Novel Type I Ribosome-inactivating Protein from the Mature Seeds of Pumpkin (*Cucurbita moschata*), and Preparation of Its Immunotoxin Against Human Melanoma Cells, *Cell Res.* 13, 369–374.

75. Koike, K., Li, W., Liu, L., Hata, E., and Nikaido, T. (2005) New Phenolic Glycosides from the Seeds of *Cucurbita moschata*, *Chem. Pharm. Bull. (Tokyo)* 53, 225–228.
76. Jung, M.J., Kang, S.S., Jung, Y.J., and Choi, J.S. (2004) Phenolic Glycosides from the Stem Bark of *Albizia julibrissin*, *Chem. Pharm. Bull. (Tokyo)* 52, 1501–1503.
77. De Leo, M., Braca, A., De Tommasi, N., Norscia, I., Morelli, I., Battinelli, L., and Mazzanti, G. (2004) Phenolic Compounds from *Baseonema acuminatum* Leaves: Isolation and Antimicrobial Activity, *Planta Med.* 70, 841–846.
78. Cottiglia, F., Bonsignore, L., Casu, L., Deidda, D., Pompei, R., Casu, M., and Floris, C. (2005) Phenolic Constituents from *Ephedra nebrodensis*, *Nat. Prod. Res.* 19, 117–123.
79. Braham, H., Mighri, Z., Jannet, H.B., Matthew, S., and Abreu, P.M. (2005) Antioxidant Phenolic Glycosides from *Moricandia arvensis*, *J. Nat. Prod.* 68, 517–522.
80. Liu, G.-W. (2001) *Chinese Herbal Medicine*, HuaXia Publishing House, Beijing (in Chinese).
81. Shri, N.G., Laxmi, N.M., and Shantosh, K.A. (1989) Corchioside A, an Orcinol Glycoside from *Curculigo orchioides*, *Phytochemistry* 28, 1771–1772.
82. Xu, J.P., and Xu, R.S. (1992) Phenyl Glycosides from *Curculigo orchioides*, *Yao Xue Xue Bao* 27, 353–357.
83. Li, N., Zhao, Y., Jia, A., Liu, Y., and Zhou, J. (2003) Study on the Chemical Constituents of *Curculigo orchioides*, *Tianran Chanwu Yanjiu Yu Kaifa* 15, 208–211.
84. Lu, H., Zhu, B., and Liang, Y. (2002) Determination of Curculigoside in *Curculigo orchioides* by HPLC, *Zhongguo Zhongyao Zazhi* 27, 192–194.
85. Palazzino, G., Galeffi, C., Federici, E., Delle Monache, F., Francesca Cometa, M., and Palmery, M. (2000) Benzylbenzoate and Norlignan Glucosides from *Curculigo pilosa*: Structural Analysis and *in vitro* Vascular Activity, *Phytochemistry* 55, 411–417.
86. Chen, C., Ni, W., and Mei, W. (1999) Glycosides from *Curculigo orchioides*, *Yunnan Zhi Wu Yan Jiu* 21, 521–524.
87. Fu, D.-X., Lei, G.-Q., Cheng, X.-W., Chen, J.-K., and Zhou, T.-S. (2004) Curculigoside C, a New Phenolic Glucoside from Rhizomes of *Curculigo orchioides*, *Acta Bot. Sin.* 46, 621–624.
88. Wu, H., Liu, Y., and Chen, Y. (1999) Determination of Curculigoside in Rhizomes of *Curculigo orchioides* Gaertn. by HPLC, *Yaowu Fenxi Zazhi* 19, 105–107.
89. Li, N., Tan, N.-H., and Zhou, J. (2004) New Chlorine-Containing Phenoloid from *Curculigo capitulate*, *J. Asian Nat. Prod. Res.* 6, 7–10.
90. Day, J.A., and Saunders, E. (2004) Glycosidation of Chlorophenols by *Lemna minor*, *Environ. Toxicol. Chem.* 23, 613–620.
91. Rothley, D., and Oelschlaeger, H. (1982) Unequivocal Determination of Glucuronides, in Particular of Clofibril  $\beta$ -Glucuronide, *Arch. Pharm. (Weinheim)* 315, 457–462.
92. Miller, L.P. (1957) Monosaccharide Components of an Induced Glycoside of Trichloroethyl Alcohol from Dandelion Tops, *Contrib. Boyce Thompson Inst.* 19, 113–115.
93. Day, J.A., III (2002) Formation and Fate of Chlorophenol Glycosides in an Aquatic Plant Environment, Georgia Institute of Technology, Atlanta, 255 pp. [abstracted in *Diss. Abstr. Int. B* 63, 5435 (2003)].
94. Chang, W.-L., Chen, C.-H., and Lee, S.-S. (1999) Three Novel Constituents from *Curculigo capitulata* and Revision of C-2 Stereochemistry in Nyasicoside, *J. Nat. Prod.* 62, 734–739.
95. Galeffi, C., Federici, E., Palazzino, G., and Nicoletti, M. (1997) Research on African Medicinal Plants. XXXVIII. Further Norlignans from *Hypoxis obtusa* Burch, *Gazz. Chim. Ital.* 127, 501–504.
96. Marini Bettolo, G.B., Patamia, M., Nicoletti, M., Galeffi, C., and Messana, I. (1982) Research on African Medicinal Plants. II. Hypoxoside, a New Glycoside of Uncommon Structure from *Hypoxis obtusa* Busch, *Tetrahedron* 38, 1683–1687.
97. Chang, P.C., and Wu, C.C. (1979) *Herbs of Taiwan in Color*, Vol. 1, Nan Cun (South Villa) Trade Co., Taipei, Republic of China, p. 99.
98. Morikawa, H., Kasai, R., Otsuka, H., Hirata, E., Shinzato, T., Aramoto, M., and Takeda, Y. (2004) Terpenic and Phenolic Glycosides from Leaves of *Breynia officinalis* HEMSL, *Chem. Pharm. Bull. (Tokyo)* 52, 1086–1090.
99. Perry, L.M. (1980) *Medicinal Plants of East and Southeast Asia: Attributed Properties and Uses*, The MIT Press, Cambridge.
100. Kim, J.S., Shim, S.H., Xu, Y.N., Kang, S.S., Son, K.H., Chang, H.W., Kim, H.P., and Bae, K. (2004) Phenolic Glycosides from *Pyrola japonica*, *Chem. Pharm. Bull. (Tokyo)* 52, 714–717.
101. Li, L.M., Liao, X., Peng, S.L., and Ding, L.S. (2005) Chemical Constituents from the Seeds of *Ziziphus jujube* var. *spinosa* (Bunge) Hu, *J. Integrative Plant Biol.* (formerly *Acta Bot. Sin.*) 47, 494–498.
102. Ouyang, M.-A., He, Z.-D., and Wu, C.-L. (2003) Anti-oxidative Activity of Glycosides from *Ligustrum sinense*, *Nat. Prod. Res.* 17, 381–387.
103. Nagatani, Y., Warashina, T., and Noro, T. (2002) Studies on the Constituents from the Aerial Part of *Baccharis dracunculifolia* DC. II, *Chem. Pharm. Bull. (Tokyo)* 50, 583–589.
104. Zhang, W., Chen, W., Wang, Y., Liu, W., Kong, D., and Li, H. (2001) Two New Glycosides from *Erigeron breviscapus* (Vant.) Hand.-Mazz, *Zhongguo Zhongyao Zazhi* 26, 689–690.
105. Chen, X.C., and Jia, Z.J. (2000) Two New Glycosides from *Rubus amabilis*, *Chin. Chem. Lett.* 11, 897–900.
106. Zhang, W., Tam, H.A., Thi, B., Chen, W., Kong, D., Li, H., Wang, Y., and Fouraste, I. (2000) Two New Glycosides from *Erigeron breviscapus*, *J. Chinese Pharmaceut. Sci.* 9, 122–124.
107. Kitajima, J., Ishikawa, T., and Tanaka, Y. (1998) Water-Soluble Constituents of Fennel. II. Four *erythro*-Anethole Glycol Glycosides and Two *p*-Hydroxyphenylpropylene Glycol Glycosides, *Chem. Pharm. Bull. (Tokyo)* 46, 1591–1594.
108. Yoshikawa, K., Eiko, K., Mimura, N., Kondo, Y., and Arihara, S. (1998) Hovetrichosides C–G, Five New Glycosides of Two Auronols, Two Neolignans, and a Phenylpropanoid from the Bark of *Hovenia trichocarea*, *J. Nat. Prod.* 61, 786–790.
109. Rashid, M.A., Gustafson, K.R., Cardellina, J.H., II, and Boyd, M.R. (1996) A Benzoic Acid Glycoside from *Geniostoma antherotrichum*, *Phytochemistry* 41, 1205–1207.
110. Xiao, K., Xuan, L., Xu, Y., and Bai, D. (2002) Novel Stilbene Glycosides from *Polygonum multiflorum*, *Acta Bot. Sin.* 44, 1491–1494.
111. Heitzman, M.E., Neto, C.C., Winiarz, E., Vaisberg, A.J., and Hammond, G.B. (2005) Ethnobotany, Phytochemistry and Pharmacology of *Uncaria* (Rubiaceae), *Phytochemistry* 66, 5–29.
112. Kong, J.M., Goh, N.K., Chia, L.S., and Chia, T.F. (2003) Recent Advances in Traditional Plant Drugs and Orchids, *Acta Pharmacol. Sin.* 24, 7–21.
113. Lu, Y.R., and Foo, L.Y. (2002) Polyphenolics of *Salvia*—A Review, *Phytochemistry* 59, 117–140.
114. Lovkova, M.Ya., Buzuk, G.N., Sokolova, S.M., and Kliment'eva, N.I. (2001) Chemical Features of Medicinal Plants (review), *Appl. Biochem. Microbiol.* 37, 229–237.
115. Agarwal, S.K., Singh, S.S., Lakshmi, V., Vermam, S., and Kumar, S. (2001) Chemistry and Pharmacology of Rhubarb (*Rheum* species)—A Review, *J. Scient. Indust. Res.* 60, 1–9.
116. Von Kruedener, S., Schneider, W., and Elstner, E. (1995) A Combination of *Populus tremula*, *Solidago virgaurea* and *Fraxinus excelsior* as an Antiinflammatory and Antirheumatic Drug—A Short Review, *Arzneim. Forsch./Drug Res.* 45, 169–171.



117. Ahmad, Y.U., and Rahman, A. (1994) *Handbook of Natural Products Data*, Vol. 2, Elsevier, Amsterdam, The Netherlands.
118. Bandyukova, V.A. (1983) Plant Phenol Acids, Their Esters and Glycosides, *Khim. Prirod. Soed. (USSR)* 3, 263–273.
119. Miller, L.P. (1973) Glycosides, *Phytochemistry* 1, 297–375.
120. Hopkinson, S.M. (1969) Chemistry and Biochemistry of Phenolic Glycosides, *Quart. Rev. Chem. Soc.* 23, 98–124.
121. Kompantsev, V.A. (1969) Phenolic Glycosides from *Salix pentandroides*, *Khim. Prirod. Soed. (USSR)* 5, 183–201.
122. Psenak, M. (1967) Phenol Glycosides: Their Significance and Structure, *Biologia (Bratislava)* 22, 704–714.
123. Kondratyuk, T.P., and Pezzuto, J.M. (2004) Natural Product Polyphenols of Relevance to Human Health, *Pharmaceut. Biol.* 42 (Suppl. S.), 46–63.
124. Matsuda, H., Kageura, T., Morikawa, T., Toguchida, I., Harima, S., and Yoshikawa, M. (2000) Effects of Stilbene Constituents from Rhubarb on Nitric Oxide Production in Lipopolysaccharide-Activated Macrophages, *Bioorg. Med. Chem. Lett.* 10, 323–327.
125. Nicolaou, K.C., Roecker, A.J., Barluenga, S., Pfefferkorn, J.A., and Cao, G.-Q. (2001) Discovery of Novel Antibacterial Agents Active Against Methicillin-Resistant *Staphylococcus aureus* from Combinatorial Benzopyran Libraries, *Chem-BioChem* 2, 460–465.
126. Schultz, T.P., Hubbard, T.F., Jr., Jin, L., Fisher, T.H., and Nicholas, D.D. (1990) Role of Stilbenes in the Natural Durability of Wood—Fungicidal Structure—Activity Relationships, *Phytochemistry* 29, 1501–1507.
127. Shen, F., Chen, S.-J., Dong, X.-J., Zhong, H., Li, Y.-T., and Cheng, G.-F. (2003) Suppression of IL-8 Gene Transcription by Resveratrol in Phorbol Ester Treated Human Monocytic Cells, *J. Asian Nat. Prod. Res.* 5, 151–157.
128. Boonlaksiri, C., Oonanant, W., Kongsaree, P., Kittakoop, P., Tanticharoen, M., and Thebtaranonth, Y. (2000) An Antimalarial Stilbene from *Artocarpus integer*, *Phytochemistry* 54, 415–417.
129. Banwell, M.G., Bezos, A., Chand, S., Dannhardt, G., Kiefer, W., Nowe, U., Parish, C.R., Savage, G.P., and Ulbrich, H. (2003) Convergent Synthesis and Preliminary Biological Evaluations of the Stilbenolignan (+/–)-Aiphanol and Various Congeners, *Org. Biomol. Chem.* 1, 2427–2429.
130. Orsini, F., Pelizzoni, F., Verotta, L., Aburjai, T., and Roger, C.B. (1997) Isolation, Synthesis, and Antiplatelet Aggregation Activity of Resveratrol 3-O- $\beta$ -D-Glucopyranoside and Related Compounds, *J. Nat. Prod.* 60, 1082–1087.
131. Pacher, T., Seger, C., Engelmeier, D., Vajrodaya, S., Hofer, O., and Greger, H. (2002) Antifungal Stilbenoids from *Stemona collinsae*, *J. Nat. Prod.* 65, 820–827.
132. Likhitwitayawuid, K., and Sritularak, B. (2001) A New Dimeric Stilbene with Tyrosinase Inhibitory Activity from *Artocarpus gomezianus*, *J. Nat. Prod.* 64, 1457–1459.
133. Dai, J.R., Hallock, Y.F., Cardellina, J.H., and Boyd, M.R. (1998) HIV-Inhibitory and Cytotoxic Oligostilbenes from the Leaves of *Hopea malibato*, *J. Nat. Prod.* 61, 351–353.
134. Hazai, L., and Hornyak, G. (1998) Photochemical Reactions of Stilbenes and Their Heterocyclic Derivatives—A Review, *Ach.-Model. Chem.* 135, 493–514.
135. Hagen, S., and Hopf, H. (1998) Modern Routes to Extended Aromatic Compounds, Carbon Rich Compounds I, *Topic. Curr. Chem.* 196, 45–89.
136. Bai, L., Masukawa, N., Yamaki, M., and Takagi, S. (1997) Two Bibenzyl Glucosides from *Pleione bulbocodioides*, *Phytochemistry* 44, 1565–1567.
137. Toshio, M., and Akira, U. (1991) Ionone and Bibenzyl Glycosides from *Epimedium grandiflorum* var. *thunbergianum*, *Phytochemistry* 30, 1727–1728.
138. Bae, K.H. (2002) *Medicinal Plants of Korea*, Kyo-Hak Publishing Co., Seoul, pp. 302.
139. Yang, H., Sung, S.H., and Kim, Y.C. (2005) Two New Hepatoprotective Stilbene Glycosides from *Acer mono* Leaves, *J. Nat. Prod.* 68, 101–103.
140. Rahman, A., Naz, H., Fadimatou, Makhmoor, T., Yasin, A., Fatima, N., Ngounou, F.N., Kimbu, S.F., Sondengam, B. L., and Choudhary, M.I. (2005) Bioactive Constituents from *Boswellia papyrifera*, *J. Nat. Prod.* 68, 189–193.
141. Roggero, J.P., and Archier, P. (1994) Quantitative Determination of Resveratrol and of One of Its Glycosides in Wines, *Sci. Aliments* 14, 99–107.
142. Vastano, B.C., Chen, Y., Zhu, N., Ho, C.T., Zhou, Z., and Rosen, R.T. (2000) Isolation and Identification of Stilbenes in Two Varieties of *Polygonum cuspidatum*, *J. Agric. Food Chem.* 48, 253–256.
143. Shi, Y.Q., Fukai, T., Sakagami, H., Kuroda, J., Miyaoka, R., Tamura, M., Yoshida, N., and Nomura, T. (2001) Cytotoxic and DNA Damage-Inducing Activities of Low Molecular Weight Phenols from Rhubarb, *Anticancer Res.* 21, 2847–2853.
144. Kimura, Y. (2003) Pharmacological Studies on Resveratrol, *Methods Find. Exp. Clin. Pharmacol.* 25, 297–310.
145. Nonaka, G., Minami, M., and Nishioka, I. (1977) Studies on Rhubarb (*Rhei rhizoma*). III. Stilbene Glycosides, *Chem. Pharm. Bull. (Tokyo)* 25, 2300–2305.
146. Gromova, A.S., Lutskii, V.I., and Tyukavkina, N.A. (1979) Stilbenes from the Bark of Some *Pinaceae* Species, *Koksnes Kimija* 3, 103–109.
147. Pearson, T.W., Kriz, G.S., Jr., and Taylor, R.J. (1977) Absolute Identification of Hydroxystilbenes, Chemical Markers in Engelmann spruce, *Wood Sci.* 10, 93–98.
148. Gromova, A.S., Tyukavkina, N.A., Lutskii, V.I., Kalabin, G.A., and Kushnarev, D.F. (1975) Hydroxystilbenes of the Inner Bark of *Pinus sibirica*, *Khim. Prirod. Soed. (USSR)* 11, 677–682.
149. Bombardelli, E., Martinelli, E.M., and Mustich, G. (1975) Plants of Mozambique. IX. New Hydroxystilbene Glycoside from *Terminalia sericea*, *Fitoterapia* 46, 199–200.
150. Nkiliza, J. (2003) Rhubarb Extract, Compositions Obtained from This Extract, Procedure for Obtaining, and Uses, French Patent: FR 2835185, 20 pp.
151. Kimura, Y., and Okuda, H. (2000) Effects of Naturally Occurring Stilbene Glucosides from Medicinal Plants and Wine, on Tumor Growth and Lung Metastasis in Lewis Lung Carcinoma-Bearing Mice, *J. Pharm. Pharmacol.* 52, 1287–1295.
152. Zhang, L., Li, L., and Li, Y. (2004) Mechanism of the Protection of Effective Component of Tuber Fleeceflower Root Stilbene Glycoside on Nerve Cells, *Zhongguo Linchuang Kangfu* 8, 118–120.
153. Chen, W., Li, L., Yin, X., Yang, G., Qiao, C., Zhang, W., and Chen, H. (2001) Tetrahydroxy-diphenylethylene Glycosides for Preventing and Treating Cardiovascular and Cerebrovascular Diseases, Chinese Patent: CN 1303858, 11 pp.
154. Matsuda, H., Tewtrakul, S., Morikawa, T., and Yoshikawa, M. (2004) Anti-allergic Activity of Stilbenes from Korean Rhubarb (*Rheum undulatum* L.): Structure Requirements for Inhibition of Antigen-Induced Degranulation and Their Effects on the Release of TNF- $\alpha$  and IL-4 in RBL-2H3 Cells, *Bioorg. Med. Chem.* 12, 4871–4876.
155. Pan, H., and Lundgren, L.N. (1995) Phenolic Extractives from Root Bark of *Picea abies*, *Phytochemistry* 39, 1423–1428.
156. Fuendjiep, V., Wandji, J., Tillequin, F., Mulholland, D.A., Budzikiewicz, H., Fomum, Z.T., Nyemba, A.M., and Koch, M. (2002) Chalconoid and Stilbenoid Glycosides from *Guibourtia tessmanii*, *Phytochemistry* 60, 803–806.
157. Ali, Z., Ito, T., Tanaka, T., Nakaya, K.I., Murata, J., Darnaedi, D., and Iinuma, M. (2004) Acetophenone C-Glucosides and Stilbene O-Glucosides in *Upuna borneensis*, *Phytochemistry* 65, 2141–2146.
158. Waffo-Teguo, P., Lee, D., Cuendet, M., Merillon, J.-M., Pezzuto, J.M., and Kinghorn, A.D. (2001) Two New Stilbene

- Dimer Glucosides from Grape (*Vitis vinifera*) Cell Cultures, *J. Nat. Prod.* 64, 136–138.
159. Ito, T., Akao, Y., Tanaka, T., Iinuma, M., and Nozawa, Y. (2002) Vaticanol C, a Novel Resveratrol Tetramer, Inhibits Cell Growth Through Induction of Apoptosis in Colon Cancer Cell Lines, *Biol. Pharm. Bull.* 25, 147–148.
  160. Iliya, I., Tanaka, T., Furusawa, M., Ali, Z., Nakaya, K., Iinuma, M., Shirataki, Y., Murata, J., and Darnaedi, D. (2001) Four New Glucosides of Stilbene Oligomers from the Stem of *Gnetum gnemonoides*, *Heterocycles* 55, 2123–2130.
  161. Iliya, I., Tanaka, T., Iinuma, M., Furusawa, M., Ali, Z., Nakaya, K.I., Murata, J., and Darnaedi, D. (2002) Five Stilbene Glucosides from *Gnetum gnemonoides* and *Gnetum africanum*, *Helv. Chim. Acta* 85, 2394–2402.
  162. Iliya, I., Ali, Z., Tanaka, T., Iinuma, M., Furusawa, M., Nakaya, K., Murata, J., Darnaedi, D., Matsuura, N., and Ubukata, M. (2003) Stilbene Derivatives from *Gnetum gnemon* Linn, *Phytochemistry* 62, 601–606.
  163. Xiao, K., Xuan, L., Xu, Y., Bai, D., Zhong, D., Wu, H., Wang, Z., and Zhang, N. (2002) Dimeric Stilbene Glucosides from *Polygonum cuspidatum*, *Eur. J. Org. Chem.* 3, 564–568.
  164. Ono, M., Ito, Y., Kinjo, J., Yahara, S., Nohara, T., and Niiho, Y. (1995) Four New Glycosides of Stilbene Trimer from *Foeniculi fructus* (fruit of *Foeniculum vulgare* Miller), *Chem. Pharm. Bull. (Tokyo)* 43, 868–871.
  165. Xiao, K., Xuan, L., Xu, Y., and Bai, D. (2000) Stilbene Glycoside Sulfates from *Polygonum cuspidatum*, *J. Nat. Prod.* 63, 1373–1376.
  166. Ashton, P.S. (1982) Flora Malesiana, in *Spermatophyta*, Series 1, Vol. 9 (Steenis C.G.G.J., ed.), Martius Nijhoff Publishers, Leiden, The Netherlands, pp. 337–340.
  167. Ito, T., Ali, Z., Iliya, I., Furusawa, M., Tanaka, T., Nakaya, K.C., Takahashi, Y., Sawa, R., Murata, J., Darnaedi, D., et al. (2005) Occurrence of Stilbene Glucosides in *Upuna borneensis*, *Helv. Chim. Acta* 88, 23–34.
  168. Taskova, R.M., Gotfredsen, C.H., and Jensen, S.R. (2005) Chemotaxonomic Markers in Digitalideae (Plantaginaceae), *Phytochemistry* 66, 1440–1447.
  169. Li, Y., and Ohizumi, Y. (2004) Search for Constituents with Neurotrophic Factor-Potentiating Activity from the Medicinal Plants of Paraguay and Thailand, *Yakugaku Zasshi* 124, 417–424.
  170. Jimenez, C., and Riguera, R. (1994) Phenylethanoid Glycosides in Plants: Structure and Biological Activity, *Nat. Prod. Rep.* 11, 591–606.
  171. Mel'nikov, V.N., Bugorskii, P.S., and Medvedkova, V.V. (1975)  $\beta$ -Phenethyl- $\beta$ -D-glucopyranoside from the Flowers of *Rosa gallica*, *Khim. Prirod. Soed. (USSR)* 11, 807–809.
  172. Kimura, T., Utida, M., Nakajima, H., and Dombou, M. (2000) Enzymatic Synthesis of Phenethyl Glycoside and Its Application, *J. Appl. Glycosci.* 47, 55–59.
  173. Nicoletti, M., Galeffi, C., Messana, I., Garbarino, J.A., Gambaro, V., Nyandat, E., and Marini-Bettolo, G.B. (1986) New Phenylpropanoid Glucosides from *Calceolaria hypericina*, *Gazz. Chim. Ital.* 116, 431–433.
  174. Ersoz, T., Ozalp, M., Ekizoglu, M., and Calis, I. (2002) Antimicrobial Activities of the Phenylethanoid Glycosides from *Scutellaria galericulata*, *Hacettepe Univ. Eczacilik Fak. Derg. (Turkey)* 22, 1–8.
  175. Miyase, T., Ishino, M., Akahori, C., Ueno, A., Ohkawa, Y., and Tanizawa, H. (1991) Phenylethanoid Glycosides from *Plantago asiatica*, *Phytochemistry* 30, 2015–2018.
  176. Liu, D.-L., Zhang, Y., Xu, S.-X., Xu, Y., and Wang, Z.-X. (1998) Phenylethanoid Glycosides from *Forsythia suspensa* Vahl, *J. Chinese Pharmaceut. Sci.* 7, 103–105.
  177. Wang, H., Sun, Y., Ye, W.-C., Xiong, F., Wu, J.-J., Yang, C.-H., and Zhao, S.-X. (2004) Antioxidative Phenylethanoid and Phenolic Glycosides from *Picrorhiza scrophulariiflora*, *Chem. Pharm. Bull. (Tokyo)* 52, 615–617.
  178. Pettit, G.R., Numata, A., Takemura, T., Ode, R.H., Narula, A.S., Schmidt, J.M., Cragg, G.M., and Pase, C.P. (1990) Antineoplastic Agents, 107. Isolation of Acteoside and Isoacteoside from *Castilleja linariaefolia*, *J. Nat. Prod.* 53, 456–458.
  179. Diaz, A.M., Abad, M.J., Fernandez, L., Silvan, A.M., De Santos, J., and Bermejo, P. (2004) Phenylpropanoid Glycosides from *Scrophularia scorodonia*: *In vitro* Anti-inflammatory Activity, *Life Sci.* 74, 2515–2526.
  180. Chiou, W.-F., Lin, L.-C., and Chen, C.-F. (2003) The Antioxidant and Free Radical Scavenging Properties of Acteoside, *Chin. Pharm. J. (Taipei)* 55, 347–353.
  181. Boje, K., Lechtenberg, M., and Nahrstedt, A. (2003) New and Known Iridoid- and Phenylethanoid Glycosides from *Harpagophytum procumbens* and Their *in vitro* Inhibition of Human Leukocyte Elastase, *Planta Med.* 69, 820–825.
  182. Lee, J.Y., Woo, E.-R., and Kang, K.W. (2005) Inhibition of Lipopolysaccharide-Inducible Nitric Oxide Synthase Expression by Acteoside Through Blocking of AP-1 Activation, *J. Ethnopharmacol.* 97, 561–566.
  183. Zhang, F., Jia, Z., Deng, Z., Fan, J., Chen, J., Wu, H., Zhao, C., and Wei, Y. (2002) Inhibition of Verbascoside on the Telomerase Activity of Human Gastric Adenocarcinoma Cell MKN45, *Shijie Huaren Xiaohua Zazhi* 10, 366–367.
  184. Nagao, T., Abe, F., and Okabe, H. (2001) Antiproliferative Constituents in the Plants. 7. Leaves of *Clerodendron bungei* and Bark of *C. trichotomum*, *Biol. Pharm. Bull.* 24, 1338–1341.
  185. Xiong, Q., Hase, K., Tezuka, Y., Namba, T., and Kadota, S. (1999) Acteoside Inhibits Apoptosis in D-Galactosamine and Lipopolysaccharide-Induced Liver Injury, *Life Sci.* 65, 421–430.
  186. Schapoval, E.E.S., Winter de Vargas, M.R., Chaves, C.E.G., Bridi, R., Zuanazzi, J.A., and Henriques, A.T. (1998) Antiinflammatory and Antinociceptive Activities of Extracts and Isolated Compounds from *Stachytarpheta cayennensis*, *J. Ethnopharmacol.* 60, 53–59.
  187. Kang, K.H., Jang, S.K., Kim, B.K., and Park, M.K. (1994) Antibacterial Phenylpropanoid Glycosides from *Paulownia tomentosa* Steud., *Arch. Pharm. Res.* 17, 470–475.
  188. Pu, X., Li, X., Li, H., Tu, P., Song, Z., and Li, C. (2001) Campneoside II of *Cistanche tubulosa* (Schenk) R. Wight Protects Neurons from Apoptosis Induced by Neurotoxin 1-Methyl-4-phenylpyridinium (MPP<sup>+</sup>), *Beijing Daxue Xuebao, Yixueban* 33, 217–220.
  189. Jimenez, C., Villaverde, M.C., Riguera, R., Castedo, L., and Stermitz, F.R. (1987) Three Phenylpropanoid Glycosides from *Mussatia*, *Phytochemistry* 26, 1805–1810.
  190. Jimenez, C., Villaverde, M.C., Riguera, R., Castedo, L., and Stermitz, F.R. (1988) Five Phenylpropanoid Glycosides from *Mussatia*, *Phytochemistry* 27, 2947–2951.
  191. Jimenez, C., Villaverde, M.C., Riguera, R., Castedo, L., and Stermitz, F. (1989) Phenylpropanoid Glycosides from *Mussatia hyacinthine*, *J. Nat. Prod.* 52, 408–410.
  192. Cano, E., Veiga, M., Jimenez, C., and Riguera, R. (1990) Pharmacological Effects of Three Phenylpropanoid Glycosides from *Mussatia*, *Planta Med.* 56, 24–26.
  193. Salib, J.Y., and Michael, H.N. (2004) Cytotoxic Phenylethanoid Glycosides from *Psidium guajava* Seeds, *Phytochemistry* 65, 2091–2093.
  194. Kikuchi, M., and Yamauchi, Y. (1986) Structural Analysis on the Constituents of *Syringa* Species. II. Structures of Components from the Leaves of *Syringa reticulata* (Blume) Hara, *Ann. Rep. Tohoku Coll. Pharm.* 33, 63–68.
  195. Kikuchi, M., Yamauchi, Y., Takahashi, Y., and Sugiyama, M. (1989) Studies on the Constituents of *Syringa* Species. VIII.



- Isolation and Structures of Phenylpropanoid Glycosides from the Leaves of *Syringa reticulata* (Blume) Hara, *Yakugaku Zasshi* 109, 366–371.
196. Shen, Y.-C., Lin, S.-L., Hsieh, P.-W., and Chein, C.-C. (1996) Secoiridoid Glycosides from *Jasminum polyanthum*, *J. Chinese Chem. Soc. (Taipei)* 43, 171–176.
  197. Shen, Y.-C., Lin, S.-L., and Chein, C.-C. (1996) Jaspolyseide, a Secoiridoid Glycoside from *Jasminum polyanthum*, *Phytochemistry* 42, 1629–1631.
  198. Damtoft, S., Franzyk, H., and Jensen, S.R. (1997) Iridoid Glucosides from *Picconia excelsa*, *Phytochemistry* 45, 743–750.
  199. Machida, K., Kaneko, A., Hosogai, T., Kakuda, R., Yaoita, Y., and Kikuchi, M. (2002) Studies on the Constituents of *Syringa* Species. X. Five New Iridoid Glycosides from the Leaves of *Syringa reticulata* (Blume) Hara, *Chem. Pharm. Bull. (Tokyo)* 50, 493–497.
  200. Fukuyama, Y., Koshino, K., Hasegawa, T., Yamada, T., and Nakagawa, K. (1987) New Secoiridoid Glucosides from *Ligustrum japonicum*, *Planta Med.* 53, 427–431.
  201. LaLonde, R.T., Wong, C., and Tsai, A.I.-M. (1976) Polyglucosidic Metabolites of Oleaceae. The Chain Sequence of Oleoside Aglucon, Tyrosol, and Glucose Units in Three Metabolites from *Fraxinus americana*, *J. Am. Chem. Soc.* 98, 3007–3013.
  202. Jensen, S.R., Franzyk, H., and Wallander, E. (2002) Chemotaxonomy of the Oleaceae: Iridoids as Taxonomic Markers, *Phytochemistry* 60, 213–231.
  203. Kikuchi, M., and Yamauchi, Y. (1985) Studies on the Constituents of *Osmanthus* Species. III. On the Components of the Leaves of *Osmanthus ilicifolius* (Hassk.) Mouillefert, *Yakugaku Zasshi* 105, 442–448.
  204. Kuwajima, H., Uemura, T., Takaishi, K., Inoue, K., and Inouye, H. (1988) Monoterpene Glucosides and Related Natural Products. Part 60. A Secoiridoid Glucoside from *Olea europaea*, *Phytochemistry* 27, 1757–1759.
  205. Ryan, D., Antolovich, M., Herlt, T., Prenzler, P.D., Lavee, S., and Robards, K. (2002) Identification of Phenolic Compounds in Tissues of the Novel Olive Cultivar Hardy's Mammoth, *J. Agric. Food Chem.* 50, 6716–6724.
  206. Ryan, D., Prenzler, P.D., Lavee, S., Antolovich, M., and Robards, K. (2003) Quantitative Changes in Phenolic Content During Physiological Development of the Olive (*Olea europaea*) Cultivar Hardy's Mammoth, *J. Agric. Food Chem.* 51, 2532–2538.
  207. Lee-Huang, S., Zhang, L., Huang, P.L., Chang, Y.-T., and Huang, P.L. (2003) Anti-HIV Activity of Olive Leaf Extract (OLE) and Modulation of Host Cell Gene Expression by HIV-1 Infection and OLE Treatment, *Biochem. Biophys. Res. Commun.* 307, 1029–1037.
  208. Sugiyama, M., Machida, K., Matsuda, N., and Kikuchi, M. (1993) Studies on the Constituents of *Osmanthus* Species. Part 15. A Secoiridoid Glycoside from *Osmanthus asiaticus*, *Phytochemistry* 34, 1169–1170.
  209. Kikuchi, M., and Kakuda, R. (1999) Studies on the Constituents of *Ligustrum* species. XIX. Structures of Iridoid Glucosides from the Leaves of *Ligustrum lucidum* AIT, *Yakugaku Zasshi (J. Pharm. Soc. Japan)* 119, 444–450.
  210. Tattini, M., Galardi, C., Pinelli, P., Massai, R., Remorini, D., and Agati, G. (2004) Differential Accumulation of Flavonoids and Hydroxycinnamates in Leaves of *Ligustrum vulgare* Under Excess Light and Drought Stress, *New Phytol.* 163, 547–561.
  211. Kikuchi, M., and Yamauchi, Y. (1985) Studies on the Constituents of *Ligustrum* Species. XI. On the Secoiridoids of the Fruits of *Ligustrum japonica* Thunb. and *L. lucidum* Ait, *Yakugaku Zasshi* 105, 142–147.
  212. Park, H.J., Lee, M.S., Lee, K.T., Sohn, I.C., Han, Y.N., and Miyamoto, K. (1999) Studies on Constituents with Cytotoxic Activity from the Stem Bark of *Syringa velutina*, *Chem. Pharm. Bull. (Tokyo)* 47, 1029–1031.
  213. Tanahashi, T., Shimada, A., Nagakura, N., and Nayeshiro, H. (1992) Jasamplexosides A, B and C: Novel Dimeric and Trimeric Secoiridoid Glucosides from *Jasminum amplexicaule*, *Planta Med.* 58, 552–555.
  214. Tanahashi, T., Shimada, A., Nagakura, N., Inoue, K., Kuwajima, H., Takashi, K., and Chen, C.C. (1993) A Secoiridoid Glucoside from *Fraxinus insularis*, *Phytochemistry* 33, 397–400.
  215. Iossifova, T., Mikhova, B., and Kostova, I. (1995) A Secoiridoid Dilactone from *Fraxinus ornus* Bark, *Monatsh. Chem.* 126, 1257–1264.
  216. Benkrief, R., Ranarivelo, Y., Skaltsounis, A.-L., Tillequin, F., Koch, M., Pusset, J., and Sevenet, T. (1998) Monoterpene Alkaloids, Iridoids and Phenylpropanoid Glycosides from *Osmanthus austrocaledonica*, *Phytochemistry* 47, 825–832.
  217. Tanahashi, T., Watanabe, H., Itoh, A., Nagakura, N., Inoue, K., Ono, M., Fujita, T., and Chen, C.C. (1992) A Secoiridoid Glucoside from *Fraxinus formosana*, *Phytochemistry* 31, 2143–2145.
  218. He, Z.D., Ueda, S., Inoue, K., Akaji, M., Fujita, T., and Yang, C. (1994) Secoiridoid Glucosides from *Fraxinus malacophylla*, *Phytochemistry* 35, 177–181.
  219. Ivanovska, N., Iossifova, T., and Kostova, I. (1996) Complement-Mediated Anti-inflammatory Action of Extracts and Pure Secoiridoids Isolated from *Fraxinus* species, *Phytotherapy Res.* 10, 555–558.
  220. Kuwajima, H., Morita, M., Takaishi, K., Inoue, K., Fujita, T., He, Z.D., and Yang, C.R. (1992) Secoiridoid, Coumarin and Secoiridoid-Coumarin Glucosides from *Fraxinus chinensis*, *Phytochemistry* 31, 1277–1280.
  221. Damtoft, S., Franzyk, H., and Jensen, S.R. (1995) Biosynthesis of Iridoids in *Syringa* and *Fraxinus*: Carbocyclic Iridoid Precursors, *Phytochemistry* 40, 785–792.
  222. Asaka, Y., Kamikawa, T., Tokoroyama, T., and Kubota, T. (1970) The Structure and Absolute Configuration of Syringopicroside. A New Iridoid Glucoside from *Syringa vulgaris* L., *Tetrahedron* 26, 2365–2370.
  223. Kikuchi, M., Yamauchi, Y., and Sugiyama, M. (1989) The Constituents of *Syringa* species. IX. Structure of Oxidative Products from Syringopicroside with Perbenzoic Acid, *Ann. Rep. Tohoku Coll. Pharm.* 36, 97–104.
  224. Shen, Y.C., and Chen, C.H. (1989) Novel Secoiridoid Lactones from *Jasminum multiflorum*, *J. Nat. Prod.* 52, 1060–1070.
  225. Shen, Y.-C., and Chen, C.-H. (1994) A New Secoiridoid Lactone from *Jasminum multiflorum*, *J. Chin. Chem. Soc.* 41, 473–476.
  226. Kurkin, V.A. (2003) Phenylpropanoids from Medicinal Plants: Distribution, Classification, Structural Analysis, and Biological Activity, *Chem. Nat. Comp.* 39, 123–153.
  227. Pan, J., Yuan, C., Lin, C., Jia, Z., and Zheng, R. (2003) Pharmacological Activities and Mechanisms of Natural Phenylpropanoid Glycosides, *Pharmazie* 58, 767–775.
  228. Milkowska-Leyck, K., Borkowski, B., and Strzelecka, H. (2001) Selected Polyphenol Compounds. Part IV. Characteristics and Occurrence of Phenylpropanoid Glycosides, *Herba Polonica* 47, 225–244.
  229. Sawabe, A., Kumamoto, H., and Matsubara, Y. (1998) Bioactive Glycosides in Citrus Fruit Peels, *Kinki Daigaku Nogaku Sogo Kenkyusho Hokoku* 6, 57–67.
  230. Cometa, F., Tomassini, L., Nicoletti, M., and Pieretti, S. (1993) Phenylpropanoid Glycosides. Distribution and Pharmacological Activity, *Fitoterapia* 64, 195–217.
  231. He, Z., and Yang, C. (1989) The Advances of Phenylpropanoid Glycosides in Plants, *Tianran Chanwu Yanjiu Yu Kaifa* 1, 29–41.
  232. Kucinskaite, A., Briedis, V., and Savickas, A. (2004) Experimental Analysis of Therapeutic Properties of *Rhodiola rosea* L. and Its Possible Application in Medicine, *Medicina (Kaunas)* 40, 614–619.

233. Sokolov, S.Ya., Ivashin, V.M., Zapesochnaya, G.G., Kurkin, V.A., and Shchavilinskii, A.N. (1985) Studies of Neurotropic Activity of New Compounds Isolated from *Rhodiola rosea*, *Khim. Farmatsev. Z. (USSR)* 19, 1367–1371.
234. Sawabe, A., Matsubara, Y., Iizuka, Y., and Okamoto, K. (1988) Physiologically Active Substances in Citrus Fruit Peels. XIII. Structure and Physiological Activity of Phenyl Propanoid Glycosides in the Lemon (*Citrus limon* Burm. f.), Unshiu (*Citrus unshiu*), and Kinkan (*Fortunella japonica*), *Nippon Nogeikagaku Kaishi* 62, 1067–1071.
235. Matsubara, Y., Kumamoto, H., Sawabe, A., Iizuka, Y., and Okamoto, K. (1985) Structure and Physiological Activity of Phenyl Propanoid Glycosides in Lemon, Unshiu and Orange Peelings, *Kinki Daigaku Igaku Zasshi* 10, 51–58.
236. Sawabe, A. (1996) Bioactive Compounds in Citrus Fruit Peels, *Foods Food Ingrid. J. Japan* 169, 37–44.
237. Matsubara, Y., Yusa, T., Sawabe, A., Iizuka, Y., and Okamoto, K. (1991) Studies on Physiologically Active Substances in Citrus Fruit Peel. Part XX. Structure and Physiological Activity of Phenyl Propanoid Glycosides in Lemon (*Citrus limon* Burm. f.) Peel, *Agric. Biol. Chem.* 55, 647–650.
238. Afifi, M.S., Lahloub, M.B., El-Khayaat, S.A., Anklin, C.G., Ruegger, H., and Sticher, O. (1993) Crenatoside: A Novel Phenylpropanoid Glycoside from *Orobancha crenata*, *Planta Med.* 59, 359–362.
239. Nakanishi, T., Iida, N., Inatomi, Y., Murata, H., Inada, A., Murata, J., Lang, F.A., Iinuma, M., and Tanaka, T. (2004) New Neolignan and Phenylpropanoid Glycosides in *Juniperus communis* var. *depressa*, *Heterocycles* 63, 2573–2580.
240. Sosa, A. (1934) Biochemical Study of *Betula alba* L. A New Heteroside: Betuloside, and Its Prosthetic Group, Betuligenol, *Rev. Acad. Cienc. (Madrid)* 31, 81–100.
241. Kiem, P.V., Minh, C.V., Dat, N.T., Cai, X.F., Lee, J.J., and Kim, Y.H. (2003) Two New Phenylpropanoid Glycosides from the Stem Bark of *Acanthopanax trifoliatum*, *Arch. Pharm. Res.* 26, 1014–1017.
242. Miyazawa, M. (2001) Biotransformation of Lignans and Neolignans, *Curr. Org. Chem.* 5, 975–986.
243. Ramos, A.C., De Clairac, R.P.L., and Medarde, M. (1999) Heterolignans, *Heterocycles* 51, 1443–1469.
244. Damayanthi, Y., and Lown, J.W. (1998) Podophyllotoxins: Current Status and Recent Developments, *Curr. Med. Chem.* 5, 205–252.
245. Begum, A.N., Nicolle, C., Mila, I., Lapierre, C., Nagano, K., Fukushima, K., Heinonen, S., Adlercreutz, H., Remesy, C., and Scalbert, A. (2004) Dietary Lignins Are Precursors of Mammalian Lignans in Rats, *J. Nutr.* 134, 120–127.
246. Wang, L.Q. (2002) Mammalian Phytoestrogens: Enterodiol and Enterolactone, *J. Chromatogr. B* 777, 289–309.
247. Qu, H.Y., Madl, R.L., Takemoto, D.J., Baybutt, R.C., and Wang, W.Q. (2005) Lignans Are Involved in the Antitumor Activity of Wheat Bran in Colon Cancer SW480 Cells, *J. Nutr.* 135, 598–602.
248. Ranich, T., Bhatena, S.J., and Velasquez, M.T. (2001) Protective Effects of Dietary Phytoestrogens in Chronic Renal Disease, *J. Ren. Nutr.* 11, 183–193.
249. Kitts, D.D., Yuan, Y.V., Wijewickreme, A.N., and Thompson, L.U. (1999) Antioxidant Activity of the Flaxseed Lignan Secoisolariciresinol Diglycoside and Its Mammalian Lignan Metabolites Enterodiol and Enterolactone, *Mol. Cell. Biochem.* 202, 91–100.
250. Yahara, S., Nakazono, M., Tutumi, H., and Nohara, T. (1992) Lignans from Leaves of *Laurus nobilis* L., *Shoyakugaku Zasshi* 46, 184–186.
251. Yuan, Z., Tezuka, Y., Fan, W., Kadota, S., and Li, X. (2002) Constituents of the Underground Parts of *Glehnia littoralis*, *Chem. Pharm. Bull. (Tokyo)* 50, 73–77.
252. Min, B.-S., Na, M.-K., Oh, S.-R., Ahn, K.-S., Jeong, G.-S., Li, G., Lee, S.-K., Joung, H., and Lee, H.-K. (2004) New Furofuran and Butyrolactone Lignans with Antioxidant Activity from the Stem Bark of *Styrax japonica*, *J. Nat. Prod.* 67, 1980–1984.
253. Tan, X.Q., Chen, H.-S., Liu, R.-H., Tan, C.-H., Xu, C.-L., Xuan, W.-D., and Zhang, W.-D. (2005) Lignans from *Trachelospermum jasminoides*, *Planta Med.* 71, 93–95.
- 253a. Tanaka, T., Ikeda, T., Kaku, M., Zhu, X.-H., Okawa, M., Yokomizo, K., Uyeda, M., and Nohara, T. (2004) A New Lignan Glycoside and Phenylethanoid Glycosides from *Strobilanthes cusia* Bremek, *Chem. Pharm. Bull.* 52, 1242–1245.
254. Kraft, C., Jenett-Siems, K., Kohler, I., Tofern-Reblin, B., Siems, K., Bienzle, U., and Eich, E. (2002) Antiplasmodial Activity of Sesquiolignans and Sesqueneolignans from *Bonamia spectabilis*, *Phytochemistry* 60, 167–173.
255. Su, B., Zhu, Q., Gao, K., Yuan, C., and Jia, Z. (1999) Lignan and Phenylpropanoid Glycosides from *Lancea tibetica* and Their Antitumor Activity, *Planta Med.* 65, 558–561.
256. Kuriyama, K., Tsuchiya, K., and Murui, T. (1995) Generation of New Lignan Glucosides During Germination of Sesame Seeds, *Nippon Nogeikagaku Kaishi* 69, 685–693.
257. Haensel, R., Schulz, H., and Leuckert, C. (1964) The Lignane Glycoside Arctiin as a Chemotaxonomic Characteristic in the Compositae Family, *Z. Naturforsch B* 19, 727–734.
258. Jang, Y.P., Kim, S.R., and Kim, Y.C. (2001) Neuroprotective Dibenzylbutyrolactone Lignans of *Torreya nucifera*, *Planta Med.* 67, 470–472.
259. Wang, C.Z., and Jia, Z.J. (1996) Neolignan Glucosides from *Pedicularis torta*, *Chin. Chem. Lett.* 7, 145–148.
260. Wang, C., and Jia, Z. (1997) Lignan, Phenylpropanoid and Iridoid Glycosides from *Pedicularis torta*, *Phytochemistry* 45, 159–166.
261. El Gamal, A.A., Takeya, K., Itokawa, H., Halim, A.F., Amer, M.M., and Saad, H.-E.A. (1997) Lignan Bis-glucosides from *Galium sinaicum*, *Phytochemistry* 45, 597–600.
262. Morikawa, T., Sun, B., Matsuda, H., Wu, L.J., Harima, S., and Yoshikawa, M. (2004) Bioactive Constituents from Chinese Natural Medicines. XIV. New Glycosides of  $\beta$ -Carboline-Type Alkaloid, Neolignan, and Phenylpropanoid from *Stellaria dichotoma* L. var. *lanceolata* and Their Antiallergic Activities, *Chem. Pharm. Bull. (Tokyo)* 52, 1194–1199.
263. Lee, I.G., Lee, S.J., Yoo, I.D., Yoo, I.J., and Yoon, B.S. (2001) Novel Lignan Glycoside Having Inhibition Activity of Lipid Peroxidation, Korean Patent KR 2001002647.
264. Kawagishi, S., Oosawa, T., and Katsuzaki, H. (1994) Pinoresinol Glycoside, Sesame Seed Extract Containing the Glycoside, and Its Use for Preventing Oxidation of Lipids, Japanese Patent JP 06116282, 5 pp.
265. Katsuzaki, H., Kawakishi, S., and Osawa, T. (1993) Structure of Novel Antioxidative Lignan Triglycoside Isolated from Sesame Seed, *Heterocycles* 36, 933–936.
266. Hou, C.C., Lin, S.J., Cheng, J.T., and Hsu, F.L. (2003) Antidiabetic Dimeric Guianolides and a Lignan Glycoside from *Lactuca indica*, *J. Nat. Prod.* 66, 625–629.
267. Cho, J.Y., Kim, A.R., and Park, M.H. (2001) Lignans from the Rhizomes of *Coptis japonica* Differentially Act as Anti-inflammatory Principles, *Planta Med.* 67, 312–316.
268. Ward, R.S. (1999) Lignans, Neolignans and Related Compounds, *Nat. Prod. Rep.* 16, 75–96.
269. Zainullin, R.A., Kukovinets, O.S., Kunakova, R.V., and Tolstikov, G.A. (2001) Naphthalene Oxidation and Reduction Reactions: A Review, *Petroleum Chem. (Russia)* 41, 143–158.
270. Simon, M.A., Bonner, J.S., McDonald, T.J., and Autenrieth, R.L. (1999) Bioaugmentation for the Enhanced Bioremediation of Petroleum in a Wetland, *Polycyc. Aromat. Comp.* 14, 231–239.
271. Bode, H.B., Wegner, B., and Zeeck, A. (2000) Biosynthesis of

- Cladospirone Bisepoxide, a Member of the Spirobisnaphthalene Family, *J. Antibiot.* 53, 153–157.
272. Watanabe, A., and Ebizuka, Y. (2002) A Novel Hexaketide Naphthalene Synthesized by a Chimeric Polyketide Synthase Composed of Fungal Pentaketide and Heptaketide Synthases, *Tetrahedron Lett.* 43, 843–846.
273. Davis, P.H. (1988) *Flora of Turkey and the East Aegean Islands*, University Press, Edinburgh, United Kingdom.
274. Kuruüzüm, A., Demirezer, L.Ö., Bergere, I., and Zeeck, A. (2001) Two New Chlorinated Naphthalene Glycosides from *Rumex patientia*, *J. Nat. Prod.* 64, 688–690.
275. Cichewicz, R.H., and Nair, M.G. (2002) Isolation and Characterization of Stelladerol, a New Antioxidant Naphthalene Glycoside, and Other Antioxidant Glycosides from Edible Daylily (*Heemerocallis*) Flowers, *J. Agric. Food Chem.* 50, 87–91.
276. Demirezer, Ö., Kuruüzüm, A., Bergere, I., Schiewe, H.J., and Zeeck, A. (2001) Five Naphthalene Glycosides from the Roots of *Rumex patientia*, *Phytochemistry* 56, 399–402.
277. Choi, J.S., Jung, J.H., Lee, H.J., Lee, J.H., and Kang, S.S. (1995) A Naphthalene Glycoside from *Cassia tora*, *Phytochemistry* 40, 997–999.
278. Cichewicz, R.H., Lim, K.C., McKerrow, J.H., and Nair, M.G. (2002) Kwanzoquinones A–G and Other Constituents of *Hemerocallis fulva* ‘Kwanzo’ Roots and Their Activity Against the Human Pathogenic Trematode *Schistosoma mansoni*, *Tetrahedron* 58, 8597–8606.
279. Pathak, A., Kulshreshtha, D.K., and Maurya, R. (2004) Coumaroyl Triterpene Lactone, Phenolic and Naphthalene Glycoside from Stem Bark of *Diospyros angustifolia*, *Phytochemistry* 65, 2153–2158.
280. Wessels, P.L., Holzapfel, C.W., van Wyk, B.-E., and Marais, W. (1996) Plicataloside, an *O,O*-Diglycosylated Naphthalene Derivative from *Aloe plicatilis*, *Phytochemistry* 41, 1547–1551.
281. Cai, L., Wei, G.-X., Van der Bijl, P., and Wu, C.D. (2000) Namibian Chewing Stick, *Diospyros lycioides*, Contains Antibacterial Compounds Against Oral Pathogens, *J. Agric. Food Chem.* 48, 909–914.
282. Rauwald, H.W., and Just, H.D. (1983) Study of the Constituents of Buckthorn Cortex. III. A New Lactonic Naphthalene Glycoside from the Cortex of *Rhamnus catharticus* L., *Arch. Pharm. (Weinheim)* 316, 409–412.
283. Lin, C.N., and Wei, B.L. (1994) The Constituents of Formosan *Rhamnus* Species. 8. Flavonol and Naphthalene Glycosides from *Rhamnus nakaharai*, *J. Nat. Prod.* 57, 294–297.
284. Lemli, J., Toppet, S., Cuveele, J., and Janssen, G. (1981) Naphthalene Glycosides in *Cassia senna* and *Cassia angustifolia*. Studies in the Field of Drugs Containing Anthracene Derivatives. XXXII, *Planta Med.* 43, 11–17.
285. Liou, M.-J., Teng, C.-M., and Wu, T.-S. (2002) Constituents from *Rubia ustulata* Diels and *R. yunnanensis* Diels and Their Antiplatelet Aggregation Activity, *J. Chin. Chem. Soc. (Taipei)* 49, 1025–1030.
286. Auvray, P., Bichat, F., and Genne, P. (2000) Preclinical Evaluation of Aromatase Inhibitors Antitumor Activity, *Bull. Cancer* 87, 7–22.
287. Sendelbach, L.E. (1989) A Review of the Toxicity and Carcinogenicity of Anthraquinone Derivatives, *Toxicology* 57, 227–240.
288. Sissi, C., and Palumbo, M. (2004) Antitumor Potential of Azabioisosterism in Anthracenedione-Based Drugs, *Curr. Top. Med. Chem.* 4, 219–230.
289. De Witte, P. (1993) Metabolism and Pharmacokinetics of Anthranoids, *Pharmacology* 47 (Suppl. 1), 86–97.
290. Franz, G. (1993) The Senna Drug and Its Chemistry, *Pharmacology* 47 (Suppl. 1), 2–6.
291. Koyama, J., Morita, I., Kawanishi, K., Tagahara, K., and Kobayashi, N. (2003) Capillary Electrophoresis for Simultaneous Determination of Emodin, Chrysophanol, and Their 8- $\beta$ -D-Glucosides, *Chem. Pharm. Bull. (Tokyo)* 51, 418–420.
292. Yin, J., and Guo, L. (1993) *Modern Study of Chinese Drugs and Clinical Applications*, Xueyuan Press, Beijing, China (in Chinese).
293. Frew, T., Powis, G., Berggren, M., Abraham, R.T., Ashendel, C.L., Zalkow, L.H., Hudson, C., Qazia, S., Gruszecka-Kowalik, E., and Merriman, R. (1994) A Multiwell Assay for Inhibitors of Phosphatidylinositol-3-kinase and the Identification of Natural Product Inhibitors, *Anticancer Res.* 14, 2425–2458.
294. Krenn, L., Presser, A., Pradhan, R., Bahr, B., Paper, D.H., Mayer, K.K., and Kopp, B. (2003) Sulfemodin 8-*O*- $\beta$ -D-Glucoside, a New Sulfated Anthraquinone Glycoside, and Antioxidant Phenolic Compounds from *Rheum emodi*, *J. Nat. Prod.* 66, 1107–1109.
295. Koyama, J., Morita, I., Fujiyoshi, H., and Kobayashi, N. (2005) Simultaneous Determination of Anthraquinones, Their 8- $\beta$ -D-Glucosides, and Sennosides of *Rhei rhizoma* by Capillary Electrophoresis, *Chem. Pharm. Bull. (Tokyo)* 53, 573–575.
296. Van Gorkom, B.A.P., Vries, E.G.E., Karrenbeld, A., and Kleibeuker, J.H. (1999) Review Article: Anthranoid Laxatives and Their Potential Carcinogenic Effects, *Aliment. Pharmacol. Therapeut.* 13, 443–452.
297. Newall, C.A., Anderson, L.A., and Phillipson, J.D. (1996) *Herbal Medicines: A Guide for Health Care Professionals*, The Pharmaceutical Press, London.
298. Wichtl, M., and Bisset, N.G. (eds., English transl. by Bisset, N.G.) (1994) *Rhamni purshiani* Cortex—Cascara Bark, in *Herbal Drugs and Phytopharmaceuticals*, pp. 90–91, CRC Press, Stuttgart.
299. Van Os, F.H. (1976) Anthraquinone Derivatives in Vegetable Laxatives, *Pharmacology* 14 (Suppl. 1), 7–17.
300. Che, Q.M., Akao, T., Hattori, M., Tsuda, Y., Namba, T., and Kobashi, K. (1991) Barbaloin Stimulates Growth of *Eubacterium* sp. Strain BAR, a Barbaloin-Metabolizing Bacterium from Human Feces, *Chem. Pharm. Bull. (Tokyo)* 39, 757–760.
301. Alves, D.S., Perez-Fons, L., Estepa, A., and Micol, V. (2004) Membrane-Related Effects Underlying the Biological Activity of the Anthraquinones Emodin and Barbaloin, *Biochem. Pharmacol.* 68, 549–561.
302. Hernandez-Medel, M.R., and Pereda-Miranda, R. (2002) Cytotoxic Anthraquinone Derivatives from *Picramnia antidesma*, *Planta Med.* 68, 556–558.
303. Qhotsokoane-Lusunzi, M.A., and Karuso, P. (2001) Secondary Metabolites from Basotho Medicinal Plants. II. *Bulbine capitata*, *Aust. J. Chem.* 54, 427–430.
304. Palm, C., Bjork, O., Bjorkholm, M., and Eksborg, S. (2001) Quantification of Doxorubicin in Plasma—A Comparative Study of Capillary and Venous Blood Sampling, *Anticancer Drugs* 12, 859–864.
305. Demirezer, L.O., Kuruuzum-Uz, A., Bergere, I., Schiewe, H.-J., and Zeeck, A. (2001) The Structures of Antioxidant and Cytotoxic Agents from Natural Source: Anthraquinones and Tannins from Roots of *Rumex patientia*, *Phytochemistry* 58, 1213–1217.
306. Qhotsokoane-Lusunzi, M.A., and Karuso, P. (2001) Secondary Metabolites from Basotho Medicinal Plants. I. *Bulbine narcissifolia*, *J. Nat. Prod.* 64, 1368–1372.
307. Tunman, T. (1907) Alder Buckthorn Bark and Its Glucoside, *Chem. Zentralbl. (Germany)* 48, 99–103.
308. Schmidt, E. (1909) Contributions on the Rhamnosides, *Arch. Pharm. (Germany)* 246, 214–224.
309. Krasovskii, N. (1913) Rhamnoxanthin from *Rhamnus cathartica* and Frangulin from *Rhamnus frangula*, *Zh. Russ. Fiziko-Khim, Obshchest. (Russia)* 45, 188–193.
310. Koch, W. (2001) Studies on Biologically Active Substances of *Hypericum* L. Species, *Farmatsevt. Zh. (Kiev)* 4, 50–55.
311. Muhlemann, H. (1955) Anthraquinone and Anthraquinone Glycosides. XVII, *Pharm. Acta Helv.* 30, 350–355.



312. Francis, G.W., Aksnes, D.W., and Holt, O. (1998) Assignment of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR Spectra of Anthraquinone Glycosides from *Rhamnus frangula*, *Magnet. Reson. Chem.* **36**, 769–772.
313. Chung, M.I., Gan, K.H., Lin, C.N., Ko, F.N., and Teng, C.M. (1993) Antiplatelet Effects and Vasorelaxing Action of Some Constituents of Formosan Plants, *J. Nat. Prod.* **56**, 929–934.
314. Wei, B.L., Lu, C.M., Tsao, L.T., Wang, J.P., and Lin, C.N. (2001) *In vitro* Anti-inflammatory Effects of Quercetin 3-*O*-Methyl Ether and Other Constituents from *Rhamnus* Species, *Planta Med.* **67**, 745–747.
315. Uesato, S., Tokunaga, T., and Takeuchi, K. (1998) Novel Angucycline Compound with Both Antigastrin and Gastric Mucosal Protective Activities, *Bioorg. Med. Chem. Lett.* **8**, 1969–1972.
316. Drautz, H., Zahner, H., Rohr, J., and Zeeck, A. (1986) Metabolic Products of Microorganisms. 234. Urdamycins, New Angucycline Antibiotics from *Streptomyces fradiae*. I. Isolation, Characterization and Biological Properties, *J. Antibiot.* **39**, 1657–1669.
317. Rohr, J., and Zeeck, A. (1987) Metabolic Products of Microorganisms. 240. Urdamycins, New Angucycline Antibiotics from *Streptomyces fradiae*. II. Structural Studies of Urdamycins B to F, *J. Antibiot.* **40**, 459–467.
318. Henkel, T., Rohr, J., Beale, J.M., and Schwenen, L. (1990) Landomycins, New Angucycline Antibiotics from *Streptomyces* sp. I. Structural Studies on Landomycins A–D, *J. Antibiot.* **43**, 492–503.
319. Matseliukh, B.P., Konovalova, T.A., Polishchuk, L.V., and Bambura, O.I. (1998) The Sensitivity to Landomycins A and E of Streptomycetes, Producers of Polyketide Antibiotics, *Mikrobiol. Zh. (Kiev)* **60**, 31–36.
320. Liu, H.-W., Sherman, D.H., and Zhao, L. (2002) Recombinant Bacteria Producing Substances with Altered Sugar Moieties and Their Use for Production of These Substances, U.S. Patent Application A2 20020411, 174 pp.
321. Ishimaru, T., Kanamaru, T., Takahashi, T., Ohta, K., and Okazaki, H. (1982) Inhibition of Prolyl Hydroxylase Activity and Collagen Biosynthesis by the Anthraquinone Glycoside, P-1894B, an Inhibitor Produced by *Streptomyces albogriseolus*, *Biochem. Pharmacol.* **31**, 915–919.
322. Bechthold, A., Domann, S., Faust, B., Hoffmeister, D., Stockert, S., Trefzer, A., Weitnauer, G., and Westrich, L. (1999) Glycosylated Natural Products: Perspectives for Combinatorial Biosynthesis, *Chemother. J.* **8**, 130–135.
323. Singh R, Geetanjali, Chauhan SMS. (2004) 9,10-Anthraquinones and Other Biologically Active Compounds from the Genus *Rubia*, *Chem. Biodiver.* **1**, 1241–1264.
324. Krenn, L., Pradhan, R., Presser, A., Reznicek, G., and Kopp, B. (2004) Anthrone C-Glucosides from *Rheum emodi*, *Chem. Pharm. Bull. (Tokyo)* **52**, 391–393.
325. Muhlemann, H. (1949) Anthraquinone and Anthraquinone Glycoside. VI. Anthrone and Anthranol Glycosides, *Pharm. Acta Helv.* **24**, 343–351.
326. Fairbairn, J.W. (1949) Active Constituents of Vegetable Purgatives Containing Anthracene Derivatives. I. Glycosides and Aglycones, *J. Pharm. Pharmacol.* **1**, 683–692.

[Received July 11, 2005; accepted July 26, 2005]