Correction to: Treatment outcomes for acute type A aortic dissection with patent false lumen in patients over the age of 80

Nobuo Kondo¹ · Kentaro Tamura¹ · Arudo Hiraoka¹ · Toshinori Totsugawa¹ · Genta Chikazawa¹ · Taichi Sakaguchi¹ · Hidenori Yoshitaka¹

Published online: 29 February 2020
© The Japanese Association for Thoracic Surgery 2020

Correction to:
General Thoracic and Cardiovascular Surgery
(2019) 67:765–772
<https://doi.org/10.1007/s11748-019-01111-9>

In the original publication of the article, Table 1 was published with errors. The correct Table 1 is given in this correction.

The original article can be found online at <https://doi.org/10.1007/s11748-019-01111-9>.

✉ Nobuo Kondo
kondounobuo0515@gmail.com

¹ Department of Cardiovascular Surgery, The Sakakibara Heart Institute of Okayama, 2-5-1, Nakai-cho, Kita-Ku, Okayama 700-0804, Japan

Table 1 Demographic and characteristics of patients

Variable	Total (n = 67)	Surgical (n = 44)	Medical (n = 23)	p value
Age (years)	85.1 ± 4.7	83.5 ± 2.9	88.3 ± 5.9	< 0.05
Men	15 (22.4)	12 (27.2)	3 (13.0)	0.20
Hypertension	44 (65.7)	29 (65.9)	15 (65.2)	0.45
Diabetes mellitus	5 (7.5)	4 (9.1)	1 (4.3)	0.38
Dislipidemia	14 (20.9)	9 (20.5)	5 (21.7)	0.89
Smoking	12 (17.9)	9 (20.5)	3 (13.0)	0.26
COPD	6 (9.0)	5 (11.4)	1 (4.3)	0.26
Creatinine (mg/dL)	0.96 ± 0.78	1.06 ± 0.92	0.78 ± 0.28	0.86
HD	4 (6.0)	4 (9.1)	0 (0)	0.15
previous cardiac surgery	6 (9.0)	6 (13.6)	0 (0)	0.05
frailty	4.43 ± 1.70	4.09 ± 1.63	5.08 ± 1.70	< 0.05
De Bakey				
Type I	41 (61.2)	30 (68.2)	11 (47.9)	0.10
Type II	12 (17.9)	10 (22.7)	2 (8.7)	0.09
Type IIIR	9 (13.4)	4 (9.1)	5 (21.7)	0.23
Unknown	5 (7.5)	0	5 (21.7)	< 0.05
Patent false part				
Ascending	67 (100)	44 (100)	23 (100)	–
Arch	49	29	20	0.06
Descending	27	16	11	0.36
Abdominal	14	6	8	< 0.05
Complicated dissection				
Brain ischemia	8 (11.9)	7 (15.9)	1 (4.3)	0.12
Coronary ischemia	3 (4.5)	2 (4.5)	1 (4.3)	0.69
Visceral ischemia	2 (2.9)	2 (4.5)	0 (0)	0.39
Limb ischemia	4 (5.9)	3 (6.8)	1 (4.3)	0.52
Aortic valve regurgitation (≥ moderate)	9 (13.3)	9 (20.4)	0 (0)	0.16
Cardiac tamponade	22 (32.8)	16 (36.4)	6 (26.1)	0.54

Continuous data are presented as mean ± SD and categorical data as n (%)

COPD chronic obstructive pulmonary disease, HD hemodialysis