$\delta^{15}\text{N}$ -stable isotope analysis of NH_x : An overview on analytical measurements, source sampling and its source apportionment

Noshan Bhattarai^{1,2}, Shuxiao Wang (✉)^{1,2}, Yuepeng Pan³, Qingcheng Xu^{1,2}, Yanlin Zhang⁴, Yunhua Chang⁴, Yunting Fang⁵

¹ State Key Joint Laboratory of Environmental Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing 100084, China

² State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex, Beijing 100084, China

³ State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

⁴ Yale-NUIST Center on Atmospheric Environment, Nanjing University of Information Science & Technology, Nanjing 210044, China

⁵ CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

HIGHLIGHTS

- Challenges in sampling of NH_3 sources for $\delta^{15}\text{N}$ analysis are highlighted.
- Uncertainties in the isotope-based source apportionment of NH_3 and NH_4^+ are outlined.
- Characterizing dynamic isotopic fractionation may reduce uncertainties of NH_x science.

ARTICLE INFO

Article history:

Received 19 November 2020

Revised 28 January 2021

Accepted 29 January 2021

Available online 15 March 2021

Keywords:

Aerosol ammonium
Atmospheric gaseous ammonia
Isotope fingerprinting
Isotope-based source apportionment
Ammonia gas-to-particle conversion

GRAPHIC ABSTRACT

ABSTRACT

Agricultural sources and non-agricultural emissions contribute to gaseous ammonia (NH_3) that plays a vital role in severe haze formation. Qualitative and quantitative contributions of these sources to ambient $\text{PM}_{2.5}$ (particulate matter with an aerodynamic equivalent diameter below $2.5\ \mu\text{m}$) concentrations remains uncertain. Stable nitrogen isotopic composition ($\delta^{15}\text{N}$) of NH_3 and NH_4^+ ($\delta^{15}\text{N}(\text{NH}_3)$ and $\delta^{15}\text{N}(\text{NH}_4^+)$, respectively) can yield valuable information about its sources and associated processes. This review provides an overview of the recent progress in analytical techniques for $\delta^{15}\text{N}(\text{NH}_3)$ and $\delta^{15}\text{N}(\text{NH}_4^+)$ measurement, sampling of atmospheric NH_3 and NH_4^+ in the ambient air and their sources signature (e.g., agricultural vs. fossil fuel), and isotope-based source apportionment of NH_3 in urban atmosphere. This study highlights that collecting sample that are fully representative of emission sources remains a challenge in fingerprinting $\delta^{15}\text{N}(\text{NH}_3)$ values of NH_3 emission sources. Furthermore, isotopic fractionation during NH_3 gas-to-particle conversion under varying ambient field conditions (e.g., relative humidity, particle pH, temperature) remains unclear, which indicates more field and laboratory studies to validate theoretically predicted isotopic fractionation are required. Thus, this study concludes that lack of refined $\delta^{15}\text{N}(\text{NH}_3)$ fingerprints and full understanding of isotopic fractionation during aerosol formation in a laboratory and field conditions is a limitation for isotope-based source apportionment of NH_3 . More experimental work (in chamber studies) and theoretical estimations in combinations of field verification are necessary in characterizing isotopic fractionation under various environmental and atmospheric neutralization conditions, which would help to better interpret isotopic data and our understanding on NH_x ($\text{NH}_3 + \text{NH}_4^+$) dynamics in the atmosphere.

© Higher Education Press 2021

1 Introduction

Atmospheric ammonia (NH_3) plays a vital role in severe haze formation in the urban (Ye et al., 2011; Behera et al., 2013; Wang et al., 2015; Wei et al., 2015; Wang et al., 2016; Zhao et al., 2016; Ge et al., 2019; Jiang et al., 2021) and rural environment (Meng et al., 2018; Xu et al., 2020; Douglas et al., 2021). Despite the role of NH_3 in ion chemistry of $\text{PM}_{2.5}$ (particulate matter with an aerodynamic equivalent diameter below 2.5 μm), which is due to the considerable contribution from NH_3 gas-to-particle conversion to inorganic secondary aerosol formation (Behera and Sharma, 2012; Behera et al., 2013; Huang et al., 2014), it is a largely unregulated pollutant in many regions of the world including China and the USA. Understanding the role of precursor (e.g., NH_3) in atmospheric chemistry and its contribution to fine particles is the key to designing effective air pollution control policies (e.g., Ding et al., 2019).

The stable nitrogen isotopic composition ($\delta^{15}\text{N}$) of NH_3 (gas) and aerosol ammonium (NH_4^+) in the ambient air ($\delta^{15}\text{N}(\text{NH}_3)$ and $\delta^{15}\text{N}(\text{NH}_4^+)$, respectively) can yield valuable information about its sources and the processes they resulted from beyond what concentration alone could provide (Felix et al., 2014; Felix et al., 2017). For example, Pan et al. (2016) utilized isotope analysis method for source apportionment of NH_4^+ based on predicted initial NH_3 (calculated as initial $\delta^{15}\text{N}(\text{NH}_3) = \delta^{15}\text{N}(\text{NH}_4^+)$ (measured) $- \varepsilon_{\text{NH}_4^+ - \text{NH}_3} \times (1-f)$, where ε is the isotopic enrichment factor and f is the amount of NH_3 converted to NH_4^+ calculated as $(\text{NH}_4^+ / (\text{NH}_4^+ + \text{NH}_3))$ for concurrently measured NH_3 and NH_4^+). Their study and an earlier study from Felix et al. (2014) highlighted the importance of fossil fuel related sources (e.g., vehicular exhaust, NH_3 slip from Selective Catalytic Reduction (SCR) equipped power plants, and coal combustion) based on isotope-based source apportionment of $\delta^{15}\text{N}(\text{NH}_4^+)$ and $\delta^{15}\text{N}(\text{NH}_3)$ values, respectively. Thereafter a number of studies suggested that fossil fuel related sources may be an important source of NH_4^+ and NH_3 in urban atmosphere in China based on source apportionment of $\delta^{15}\text{N}(\text{NH}_4^+)$ values (Liu et al., 2018; Pan et al., 2018a; Wu et al., 2019a; Wu et al., 2019b; Bhattarai et al., 2020; Wu et al., 2020; Xiao et al., 2020; Zhang et al., 2020c) and $\delta^{15}\text{N}(\text{NH}_3)$ values (Chang et al., 2016; Chang et al., 2019; Bhattarai et al., 2020; Zhang et al., 2020a; Zhang et al., 2020b), respectively.

Isotope-based source apportionment of NH_4^+ and NH_3 utilizing $\delta^{15}\text{N}(\text{NH}_4^+)$ values or $\delta^{15}\text{N}(\text{NH}_3)$ values would require relatively distinct source signature of NH_3 emission sources. After early research by Moore (1974), a number of studies (e.g., Freyer, 1978; Felix et al., 2013; Chang et al., 2016) have been carried out on characterizing $\delta^{15}\text{N}(\text{NH}_3)$ source signature of gaseous NH_3 . However, variability and

overlapping exist between previously reported $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for agricultural sources (e.g., livestock waste, fertilizer application), thus this study tries to answer if those source signatures (e.g., livestock waste (−56.1 ‰ to −8.9 ‰) (Freyer, 1978; Heaton, 1987; Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020)) are representative of the original emission sources for different NH_3 collection methods (e.g., active vs. passive). Moreover, this study investigates if previously reported source signatures of NH_3 emission sources are representative for tracer studies.

Source apportionment of NH_4^+ from $\delta^{15}\text{N}(\text{NH}_4^+)$ values requires consideration of equilibrium isotope effect of NH_3 (gas) to NH_4^+ (particle) in the air (Altieri et al., 2014; Pan et al., 2016). Notably, some progresses have been made in the past years, including but not limited to analytical techniques for $\delta^{15}\text{N}(\text{NH}_4^+)/\delta^{15}\text{N}(\text{NH}_3)$ measurement (Zhang et al., 2007; Felix et al., 2013; Liu et al., 2014), characterizing source signature of NH_3 emissions sources (e.g., fertilizer use vs. vehicular emissions reported by Felix et al. (2013)) and theoretical estimation of temperature dependent equilibrium isotopic enrichment factor ($\varepsilon_{\text{NH}_4^+ - \text{NH}_3} (\text{‰}) = 1000 \times (\alpha_{\text{NH}_4^+ - \text{NH}_3} - 1)$, where $\alpha_{\text{NH}_4^+ - \text{NH}_3}$ is the isotopic fractionation factor) for phase specific NH_3 (gas) to NH_4^+ (solid or aqueous) conversion in the atmosphere (Urey, 1947; Walters et al., 2019b). Research by Pan et al. (2016) and Chang et al. (2016) has enabled isotope-based source apportionment of ambient NH_4^+ utilizing $\delta^{15}\text{N}(\text{NH}_4^+)$ values (e.g., Pan et al., 2018a; Wu et al., 2019a), and of ambient NH_3 utilizing $\delta^{15}\text{N}(\text{NH}_3)$ values (e.g., Chang et al., 2019; Zhang et al., 2020a), respectively. However, there are some potential uncertainties in the current approach of isotope-based source apportionment of NH_4^+ and NH_3 , which are highlighted in this study.

In short, we review the recent isotope related progress that has been made in collection of NH_x ($\text{NH}_3 + \text{NH}_4^+$) for $\delta^{15}\text{N}(\text{NH}_4^+)/\delta^{15}\text{N}(\text{NH}_3)$ analysis (second section), isotope source signature of major NH_3 emissions sources (third section), and potential uncertainties in current approach of isotope-based source apportionment of NH_4^+ and NH_3 (fourth section). Outlook for further research needs including suggestions on reducing potential uncertainty are provided in the fifth section.

2 Collection of NH_x for $\delta^{15}\text{N}(\text{NH}_4^+)/\delta^{15}\text{N}(\text{NH}_3)$ analysis

2.1 Collection of NH_x for isotope studies

A summary of advancement in the concentration measurement of NH_3 from the use of the bubbler sampling method to recent online methods can be found in Sutton et al.

(2008). This section only describes recent methods for capturing NH_3 (passive sampling device or active sampling device for simultaneous collection of NH_3 and NH_4^+) for $\delta^{15}\text{N}$ analysis.

Passive diffusion samplers operate on the principle of diffusion (e.g., Adapted Low-cost Passive High Absorption, ALPHA, or Ogawa) and are widely used for measurement of $\delta^{15}\text{N}(\text{NH}_3)$ values in ambient NH_3 (e.g., Felix et al., 2014; Chang et al., 2019) and $\delta^{15}\text{N}$ isotopic fingerprinting of NH_3 emission sources (e.g., Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020).

Replicate samples (e.g., triplicate) need to be collected in parallel with passive samplers to provide estimates of accuracy and precision of method (Tang et al., 2001; Puchalski et al., 2011). Thus, passive samplers can provide confidence in the reproducibility of collected samples. However, recent studies show that $\delta^{15}\text{N}(\text{NH}_3)$ values measured with the ALPHA passive sampler could be lower than actual values based on the comparison of weekly ALPHA sampling with denuder based active sampler (DELTA) (Pan et al., 2020b), and with honeycomb denuders in Walters et al. (2020). In this case, passive samplers may still supplement field sampling of NH_3 measurement for $\delta^{15}\text{N}(\text{NH}_3)$ analysis after correction by adding 15 % for samples collected using passive diffusion samplers (Pan et al., 2020a). Moreover, a recent laboratory-validation study on passive diffusion samplers suggest that bias may vary depending upon the amount of NH_3 collected by the sampler (Kawashima et al., 2021).

Walters and Hastings (2018) and Walters et al. (2019a) showed the suitability of the honeycomb denuder system combined with a filter pack to characterize $\delta^{15}\text{N}(\text{NH}_3)$ and $\delta^{15}\text{N}(\text{NH}_4^+)$ values when the aim is to determine both NH_3 gas and particles in the atmosphere. Honeycomb denuder combined with a filter pack sampling system has been widely used for inorganic gas and particle speciation in previous studies (Koutrakis et al., 1993; Pathak et al., 2003; Wei et al., 2015; Zhao et al., 2016). Walters et al. (2019a) indicates that a Nylon filter with an acid-coated back up filter can be used for collection to determine complete inorganic particulate concentrations for thermodynamic modeling with additional $\delta^{15}\text{N}$ measurement of speciated NH_x , whereas only an acid-coated filter could be used if $\delta^{15}\text{N}(\text{NH}_x)$ measurement were to be the primary goal. A challenge remains in field deployment of active samplers if the site is located far away due to their requirement for electricity.

2.2 Analytical techniques of $\delta^{15}\text{N}(\text{NH}_4^+)/\delta^{15}\text{N}(\text{NH}_3)$

The isotopic analysis of N relies on the generation of a stable gas as the analyte for isotope ratio mass spectrometry (IRMS). For atmospheric NH_3 samples collected using passive diffusion samplers, the sample concentration is often insufficient to satisfy conventional isotope analysis via Elemental Analyzer combustion which analyzes N_2 as an end product and requires greater than 1000 nmol N per sample for $\delta^{15}\text{N}$ analysis. Thus, this review focuses on recent progress in isotope analysis methods for small concentration (10–20 $\mu\text{mol/L}$) and low volume ($< 20\text{ mL}$) atmospheric $\text{NH}_3/\text{NH}_4^+$ samples. Recently, more sensitive isotope analysis methods (requiring only 10–60 nmole of N per analysis) have been developed to determine $\delta^{15}\text{N}(\text{NH}_4^+)$ values for atmosphere, seawaters and soil water samples (Table 1). These methods analyze nitrous oxide (N_2O) as an end product using continuous flow purge and a cryogenically trapped system coupled with IRMS (PT-IRMS) and are: 1) bromate oxidation and azide reduction method (Zhang et al., 2007), 2) coupled bromate oxidation and denitrifier method (Felix et al., 2013), and 3) bromate oxidation and hydroxylamine reduction method (Liu et al., 2014). Briefly, NH_4^+ in the sample are oxidized to nitrite (NO_2^-) using hypobromite (BrO^-) (Zhang et al., 2007), which is then converted to N_2O using either sodium azide (NaN_3) (Method 1) or denitrifying bacteria (Sigman et al., 2001) (Method 2) or hydroxylamine (NH_2OH) (Method 3) (Liu et al., 2014) under strongly acid conditions. The principle of each method can be found in the respective references and so they are not elaborated here.

The ‘isotopic composition’ of a sample relative to a reference standard (atmospheric N_2) are reported using delta notation (δ) in units of per mil (‰) as follows (Eq. (1)):

$$\delta^{15}\text{N}(\text{‰}) = \frac{(^{15}\text{N}/^{14}\text{N})_{\text{sample}} - (^{15}\text{N}/^{14}\text{N})_{\text{standard}}}{(^{15}\text{N}/^{14}\text{N})_{\text{standard}}} \times 1000. \quad (1)$$

Samples should be diluted to optimum NH_4^+ concentration for precise oxidation and reduction yields if NH_4^+ concentration is above the selected experimental method’s working concentration range (Table 1). The dilution is necessary to reduce undesirable isotope fractionation (Zhang et al., 2007). All three methods greatly facilitate

Table 1 Summary of the parameters of $\delta^{15}\text{N}(\text{NH}_3)/\delta^{15}\text{N}(\text{NH}_4^+)$ laboratory experimental methods

No.	Method	NH_4^+ ($\mu\text{mol/L}$)	N_2O (nmole)	Volume (mL)	Standards replicates ($\pm 1\sigma$ ‰)	References
1	Hypobromite-azide	0.5–10	10	20	0.3	Zhang et al.(2007)
2	Coupled denitrifier	10	20	20	0.7	Felix et al. (2013)
3	Hypobromite-hydroxylamine	10–20	60	4	0.3	Liu et al. (2014)

field sampling and in situ measurements due to its superiority in terms of low volume required (< 20 mL) and ability to measure low concentration (~ 10 – 15 $\mu\text{mol/L}$) samples with high throughput and precision. Overall, quantitative chemical transformation method of NH_4^+ to N_2O (Method 3) seems more promising due to their main advantage over use of less dangerous reagent (i.e., NH_2OH) to reduce NO_2^- to N_2O instead of extremely toxic azide buffer reagent (e.g., Hydrazoic acid (HN_3)), and denitrifying bacteria, which may be impractical for most laboratories. When NH_4^+ levels are too low (< 10 $\mu\text{mol/L}$), available methods can be tricky.

3 Isotope source signature of major NH_3 emission sources

3.1 Fossil fuel combustion related sources

$\delta^{15}\text{N}(\text{NH}_3)$ values of fossil fuel combustion-related sources (including emissions from coal-fired power plants and urban traffic) ($0.7\text{‰} \pm 6.5\text{‰}$) and agricultural sources ($-21.9\text{‰} \pm 8.7\text{‰}$) were significantly different (two-sample Welch's *t*-test, $p < 0.00001$) after correction of $\delta^{15}\text{N}(\text{NH}_3)$ values for systematic bias in passive diffusion samplers (Pan et al., 2020b; Walters et al., 2020), allowing its potential use in tracing fossil fuel-related sources and agricultural sources of NH_3 in the atmosphere. The heavier $\delta^{15}\text{N}(\text{NH}_3)$ from fossil fuel sources (-17.8‰ to 9.2‰ (measured) or -16.1‰ to 12.8‰ (after correction)) (Freyer, 1978; Heaton, 1987; Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020; Walters et al., 2020) is potentially due to NH_3 formation during high-temperature combustion and associated fractionation (Felix et al., 2013), compared to lighter $\delta^{15}\text{N}(\text{NH}_3)$ values observed in agricultural sources (-56.1‰ to -8.9‰ or -41.1‰ to -7.8‰ after correction) (Freyer, 1978; Heaton, 1987; Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020) (Fig. 1).

The process that leads to NH_3 emissions from three-way catalytic convertors (TWC) in automobiles and from NH_3 slip from industrial/power plants could likely produce slightly depleted values of $\delta^{15}\text{N}(\text{NH}_3)$ due to fractionation similar to depletion found in NO produced by catalytic reduction (Walters et al., 2015). However, there is a lack of studies assessing potential fractionation in $\delta^{15}\text{N}(\text{NH}_3)$ during catalytic reductions, which means it is not possible to fully explain the mechanism. It is unknown whether or in what ways the reaction pathway and exhaust gas composition affect the $\delta^{15}\text{N}(\text{NH}_3)$ value for NH_3 emitted from automobiles. However, previous studies suggest that the amount of $\delta^{15}\text{N}$ depletion or enrichment could depend on the variations in combustion-related factors (e.g., temperature of the catalyst, catalytic converter efficiency) within each vehicle (Pan et al., 2016; Berner and Felix, 2020).

Importantly, two-sample Welch's *t*-test indicates ($p = 0.37$) and suggests $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for urban traffic samples collected in road tunnels with honeycomb denuders (2.1‰ to 9.2‰) (Walters et al., 2020) falls within the range of $\delta^{15}\text{N}(\text{NH}_3)$ source signature for urban traffic characterized with passive samplers (corrected) (-2.8‰ to 12.8‰) (Felix et al., 2013; Chang et al., 2016). The variability observed in $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for urban traffic (-2.8‰ to 12.8‰) could be likely due to the difference in active and passive collection methods. However, there is still mismatch between $\delta^{15}\text{N}(\text{NH}_3)$ values of tailpipe emissions in field study (-9.3‰ to 9.0‰ , $-1.9\text{‰} \pm 5.5\text{‰}$) (Berner and Felix, 2020) and validated sampling techniques using citric-acid coated honeycomb denuders (2.1‰ to 9.2‰ , $4.2\text{‰} \pm 1.8\text{‰}$) (two-sample Welch's *t*-test, $p < 0.01$) (Walters et al., 2020) (Fig. 1). This suggests more experimental and field studies to ensure representativeness of $\delta^{15}\text{N}(\text{NH}_3)$ source signature for vehicular emissions are required in future studies.

3.2 Agricultural sources

Large variability and overlapping exist between $\delta^{15}\text{N}(\text{NH}_3)$ source signature of livestock waste that ranged from -56.1‰ to -8.9‰ ($-35.4\text{‰} \pm 11.5\text{‰}$) which could be due to different collection methods (e.g., 0.01 M H_2SO_4 scrubber (-12.6‰ to -10.3‰) (Freyer, 1978), HCl scrubber (-15.2‰ and -8.9‰) (Heaton, 1987), and passive diffusion samplers (-56.1‰ to -22.8‰) (Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020)) utilized in NH_3 sampling. Indeed, $\delta^{15}\text{N}(\text{NH}_3)$ source signature for livestock waste (-15.2‰ to -8.9‰) collected using active collection methods (e.g., 0.01 M H_2SO_4 scrubber (Freyer, 1978) and HCl scrubber (Heaton, 1987)) were within the range of $\delta^{15}\text{N}(\text{NH}_3)$ values (-41.1‰ to -7.8‰) collected with passive diffusion samplers (Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020) that were corrected for systematic bias in passive diffusion samplers (Fig. 1). However, $\delta^{15}\text{N}(\text{NH}_3)$ source signature for livestock waste (-15.2‰ to -8.9‰ , $-11.7\text{‰} \pm 2.4\text{‰}$) collected using active collection methods were statistically significantly different from source signature (-41.1‰ to -7.8‰ , $-18.3\text{‰} \pm 7.7\text{‰}$) characterized with passive samplers (corrected) (two-sample Welch's *t*-test, $p < 0.01$).

For fertilizer use, $\delta^{15}\text{N}(\text{NH}_3)$ source signatures ranged from -52.0‰ to -35.0‰ (Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020) where NH_3 were collected using passive diffusion samplers. The variability between $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for fertilizer use (-52.0‰ to -35.0‰) could be due to the difference in urea fertilizer types (Bateman and Kelly, 2007), urea application rates (Ti et al., 2021) and potential equilibrium and kinetic fractionation (Elliott et al., 2019) during volatilization in those field studies (Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020). $\delta^{15}\text{N}(\text{NH}_3)$ source signature for

Fig. 1 Summary of previously reported $\delta^{15}\text{N}$ source signature of NH_3 ($\delta^{15}\text{N}(\text{NH}_3)$) that were collected with passive or active sampling techniques. Dashed horizontal lines classify major sources of NH_3 based on different sample collection methods utilized in previous studies. The $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for sources listed in the figure (from top to bottom) are as following: Marine (-10.2‰ to 2.2‰) (Felix et al., 2013), livestock waste (active) (-15.2‰ to -8.9‰), livestock waste (passive) (-56.1‰ to -22.8‰), fertilizer use (passive) (-52.0‰ to -35.0‰) (Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020), urban waste (passive) including human excreta (-39.6‰ to -37.3‰), solid waste (-37.6‰ to -29.9‰) and waste water (-42.0‰ to -39.2‰) (Chang et al., 2016), coal-fired power plants (CFPP/active) which includes NH_3 slip from SCR equipped CFPP (-16.1‰ to -5.6‰) (Felix et al., 2013; Bhattarai et al., 2020) and coal combustion (-7.2‰ to 2‰) (Freyer, 1978), tailpipe emissions (active) (-9.3‰ to 9.0‰) (Berner and Felix, 2020), urban traffic (tunnel/active) (2.1‰ to 9.2‰) (Walters et al., 2020), and urban traffic (tunnel/passive) (-17.8‰ to -2.2‰) (Felix et al., 2013; Chang et al., 2016; Walters et al., 2020). $\delta^{15}\text{N}(\text{NH}_3)$ values of different sources were corrected for low bias in passive samplers by adding 15‰ to measured $\delta^{15}\text{N}(\text{NH}_3)$ values (Pan et al., 2020b; Walters et al., 2020).

fertilizer use that were corrected for systematic bias in passive diffusion samplers (-37.0‰ to -19.9‰ , $-28.3\text{‰} \pm 5.8\text{‰}$) were significantly different from livestock waste collected using passive diffusion samplers (-41.1‰ to -7.8‰ , $-18.3\text{‰} \pm 7.7\text{‰}$) (two-sample Welch's t -test, $p < 0.00001$) or scrubbers (-15.2‰ to -8.9‰ , $-11.7\text{‰} \pm 2.4\text{‰}$) (two-sample Welch's t -test, $p < 0.00001$) (Fig. 1), allowing its potential use in tracing agricultural sources of NH_3 in the atmosphere.

Volatilization is the driving factor in deriving NH_3 from agricultural sources such as livestock waste and fertilizer application. Figure 2 shows the overlapping of instantaneous $\delta^{15}\text{N}(\text{NH}_3)$ values of liquid manure (-31.0‰ to

-15.0‰) (Hristov et al., 2009) with the field-based $\delta^{15}\text{N}(\text{NH}_3)$ source signature for livestock waste (-41.1‰ to -7.8‰) (Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020), characterized with passive diffusion samplers corrected for systematic bias during the initial stages of incubation (e.g., 14 days). $\delta^{15}\text{N}(\text{NH}_3)$ source signature for livestock waste are more depleted in ^{15}N during initial days of incubation as the lighter isotope molecules tend to react faster when volatilization of $\text{NH}_3\text{-N}$ is most intensive (Hristov et al., 2009; Lee et al., 2011).

Notably, field-based $\delta^{15}\text{N}(\text{NH}_3)$ source signature for livestock waste were more depleted (-41.1‰ to -7.8‰) than the instantaneous $\delta^{15}\text{N}(\text{NH}_3)$ values of liquid manure

Fig. 2 Instantaneous $\delta^{15}\text{N}(\text{NH}_3)$ values in an NH_3 volatilization experiment with liquid manure during the incubation periods (14 to 30 days in total) (Hristov et al., 2009; Lee et al., 2011) and range of $\delta^{15}\text{N}(\text{NH}_3)$ values previously observed during field studies (-56.1‰ to -22.8‰ and -41.1‰ to -7.8‰ (corrected)) (Freyer, 1978; Heaton, 1987; Felix et al., 2013; Chang et al., 2016; Bhattarai et al., 2020) for livestock waste (shaded area). During volatilization experiment, NH_3 released from liquid manure was incubated and absorbed in a $0.5\text{ mol/L H}_2\text{SO}_4$ scrubber to characterize $\delta^{15}\text{N}(\text{NH}_3)$ values in previous studies (Hristov et al., 2009; Lee et al., 2011).

(-31.0‰ (Hristov et al., 2009)/ -22.5‰ (Lee et al., 2011)/ -15.1‰ (Lee et al., 2011)) measured on the first or second day during various incubation experiments (Fig. 2) and, than $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for livestock waste collected with scrubbers (-15.2‰ to -8.9‰) (Freyer, 1978; Heaton, 1987). This discrepancy in $\delta^{15}\text{N}(\text{NH}_3)$ source signatures in previous studies is likely due to difference in sample collection intervals (e.g., (-27.5‰ to -12.1‰) 8–12 h (Chang et al., 2016; Bhattarai et al., 2020)/(-41.1‰ to -7.8‰) two–four weeks (Felix et al., 2013)) because the dynamics of isotopic fractionation could affect $\delta^{15}\text{N}(\text{NH}_3)$ values due to volatilization during longer sampling periods (Högberg, 1997; Lee et al., 2011). Shorter sampling intervals could be appropriate to characterize $\delta^{15}\text{N}(\text{NH}_3)$ values in field conditions for livestock waste with either passive or active collection methods to reduce potential isotope effects. This requires further investigation with concurrent sampling of laboratory validated active sampler and passive sampler, which could additionally provide more understanding on representativeness of livestock waste sample for characterizing $\delta^{15}\text{N}(\text{NH}_3)$ values. This is because the sampling bias between the active collection method and passive sampler may vary depending upon the amount of NH_3 collected by

the sampler (Kawashima et al., 2021), which depends on sampling intervals.

4 Potential uncertainties in current approach of isotope-based source apportionment of NH_x

4.1 Source categories and their isotopic composition values in the stable isotope mixing model

Some studies have assumed three main sources of NH_3 or NH_4^+ (Pan et al., 2016; Pan et al., 2018a; Pan et al., 2018b; Berner and Felix, 2020; Xiao et al., 2020; Zhang et al., 2020c) to resolve measured $\delta^{15}\text{N}(\text{NH}_3)$ value or initial $\delta^{15}\text{N}(\text{NH}_3)$ value derived from $\delta^{15}\text{N}(\text{NH}_4^+)$ value in urban atmosphere. Other studies have assumed four main sources of NH_3 or NH_4^+ (Chang et al., 2016; Liu et al., 2018; Chang et al., 2019; Wu et al., 2019a; Wu et al., 2019b; Bhattarai et al., 2020; Zhang et al., 2020a; Zhang et al., 2020b; Zhang et al., 2020c) in ambient atmosphere (see Supplementary data in Tables A1 and A2).

Isotopic mass balance approach would provide feasible mixing solutions when several sources of NH_3 are assumed

as possible sources and only a single isotope system (e.g., $\delta^{15}\text{N}$ of NH_3) is available (Phillips and Gregg, 2001, 2003). When there are too many sources, the $\delta^{15}\text{N}(\text{NH}_3)$ source signature of sources can be combined if they fall within a similar range and result from similar processes (Phillips et al., 2014). For example, NH_3 emitted from the volatilization process in livestock waste and fertilizer application may have similar $\delta^{15}\text{N}(\text{NH}_3)$ source signatures, so they can be combined. To ensure the qualitative representation of the major NH_3 emission sources, assumed sources and their $\delta^{15}\text{N}(\text{NH}_3)$ source signature should be carefully identified.

It should be noted that mean $\delta^{15}\text{N}(\text{NH}_3)$ source signature of sources may be derived from Keeling plots (Keeling, 1958, 1961), where y-intercept of linear regression between $\delta^{15}\text{N}$ values and $1/\text{concentrations}$ could characterize the mean $\delta^{15}\text{N}$ source signature. However, it is not certain whether mean $\delta^{15}\text{N}$ source signature values derived from Keeling plots (Phillips and Gregg, 2001) or the arithmetic mean of $\delta^{15}\text{N}(\text{NH}_3)$ values as adopted in various studies (Chang et al., 2016; Pan et al., 2016; Stratton et al., 2019; Berner and Felix, 2020) represent the true isotopic composition of NH_3 emissions sources, to be utilized in the isotopic mixing model for source apportionment. This might need further investigation as it has implication for source apportionment studies utilizing isotope-based method. Limited availability of $\delta^{15}\text{N}$ source signature of NH_3 emission sources and their corresponding concentration data may limit wide utilization of Keeling plots in determining the mean of $\delta^{15}\text{N}(\text{NH}_3)$ values for numerous sources.

4.2 Uncertainty from measured/corrected $\delta^{15}\text{N}(\text{NH}_3)$ values collected using passive samplers

Previous studies (Chang et al., 2016; Felix et al., 2017; Chang et al., 2019; Berner and Felix, 2020; Zhang et al., 2020a; Zhang et al., 2020b) have shown that non-agricultural sources such as fossil fuel sources (e.g., vehicle exhaust) and urban waste volatilized sources (human excreta, solid waste and wastewater) are important source of NH_3 in urban areas of China and the USA (Table 2). This is without consideration of potential fractionation in $\delta^{15}\text{N}(\text{NH}_3)$ values characterized for ambient NH_3 samples collected with passive diffusion samplers. Notably, based on recent studies (Pan et al., 2020b; Walters et al., 2020), the contribution of non-agricultural sources (e.g., fossil fuel related sources, urban waste) is likely underestimated with isotope-based source apportionment of NH_3 when $\delta^{15}\text{N}(\text{NH}_3)$ values in ambient NH_3 and $\delta^{15}\text{N}(\text{NH}_3)$ source signatures are corrected for systematic low bias (Bhattarai et al., 2020; Pan et al., 2020a). However, it is not clear if the bias is significantly different in the shorter sampling intervals (e.g., few hours) for various sources when the concentration is high (e.g.,

livestock waste, urban wastes). For example, a recent study suggests that the bias can vary depending on amount of NH_3 collected by the passive diffusion sampler (Kawashima et al., 2021). Thus, $\delta^{15}\text{N}(\text{NH}_3)$ source signatures for some NH_3 emissions sources (e.g., livestock waste, urban wastes) probably need to be more carefully corrected for systematic bias before mean isotopic composition can be determined for tracer studies.

On the other hand, a recent study (Zhang et al., 2020a) suggested that the contribution of non-agricultural sources (e.g., vehicular emissions, NH_3 slip, urban waste) contribution might be underestimated with isotope-based source apportionment of NH_3 if the effect of isotopic fractionation due to $\text{NH}_3(\text{gas})$ to $\text{NH}_4^+(\text{particle})$ conversion on measured $\delta^{15}\text{N}(\text{NH}_3)$ values is not considered. Further investigation is needed to determine how atmospheric processing mechanisms will alter $\delta^{15}\text{N}(\text{NH}_3)$ values under various environmental (e.g., acidic environments) (Elliott et al., 2019) and atmospheric neutralization conditions (e.g., ammonium nitrate (NH_4NO_3) vs. ammonium sulfate ($(\text{NH}_4)_2\text{SO}_4$) during NH_3 -rich or NH_3 -poor conditions) (Moore, 1977; Stratton et al., 2019), because this is unclear.

4.3 Uncertainty in isotope-based source apportionment of particulate NH_4^+

Isotope-based source apportionment of NH_4^+ derived from $\delta^{15}\text{N}(\text{NH}_4^+)$ values requires consideration of isotopic effects to trace back sources of NH_3 (Altieri et al., 2014) under NH_3 -rich conditions. Previous study utilized 33 ‰ (based on Heaton et al. (1997)) as an isotopic fractionation factor to estimate the sources of NH_4^+ from $\delta^{15}\text{N}(\text{NH}_4^+)$ values (Kawashima and Kurahashi, 2011) without consideration of temperature effects on isotopic fractionation factor (Urey, 1947; Savard et al., 2017) and ammonium partitioning ratio ($f = \text{NH}_4^+ / (\text{NH}_4^+ + \text{NH}_3)$) (Pan et al., 2018a). Notably, concurrent measurements of NH_3 and NH_4^+ should be conducted to reduce uncertainty in isotope-based source apportionment of NH_4^+ that may derive from uncertainty in the f value (Pan et al., 2016; Huang et al., 2019).

Previous research accounted for isotope effects due to NH_3 gas-to-particle conversion in predicting initial $\delta^{15}\text{N}(\text{NH}_3)$ derived from $\delta^{15}\text{N}(\text{NH}_4^+)$ values (e.g., Pan et al., 2018a; Wu et al., 2019a; Wu et al., 2020) whereas there could be other sink processes (e.g., NH_3 dry deposition, wet deposition) that could alter $\delta^{15}\text{N}(\text{NH}_3)$ values (Felix et al., 2017; Savard et al., 2017). The mechanism of these sink processes is unclear, which remains a challenge. Thus, there is an opportunity to further investigate the fractionation mechanisms that could alter $\delta^{15}\text{N}(\text{NH}_3)$ values in the atmosphere due to sink processes during various phase transformation of NH_3 and NH_4^+ to increase the scope of related research (Elliott et al., 2019).

Table 2 Summary of recent isotope-based source apportionment studies on gaseous NH_3 derived from measured $\delta^{15}\text{N}(\text{NH}_3)$ values observed in urban areas

Location	Sampling period	Measured $\delta^{15}\text{N}(\text{NH}_3)$ (‰)	Relative contribution (%)		References
			Agricultural ^{a)}	Non-agricultural ^{b)}	
Beijing (China)	18/10/2014 to 29/11/2014	-35.0 ± 5.4 ($n = 12$)	54	46	Chang et al. (2016)
Indianapolis (USA)	07/2009 to 06/2010	-18.0 ($n = 8$)*	10	90	Felix et al. (2017)
Detroit (USA)	07/2009 to 06/2010	-14.3 ($n = 11$)*	4	96**	Felix et al. (2017)
Shanghai (China)	07/2015 to 08/2015	-31.7 ± 3.4 ($n = 10$)	47	53	Chang et al. (2019)
Texas (USA)	09/2016 to 08/2017	-19.1 ± 10.9 ($n = 22$)	45	55***	Berner and Felix (2020)
Beijing (China)	03/2016 to 03/2017	-33.2 ± 8.6 ($n = 73$)	28	72	Zhang et al. (2020a)
Beijing (China)	03/2016 to 03/2017	-33.2 ± 8.6 ($n = 73$)	34	66	Pan et al. (2020a)
Beijing (China)	13/09/2018 to 09/10/2018	-28.9 ± 1.5 ($n = 8$)	37	63	Bhattarai et al. (2020)
Beijing (China)	03/01/2017 to 25/01/2017	—	26	74	Zhang et al. (2020b)
Beijing (China)	07/02/2020 to 13/02/2020 and 25/03/2020 to 01/04/2020	—	38	62	Zhang et al. (2020b)

Notes: a) Including livestock waste and fertilizer application. b) Including urban wastes and fossil fuel related sources. "***" notes that measured $\delta^{15}\text{N}(\text{NH}_3)$ values were compared to relative proportion of emission inventory data. "***" Including contribution from biomass burning and wild fire (1%). "****" Include contribution from urban traffic. "—" Marked no information is available in the cell.

5 Conclusions and suggestions

In this article, we highlighted that the overlapping and large variability in source signatures of different sources (e.g., livestock waste, urban traffic etc.) were likely due to differences between NH_3 collection methods used in previous studies. Notably, discrepancies between $\delta^{15}\text{N}(\text{NH}_3)$ source signatures measured in the field and in the laboratory (e.g., for livestock waste or urban traffic) indicate that collecting a representative sample for multiple NH_3 emission sources remains a challenge. Moreover, this study highlights potential uncertainties in current approaches to isotope-based source apportionment of NH_4^+ and NH_3 . In this regard, the following suggestions could reduce uncertainties in future studies:

1) More field studies are necessary to refine the $\delta^{15}\text{N}(\text{NH}_3)$ source profile of NH_3 emission sources with parallel sampling where possible, with comparison between various active samplers (e.g., scrubbers, denuders) and passive samplers to verify if $\delta^{15}\text{N}(\text{NH}_3)$ measurements are representative of emission sources.

2) Nitrogen equilibrium isotope fractionation factors ($\epsilon_{\text{NH}_4^+ - \text{NH}_3}$, also known as isotopic enrichment factor) between $\text{NH}_3(\text{gas})$ and $\text{NH}_4^+(\text{particle})$ ($\Delta\delta^{15}\text{N}_{\text{NH}_4^+ - \text{NH}_3}$) ($\text{‰} = \delta^{15}\text{N}(\text{NH}_4^+) - \delta^{15}\text{N}(\text{NH}_3)$) (e.g., 10.7 ‰ to 31.4 ‰ (Walters et al., 2019a)) in a field study are not consistent

compared with modeled fractionation effects ($31 \text{ ‰} \pm 4 \text{ ‰}$) (Walters et al., 2019b). More field and laboratory studies are necessary to validate field and theoretically predicted isotopic fractionation between NH_3 gas-to-particle conversions in the atmosphere (Elliott et al., 2019). With validation, prediction of initial $\delta^{15}\text{N}(\text{NH}_3)$ from $\delta^{15}\text{N}(\text{NH}_4^+)$ values can be conducted with more certainty, for isotope-based source apportionment of NH_4^+ .

3) Transport of air mass may alter $\delta^{15}\text{N}(\text{NH}_3)$ values in ambient atmosphere (Yeatman et al., 2001; Felix et al., 2017; Savard et al., 2017) under various environmental (e.g., T , RH, pH) and atmospheric neutralization conditions (e.g., NH_3 -rich, NH_3 -poor conditions). Simultaneous measurements of $\delta^{15}\text{N}(\text{NH}_3)$ and $\delta^{15}\text{N}(\text{NH}_4^+)$ values using validated sample collection method in a vertical tower (at varying heights) could be conducted to increase understanding of such effects on $\delta^{15}\text{N}(\text{NH}_3)$ values and their isotopic fractionation. The consideration of such effects during prediction of initial $\delta^{15}\text{N}(\text{NH}_3)$ values may not only help to improve the isotope-based method but also support our understanding of NH_x dynamics in the atmosphere.

Acknowledgements This work was supported by Key Projects of National Key Research and Development Program of the Ministry of Science and Technology of China (No. 2017YFC0213005) and National Natural Science Foundation of China (Grant No. 21625701), Beijing Municipal Science and Technology Commission (No. Z191100009119001), and the Samsung Advanced Institute of Technology. The authors would like to thank Kate Smith for her help in proofreading our manuscript.

Electronic Supplementary Material Supplementary material is available in the online version of this article at <https://doi.org/10.1007/s11783-021-1414-6> and is accessible for authorized users. Supplementary material includes supplementary tables on summary of recent isotope-based source apportionment studies on ambient NH_3 derived from $\delta^{15}\text{N}(\text{NH}_3)$ values (Table A1); and summary of recent isotope-based source apportionment studies on particulate NH_4^+ derived from $\delta^{15}\text{N}(\text{NH}_4^+)$ values (Table A2).

References

- Altieri K E, Hastings M G, Peters A J, Oleynik S, Sigman D M (2014). Isotopic evidence for a marine ammonium source in rainwater at Bermuda. *Global Biogeochemical Cycles*, 28(10): 1066–1080
- Bateman A S, Kelly S D (2007). Fertilizer nitrogen isotope signatures. *Isotopes in Environmental and Health Studies*, 43(3): 237–247
- Behera S N, Sharma M (2012). Transformation of atmospheric ammonia and acid gases into components of $\text{PM}_{2.5}$: An environmental chamber study. *Environmental Science and Pollution Research International*, 19(4): 1187–1197
- Behera S N, Sharma M, Aneja V P, Balasubramanian R (2013). Ammonia in the atmosphere: A review on emission sources, atmospheric chemistry and deposition on terrestrial bodies. *Environmental Science and Pollution Research International*, 20(11): 8092–8131
- Berner A H, Felix J D (2020). Investigating ammonia emissions in a coastal urban airshed using stable isotope techniques. *Science of the Total Environment*, 707: 134952
- Bhattarai N, Wang S, Xu Q, Dong Z, Chang X, Jiang Y, Zheng H (2020). Sources of gaseous NH_3 in urban Beijing from parallel sampling of NH_3 and NH_4^+ , their nitrogen isotope measurement and modeling. *Science of the Total Environment*, 747: 141361
- Chang Y, Liu X, Deng C, Dore A J, Zhuang G (2016). Source apportionment of atmospheric ammonia before, during, and after the 2014 APEC summit in Beijing using stable nitrogen isotope signatures. *Atmospheric Chemistry and Physics*, 16(18): 11635–11647
- Chang Y, Zou Z, Zhang Y, Deng C, Hu J, Shi Z, Dore A J, Collett J L Jr (2019). Assessing contributions of agricultural and nonagricultural emissions to atmospheric ammonia in a Chinese megacity. *Environmental Science & Technology*, 53(4): 1822–1833
- Ding D, Xing J, Wang S X, Chang X, Hao J M (2019). Impacts of emissions and meteorological changes on China's ozone pollution in the warm seasons of 2013 and 2017. *Frontiers of Environmental Science & Engineering*, 13(5): 76
- Douglas P, Fecht D, Jarvis D (2021). Characterising populations living close to intensive farming and composting facilities in England. *Frontiers of Environmental Science & Engineering*, 15(3): 40
- Elliott E M, Yu Z, Cole A S, Coughlin J G (2019). Isotopic advances in understanding reactive nitrogen deposition and atmospheric processing. *Science of the Total Environment*, 662: 393–403
- Felix J D, Elliott E M, Gay D A (2017). Spatial and temporal patterns of nitrogen isotopic composition of ammonia at U.S. ammonia monitoring network sites. *Atmospheric Environment*, 150: 434–442
- Felix J D, Elliott E M, Gish T, Maghirang R, Cambal L, Clougherty J (2014). Examining the transport of ammonia emissions across landscapes using nitrogen isotope ratios. *Atmospheric Environment*, 95: 563–570
- Felix J D, Elliott E M, Gish T J, McConnell L L, Shaw S L (2013). Characterizing the isotopic composition of atmospheric ammonia emission sources using passive samplers and a combined oxidation-bacterial denitrifier approach. *Rapid Communications in Mass Spectrometry*, 27(20): 2239–2246
- Freyer H D (1978). Seasonal trends of NH_4^+ and NO_3^- nitrogen isotope composition in rain collected at Jülich, Germany. *Tellus*, 30(1): 83–92
- Ge B, Xu X, Ma Z, Pan X, Wang Z, Lin W, Ouyang B, Xu D, Lee J, Zheng M, Ji D, Sun Y, Dong H, Squires F A, Fu P, Wang Z (2019). Role of ammonia on the feedback between AWC and inorganic aerosol formation during heavy pollution in the North China Plain. *Earth and Space Science* (Hoboken, N.J.), 6(9): 1675–1693
- Heaton T H E (1987). $^{15}\text{N}/^{14}\text{N}$ ratios of nitrate and ammonium in rain at Pretoria, South Africa. *Atmospheric Environment*, 21(4): 843–852
- Heaton T H E, Spiro B, Robertson S M C (1997). Potential canopy influences on the isotopic composition of nitrogen and sulphur in atmospheric deposition. *Oecologia*, 109(4): 600–607
- Högberg P (1997). ^{15}N natural abundance in soil-plant systems. *New Phytologist*, 137(2): 179–203
- Hristov A N, Zaman S, Vander Pol M, Ndegwa P, Campbell L, Silva S (2009). Nitrogen losses from dairy manure estimated through nitrogen mass balance and chemical markers. *Journal of Environmental Quality*, 38(6): 2438–2448
- Huang R J, Zhang Y, Bozzetti C, Ho K F, Cao J J, Han Y, Daellenbach K R, Slowik J G, Platt S M, Canonaco F, Zotter P, Wolf R, Pieber S M, Bruns E A, Crippa M, Ciarelli G, Piazzalunga A, Schwikowski M, Abbaszade G, Schnelle-Kreis J, Zimmermann R, An Z, Szidat S, Baltensperger U, El Haddad I, Prevot A S (2014). High secondary aerosol contribution to particulate pollution during haze events in China. *Nature*, 514(7521): 218–222
- Huang S, Elliott E M, Felix J D, Pan Y, Liu D, Li S, Li Z, Zhu F, Zhang N, Fu P, Fang Y (2019). Seasonal pattern of ammonium ^{15}N natural abundance in precipitation at a rural forested site and implications for NH_3 source partitioning. *Environmental Pollution*, 247: 541–549
- Jiang Y Q, Xing J, Wang S X, Chang X, Liu S C, Shi A J, Liu B X, Sahu S K (2021). Understand the local and regional contributions on air pollution from the view of human health impacts. *Frontiers of Environmental Science & Engineering*, 15(5): 88
- Kawashima H, Kurahashi T (2011). Inorganic ion and nitrogen isotopic compositions of atmospheric aerosols at Yurihonjo, Japan: Implications for nitrogen sources. *Atmospheric Environment*, 45(35): 6309–6316
- Kawashima H, Ogata R, Gunji T (2021). Laboratory-based validation of a passive sampler for determination of the nitrogen stable isotope ratio of ammonia gas. *Atmospheric Environment*, 245: 118009
- Keeling C D (1958). The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas. *Geochimica et Cosmochimica Acta*, 13(4): 322–334
- Keeling C D (1961). The concentration and isotopic abundances of carbon dioxide in rural and marine air. *Geochimica et Cosmochimica Acta*, 24(3–4): 277–298
- Koutrakis P, Sioutas C T, Ferguson S M, Wolfson J D, Mulik J M, Burton R (1993). Development and evaluation of a glass honeycomb denuder/filter pack system to collect atmospheric gases and particles.

- Environmental Science & Technology, 27(12): 2497–2501
- Lee C, Hristov A N, Cassidy T, Heyler K (2011). Nitrogen isotope fractionation and origin of ammonia nitrogen volatilized from cattle manure in simulated storage. *Atmosphere*, 2(3): 256–270
- Liu D, Fang Y, Tu Y, Pan Y (2014). Chemical method for nitrogen isotopic analysis of ammonium at natural abundance. *Analytical Chemistry*, 86(8): 3787–3792
- Liu J, Ding P, Zong Z, Li J, Tian C, Chen W, Chang M, Salazar G, Shen C, Cheng Z, Chen Y, Wang X, Szidat S, Zhang G (2018). Evidence of rural and suburban sources of urban haze formation in China: a case study from the Pearl River Delta Region. *Journal of Geophysical Research, D, Atmospheres*, 123(9): 4712–4726
- Meng Z, Xu X, Lin W, Ge B, Xie Y, Song B, Jia S, Zhang R, Peng W, Wang Y, Cheng H, Yang W, Zhao H (2018). Role of ambient ammonia in particulate ammonium formation at a rural site in the North China Plain. *Atmospheric Chemistry and Physics*, 18(1): 167–184
- Moore H (1974). Isotopic measurement of atmospheric nitrogen compounds. *Tellus*, 26(1–2): 169–174
- Moore H (1977). The isotopic composition of ammonia, nitrogen dioxide and nitrate in the atmosphere. *Atmospheric Environment*, 11(12): 1239–1243
- Pan Y, Gu M, He Y, Wu D, Liu C, Song L, Tian S, Lü X, Sun Y, Song T, Walters W W, Liu X, Martin N A, Zhang Q, Fang Y, Ferracci V, Wang Y (2020a). Revisiting the concentration observations and source apportionment of atmospheric ammonia. *Advances in Atmospheric Sciences*, 37(9): 933–938
- Pan Y, Tian S, Liu D, Fang Y, Zhu X, Gao M, Gao J, Michalski G, Wang Y (2018a). Isotopic evidence for enhanced fossil fuel sources of aerosol ammonium in the urban atmosphere. *Environmental Pollution*, 238: 942–947
- Pan Y, Tian S, Liu D, Fang Y, Zhu X, Zhang Q, Zheng B, Michalski G, Wang Y (2016). Fossil fuel combustion-related emissions dominate atmospheric ammonia sources during severe haze episodes: evidence from ^{15}N -stable isotope in size-resolved aerosol ammonium. *Environmental Science & Technology*, 50(15): 8049–8056
- Pan Y P, Gu M N, Song L L, Tian S L, Wu D M, Walters W W, Yu X N, Lu X M, Ni X, Wang Y J, Cao J, Liu X J, Fang Y T, Wang Y S (2020b). Systematic low bias of passive samplers in characterizing nitrogen isotopic composition of atmospheric ammonia. *Atmospheric Research*, 243: 105018
- Pan Y P, Tian S L, Liu D W, Fang Y T, Zhu X Y, Gao M, Wentworth G R, Michalski G, Huang X J, Wang Y S (2018b). Source apportionment of aerosol ammonium in an ammonia-rich atmosphere: An isotopic study of summer clean and hazy days in urban Beijing. *Journal of Geophysical Research, D, Atmospheres*, 123(10): 5681–5689
- Pathak R K, Yao X, Lau A K H, Chan C K (2003). Acidity and concentrations of ionic species of $\text{PM}_{2.5}$ in Hong Kong. *Atmospheric Environment*, 37(8): 1113–1124
- Phillips D L, Gregg J W (2001). Uncertainty in source partitioning using stable isotopes. *Oecologia*, 127(2): 171–179
- Phillips D L, Gregg J W (2003). Source partitioning using stable isotopes: Coping with too many sources. *Oecologia*, 136(2): 261–269
- Phillips D L, Inger R, Bearhop S, Jackson A L, Moore J W, Parnell A C, Semmens B X, Ward E J (2014). Best practices for use of stable isotope mixing models in food-web studies. *Canadian Journal of Zoology*, 92(10): 823–835
- Puchalski M A, Sather M E, Walker J T, Lehmann C M, Gay D A, Mathew J, Robarge W P (2011). Passive ammonia monitoring in the United States: comparing three different sampling devices. *Journal of Environmental Monitoring*, 13(11): 3156–3167
- Savard M M, Cole A, Smirnov A, Vet R (2017). $\delta^{15}\text{N}$ values of atmospheric N species simultaneously collected using sector-based samplers distant from sources – Isotopic inheritance and fractionation. *Atmospheric Environment*, 162: 11–22
- Sigman D M, Casciotti K L, Andreani M, Barford C, Galanter M, Böhlke J K (2001). A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater. *Analytical Chemistry*, 73(17): 4145–4153
- Stratton J J, Ham J, Collett J L Jr, Benedict K, Borch T (2019). Assessing the efficacy of nitrogen isotopes to distinguish Colorado Front Range ammonia sources affecting Rocky Mountain National Park. *Atmospheric Environment*, 215: 116881
- Sutton M A, Erisman J W, Dentener F, Möller D (2008). Ammonia in the environment: From ancient times to the present. *Environmental Pollution*, 156(3): 583–604
- Tang Y S, Cape J N, Sutton M A (2001). Development and types of passive samplers for monitoring atmospheric NO_2 and NH_3 concentrations. *The Scientific World Journal*, 1: 513–529
- Ti C, Ma S, Peng L, Tao L, Wang X, Dong W, Wang L, Yan X (2021). Changes of $\delta(15)\text{N}$ values during the volatilization process after applying urea on soil. *Environmental Pollution*, 270: 116204
- Urey H C (1947). The thermodynamic properties of isotopic substances. *Journal of the Chemical Society (Resumed)*: 562–581
- Walters W W, Blum D E, Hastings M G (2019a). Selective collection of particulate ammonium for nitrogen isotopic characterization using a denuder–filter pack sampling device. *Analytical Chemistry*, 91(12): 7586–7594
- Walters W W, Chai J J, Hastings M G (2019b). Theoretical phase resolved ammonia-ammonium nitrogen equilibrium isotope exchange fractionations: Applications for tracking atmospheric ammonia gas-to-particle conversion. *ACS Earth & Space Chemistry*, 3(1): 79–89
- Walters W W, Goodwin S R, Michalski G (2015). Nitrogen stable isotope composition ($\delta^{15}\text{N}$) of vehicle-emitted NO_x . *Environmental Science & Technology*, 49(4): 2278–2285
- Walters W W, Hastings M G (2018). Collection of ammonia for high time-resolved nitrogen isotopic characterization utilizing an acid-coated honeycomb denuder. *Analytical Chemistry*, 90(13): 8051–8057
- Walters W W, Song L, Chai J, Fang Y, Colombi N, Hastings M G (2020). Characterizing the spatiotemporal nitrogen stable isotopic composition of ammonia in vehicle plumes. *Atmospheric Chemistry and Physics*, 20(19): 11551–11567
- Wang G, Zhang R, Gomez M E, Yang L, Levy Zamora M, Hu M, Lin Y, Peng J, Guo S, Meng J, Li J, Cheng C, Hu T, Ren Y, Wang Y, Gao J, Cao J, An Z, Zhou W, Li G, Wang J, Tian P, Marrero-Ortiz W, Secrest J, Du Z, Zheng J, Shang D, Zeng L, Shao M, Wang W, Huang Y, Wang Y, Zhu Y, Li Y, Hu J, Pan B, Cai L, Cheng Y, Ji Y, Zhang F, Rosenfeld D, Liss P S, Duce R A, Kolb C E, Molina M J (2016). Persistent sulfate formation from London Fog to Chinese haze.

- Proceedings of the National Academy of Sciences of the United States of America, 113(48): 13630–13635
- Wang S, Nan J, Shi C, Fu Q, Gao S, Wang D, Cui H, Saiz-Lopez A, Zhou B (2015). Atmospheric ammonia and its impacts on regional air quality over the megacity of Shanghai, China. *Scientific Reports*, 5(1): 15842
- Wei L, Duan J, Tan J, Ma Y, He K, Wang S, Huang X, Zhang Y (2015). Gas-to-particle conversion of atmospheric ammonia and sampling artifacts of ammonium in spring of Beijing. *Science China. Earth Sciences*, 58(3): 345–355
- Wu C, Wang G, Li J, Li J, Cao C, Ge S, Xie Y, Chen J, Liu S, Du W, Zhao Z, Cao F (2020). Non-agricultural sources dominate the atmospheric NH_3 in Xi'an, a megacity in the semi-arid region of China. *Science of the Total Environment*, 722: 137756
- Wu L B, Ren H, Wang P, Chen J, Fang Y T, Hu W, Ren L J, Deng J J, Song Y, Li J, Sun Y L, Wang Z F, Liu C Q, Ying Q, Fu P Q (2019a). Aerosol ammonium in the urban boundary layer in Beijing: insights from nitrogen isotope ratios and simulations in summer 2015. *Environmental Science & Technology Letters*, 6(7): 389–395
- Wu S P, Zhu H, Liu Z, Dai L H, Zhang N, Schwab J J, Yuan C S, Yan J P (2019b). Nitrogen isotope composition of ammonium in $\text{PM}_{2.5}$ in the Xiamen, China: Impact of non-agricultural ammonia. *Environmental Science and Pollution Research International*, 26(25): 25596–25608
- Xiao H W, Wu J F, Luo L, Liu C, Xie Y J, Xiao H Y (2020). Enhanced biomass burning as a source of aerosol ammonium over cities in central China in autumn. *Environmental Pollution*, 266: 115278
- Xu J, Chen J, Zhao N, Wang G, Yu G, Li H, Huo J, Lin Y, Fu Q, Guo H, Deng C, Lee S H, Chen J, Huang K (2020). Importance of gas-particle partitioning of ammonia in haze formation in the rural agricultural environment. *Atmospheric Chemistry and Physics*, 20(12): 7259–7269
- Ye X, Ma Z, Zhang J, Du H, Chen J, Chen H, Yang X, Gao W, Geng F (2011). Important role of ammonia on haze formation in Shanghai. *Environmental Research Letters*, 6(2): 024019
- Yeatman S G, Spokes L J, Dennis P F, Jickells T D (2001). Comparisons of aerosol nitrogen isotopic composition at two polluted coastal sites. *Atmospheric Environment*, 35(7): 1307–1320
- Zhang L, Altabet M A, Wu T, Hadas O (2007). Sensitive measurement of $\text{NH}_4^{+15}\text{N}/^{14}\text{N}$ ($\delta^{15}\text{NH}_4^{+}$) at natural abundance levels in fresh and saltwaters. *Analytical Chemistry*, 79(14): 5297–5303
- Zhang Y, Benedict K B, Tang A, Sun Y, Fang Y, Liu X (2020a). Persistent nonagricultural and periodic agricultural emissions dominate sources of ammonia in urban Beijing: evidence from ^{15}N stable isotope in vertical profiles. *Environmental Science & Technology*, 54(1): 102–109
- Zhang Y, Liu X, Fang Y, Liu D, Tang A, Collett J L (2020b). Atmospheric ammonia in Beijing during the COVID-19 outbreak: Concentrations, sources, and implications. *Environmental Science & Technology Letters*, 8(1): 32–38
- Zhang Z, Zeng Y, Zheng N, Luo L, Xiao H, Xiao H (2020c). Fossil fuel-related emissions were the major source of NH_3 pollution in urban cities of northern China in the autumn of 2017. *Environmental Pollution*, 256: 113428
- Zhao M, Wang S, Tan J, Hua Y, Wu D, Hao J (2016). Variation of urban atmospheric ammonia pollution and its relation with $\text{PM}_{2.5}$ chemical property in winter of Beijing, China. *Aerosol and Air Quality Research*, 16(6): 1390–1402