Correction to: Basophil Activation Test: Old and New Applications in Allergy

Oliver Hemmings^{1,2,3} · Matthew Kwok^{1,2,3} · Richard McKendry^{1,2,3} · Alexandra F. Santos^{1,2,3,4}

Published online: 27 November 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

**Correction to: Current Allergy and Asthma Reports
(2018) 18: 77**

<https://doi.org/10.1007/s11882-018-0831-5>

Under the heading *BAT in the Follow-up of Patients Submitted to Allergen-Specific Immunotherapy and Other Immunomodulatory Treatments*, the second sentence in the paragraph should read as follows:

Basophil sensitivity (as measured by CD-sens) has shown to be reduced following AIT to aeroallergens such as birch pollen [97, 98] and timothy grass pollen [99], and following AIT to insect venom [100, 101].

The fourth through sixth sentences should read as follows: Interestingly, individuals with a lower antibody-specific activity (i.e. lower percentage of allergen-specific IgE), displayed a greater efficacy to anti-IgE treatment, as determined by BAT [103]. However, the intrinsic sensitivity of the basophil itself also appears to be modified during anti-IgE therapy, and contributes, together with antibody specific activity, to alter the basophils response to challenge.

The online version of the original article can be found at <https://doi.org/10.1007/s11882-018-0831-5>

✉ Alexandra F. Santos
alexandra.santos@kcl.ac.uk

- ¹ Department of Women and Children's Health (Paediatric Allergy), School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK
- ² Peter Gorer Department of Immunobiology, School of Immunology and Microbial Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK
- ³ MRC and Asthma UK Centre in Allergic Mechanisms of Asthma, London, UK
- ⁴ Children's Allergies Department, Guy's and St. Thomas' NHS Foundation Trust, St. Thomas' Hospital, Westminster Bridge Road, London, UK