

ERRATUM

## Erratum to: Surgical Hip Dislocation for Treatment of Femoroacetabular Impingement: Factors Predicting 5-year Survivorship

Simon D. Steppacher MD, Carmen Huemmer MD, MSc,  
Joseph M. Schwab MD, Moritz Tannast MD,  
Klaus A. Siebenrock MD

Received: 24 November 2013 / Accepted: 26 November 2013 / Published online: 7 December 2013  
© The Association of Bone and Joint Surgeons® 2013

**Erratum to: Clin Orthop Relat Res**  
**DOI 10.1007/s11999-013-3268-0**

In the published study by Steppacher and colleagues, the scale of the Y-axis in Figure 4 is incorrect. The scale of the Y-axis should be ranging from 50% to 100%.

The authors apologize for the error.


---

The online version of the original article can be found under  
doi:[10.1007/s11999-013-3268-0](https://doi.org/10.1007/s11999-013-3268-0).

---

S. D. Steppacher, C. Huemmer, M. Tannast,  
K. A. Siebenrock (✉)  
Department of Orthopaedic Surgery, Inselspital,  
University of Bern, Freiburgstrasse 3010 Bern, Switzerland  
e-mail: klaus.siebenrock@insel.ch

J. M. Schwab  
Department of Orthopaedic Surgery, Medical College  
of Wisconsin, Milwaukee, WI, USA



**Fig. 4** Graph showing survival rate (conversion to THA) and followup of selected studies in the literature for open and arthroscopic (asterisk) treatment of FAI. The size indicates the number of hips

included and the gray scale indicates the percentage of labral reattachment from no reattachment (white) to 100% (black).