

## Erratum to: Mesenchymal Stem Cells in Synovial Fluid Increase After Meniscus Injury


Yu Matsukura MD, Takeshi Muneta MD, PhD,  
Kunikazu Tsuji PhD, Hideyuki Koga MD, PhD,  
Ichiro Sekiya MD, PhD

Received: 16 December 2013 / Accepted: 18 December 2013 / Published online: 3 January 2014  
© The Association of Bone and Joint Surgeons® 2013

**Erratum to: Clin Orthop Relat Res**  
DOI 10.1007/s11999-013-3418-4

In the published study by Matsukura and colleagues, the scale of the Y-axis in Figure 6 is incorrect. The scale of the Y-axis should be ranging from 0 to 1200.

The authors apologize for this error.



**Fig. 6** Colony number per synovial fluid ( $\text{mL}^{-1}$ ) was positively correlated with postinjury period ( $n = 22$ ,  $r = 0.773$ ,  $p < 0.001$ ).

The online version of the original article can be found under doi:[10.1007/s11999-013-3418-4](https://doi.org/10.1007/s11999-013-3418-4).

Y. Matsukura, T. Muneta, H. Koga  
Department of Joint Surgery and Sports Medicine, Graduate School, Tokyo Medical and Dental University, Tokyo, Japan

K. Tsuji  
Department of Cartilage Regeneration, Graduate School, Tokyo Medical and Dental University, Tokyo, Japan

I. Sekiya (✉)  
Center for Stem Cell and Regenerative Medicine, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan  
e-mail: [sekiya.arm@tmd.ac.jp](mailto:sekiya.arm@tmd.ac.jp)