Author Correction: Identification of a Novel Feruloyl Esterase by Functional Screening of a Soil Metagenomic Library

Xuanxuan Li¹ • Jia Guo¹ • Yimin Hu¹ • Yumeng Yang¹ • Junwei Jiang¹ • Fang Nan¹ • Shenglu Wu¹ • Zhihong Xin¹

Published online: 26 February 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Author Correction: Applied Biochemistry and Biotechnology (2019) 187:424–437
<https://doi.org/10.1007/s12010-018-2832-1>

The original version of this article unfortunately contained a mistake in the image and caption of Fig. 6. The corrected version of the image and caption is shown here.

The online version of the original article can be found at <https://doi.org/10.1007/s12010-018-2832-1>

✉ Zhihong Xin
xzhfood@njau.edu.cn

Xuanxuan Li
2016808110@njau.edu.cn

Jia Guo
2015208019@njau.edu.cn

Yimin Hu
2015108060@njau.edu.cn

Yumeng Yang
2015108059@njau.edu.cn

Junwei Jiang
2016108058@njau.edu.cn

Fang Nan
2016108057@njau.edu.cn

Shenglu Wu
2016108059@njau.edu.cn

¹ Key Laboratory of Food Processing and Quality Control, College of Food Science and Technology, Nanjing Agricultural University, Nanjing 210095, People's Republic of China

Fig. 6 SDS-PAGE analysis of FAE-Xuan. Lane 1: negative control (*E. coli* harboring the empty pET28a). Lane 2: crude protein extracts of recombinant cells. Lane 3: 20 mM imidazole elution. Lane 4: 100 mM imidazole elution. Lane 5: 200 mM imidazole elution. Lane 6: 300 mM imidazole elution. Lane 7: 400 mM imidazole elution. M: Protein marker Lane M: standard protein marker

The author apologizes for this oversight and for any confusion it may have caused.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.