Correction to: Pathophysiology of Hereditary Angioedema (HAE) Beyond the SERPING1 Gene

Jyoti Sharma¹ · Ankur Kumar Jindal¹ · Aaqib Zaffar Banday¹ · Anit Kaur¹ · Amit Rawat¹ · Surjit Singh¹ · Hilary Longhurst²

Published online: 17 February 2021

© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to: Clinical Reviews in Allergy & Immunology
<https://doi.org/10.1007/s12016-021-08835-8>

The original version of this article unfortunately contained a mistake. The image of Fig. 1 should be corrected. In the previous figure, we had shown that the plasma kallikrein cleaves both high molecular weight kininogen (HMWK) and low molecular weight kininogen (LMWK) to produce bradykinin. However, the plasma kallikrein activates only high molecular weight kininogen (HMWK) while low molecular weight kininogen (LMWK) is activated by tissue kallikrein. Hence we have removed low molecular weight kininogen (LMWK) in the revised figure. We wish to acknowledge Dr. Allen P. Kaplan, Medical University of South Carolina, Charleston, SC, USA for pointing out the error in Fig.1. The corrected Fig. 1 is given below.

The original article can be found online at <https://doi.org/10.1007/s12016-021-08835-8>.

✉ Ankur Kumar Jindal
ankurjindal11@gmail.com

¹ Allergy Immunology Unit, Department of Pediatrics, Postgraduate Institute of Medical Education and Research, Chandigarh, India

² Department of Clinical Immunology, University College Hospitals, London and Addenbrooke's Hospital, Cambridge, UK

The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.