Correction to: High-Mobility Group Box-1-Induced Angiogenesis After Indirect Bypass Surgery in a Chronic Cerebral Hypoperfusion Model

Shingo Nishihiro¹ · Tomohito Hishikawa¹ · Masafumi Hiramatsu¹ · Naoya Kidani¹ · Yu Takahashi¹ · Satoshi Murai¹ · Kenji Sugiu¹ · Yusuke Higaki² · Takao Yasuhara¹ · Cesario V. Borlongan³ · Isao Date¹

Published online: 26 February 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: NeuroMolecular Medicine (2019) 21:391–400
<https://doi.org/10.1007/s12017-019-08541-x>

The original version of this article unfortunately contained a mistake. Figure 5a, b were incorrect. The correct figures are given below.

The original article can be found online at <https://doi.org/10.1007/s12017-019-08541-x>.

✉ Shingo Nishihiro
n.shingo777@gmail.com

✉ Cesario V. Borlongan
cborlong@health.usf.edu

¹ Department of Neurological Surgery, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Science, 2-5-1, Shikata-cho, kita-ku, Okayama 700-8558, Japan

² Department of Pharmaceutical Analytical Chemistry, Okayama University Graduate School of Medicine, Dentistry, and Pharmaceutical Science, 1-1-1, Tsushima naka, kita-ku, Okayama 700-8530, Japan

³ Department of Neurosurgery and Brain Repair, University of South Florida Morsani College of Medicine, 12901 Bruce B Downs Blvd, Tampa, FL 33612, USA

Fig. 5 Cerebral blood flow 14 days after EMS. **a–c** Representative SPECT image obtained 14 days after EMS showing improved cerebral perfusion in both groups (A: control, B: HMGB-1, C: healthy). **d** The ratio of radiation intensity on the operated side compared to

the non-operated side was significantly higher in the HMGB1-treated group than that in the control group ($*p < 0.01$). The healthy rats show no apparent laterality of cerebral blood flow