


## Erratum to: Preliminary biochemical characterization of the novel, non-AT1, non-AT2 angiotensin binding site from the rat brain

Vardan T. Karamyan · Jason Arsenault · Emanuel Escher · Robert C. Speth

Published online: 6 May 2010  
© Springer Science+Business Media, LLC 2010

### Erratum to: Endocr DOI 10.1007/s12020-010-9328-2

In the original publication, the right panel of Figure 3 contains an error in the x axis. The corrected panel appears here.



**Fig. 3** Analysis of N-glycosylation status of the novel angiotensin binding site (representative results from three independent experiments). *Left* SDS-PAGE (7.5%) analysis and Coomassie blue staining of radio-photolabeled ( $^{125}\text{I}$ -SBpa-AngII), semipurified novel angiotensin binding site without (BS) and with (BS + PNGaseF) PNGaseF treatment. Glycoprotein fetuin was used as a positive control in this experiment. *Right* migration of  $^{125}\text{I}$  in the “BS” and “BS + PNGase” lanes of the same SDS gel ( $\sim 0.3$  cm sections)

The online version of the original article can be found under doi:  
[10.1007/s12020-010-9328-2](https://doi.org/10.1007/s12020-010-9328-2).

V. T. Karamyan  
Department of Pharmaceutical Sciences, School of Pharmacy,  
Texas Tech University HSC, Amarillo, TX 79106, USA  
e-mail: vardan.karamyan@ttuhsc.edu

J. Arsenault · E. Escher  
Department of Pharmacology, Université de Sherbrooke,  
Sherbrooke, QC, Canada  
e-mail: Jason.Arsenault@USherbrooke.ca

E. Escher  
e-mail: emanuel.escher@USherbrooke.ca

R. C. Speth (✉)  
Department of Pharmaceutical Sciences, College of Pharmacy,  
Nova Southeastern University, 3200 S. University Dr, Fort  
Lauderdale, FL 33328-2018, USA  
e-mail: Robert.speth@nova.edu