



Correction to: Influence of shunt occlusion on liver volume and functions in hyperammonemic cirrhosis patients having large porto-systemic shunts: a randomized control trial

Amar Mukund¹ · Shakti Prasad Choudhury² · Tara Prasad Tripathy¹ · Venkatesh Hosur Ananthashayana¹ · Rakesh Kumar Jagdish² · Vinod Arora² · Satender Pal Singh² · Ajay Kumar Mishra² · Shiv Kumar Sarin² 

Received: 16 September 2022 / Accepted: 16 September 2022 / Published online: 10 October 2022
© Asian Pacific Association for the Study of the Liver 2022

Correction to: Hepatology International
<https://doi.org/10.1007/s12072-022-10418-4>

The figure legends for Figs. 1 and 2 have been swapped in the original version of the article. The correct figure and legend should read as follows:

The original article can be found online at <https://doi.org/10.1007/s12072-022-10418-4>.

✉ Shiv Kumar Sarin
shivsarin@gmail.com; sksarin@ilbs.in

¹ Department of Intervention Radiology, ILBS, New Delhi, India

² Department of Hepatology, Institute of Liver and Biliary Sciences, ILBS, New Delhi 110070, India





Fig. 1 a Fluoroscopic spot image showing vascular plug (white arrow) with sclerosant in gastrosplenic shunt (arrowhead). Pre procedure volume (b) and post procedure (c) increase in volume

Fig. 2 Consort diagram depicting study design (standard medical treatment vs standard medical treatment plus shunt occlusion)



Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.