

## Erratum to: Visceral obesity stimulates anaphase bridge formation and spindle assembly checkpoint dysregulation in radioresistant oesophageal adenocarcinoma

A. M. Mongan<sup>1</sup> · N. Lynam-Lennon<sup>1</sup> · R. Casey<sup>1</sup> · S. Maher<sup>2</sup> ·  
G. Pidgeon<sup>1</sup> · J. V. Reynolds<sup>1</sup> · J. O'Sullivan<sup>1</sup>

Published online: 20 November 2015  
© Federación de Sociedades Españolas de Oncología (FESEO) 2015

### Erratum to: Clin Transl Oncol DOI 10.1007/s12094-015-1411-y

The section “Patient recruitment and anthropometry” contained the following incorrect statement:

Visceral fat area was calculated by a radiologist using a previously standardised and validated technique, and visceral fat areas in excess of 130 m<sup>2</sup> were classified as obese [19].

Correct is:

Visceral fat area was calculated by a radiologist using a previously standardised and validated technique, and patients with visceral fat areas in excess of 163.8 cm<sup>2</sup> (men) and 80.1 cm<sup>2</sup> (women) were classified as obese [19].

---

The online version of the original article can be found under doi:[10.1007/s12094-015-1411-y](https://doi.org/10.1007/s12094-015-1411-y).

---

✉ J. O'Sullivan  
osullij4@tcd.ie

<sup>1</sup> Department of Surgery, Trinity Centre for Health Sciences, St. James's Hospital, Dublin 8, Ireland

<sup>2</sup> Cancer Biology & Therapeutics Lab, School of Biological, Biomedical & Environmental Sciences, University of Hull, Cottingham road, Hull HU6 76X, UK