Correction to: CLIC1 knockout inhibits invasion and migration of gastric cancer by upregulating AMOT-p130 expression

Y. Qiu¹ · Y.-t. Mao¹ · J.-h. Zhu¹ · K. Zhao¹ · J.-f. Wang¹ · J.-m. Huang² · G.-q. Chang² · Y.-t. Guan² · F.-y. Huang² · Y.-j. Hu² · J.-q. Chen¹ · J.-l. Liu¹

Published online: 16 September 2020

© Federación de Sociedades Españolas de Oncología (FESEO) 2020

Correction to: Clinical and Translational Oncology
<https://doi.org/10.1007/s12094-020-02445-0>

In Figure 5B, the representative image of the KD-NC group was mistakenly uploaded. The authors apologize for any confusion this may have caused. The correct Fig. 5 is given below.

The original article can be found online at <https://doi.org/10.1007/s12094-020-02445-0>.

✉ J.-q. Chen
gqhans@163.com

✉ J.-l. Liu
liujinlugg@163.com

¹ Department of Gastrointestinal Surgery, The First Affiliated Hospital of Guangxi Medical University, 6 Shuangyong Road, Nanning 530021, Guangxi Zhuang Autonomous Region, People's Republic of China

² The Guangxi Medical University, Nanning, Guangxi Zhuang Autonomous Region, People's Republic of China

Fig. 5 AMOT-p130 inhibits invasion and migration of GC cells. **a** The cell migration assay showed that the number of cells passing through the membrane in the KD group was significantly higher than that in the KD-NC group of cells ($P < 0.05$), while the number of cells crossing through the membrane in the OE group was less than the OE-NC group of cells ($P < 0.05$). **b** The cell invasion assay

showed that the number of cells invaded in the KD group was higher than the KD-NC group ($P < 0.05$), while the OE group was decreased compared with the OE-NC group ($P < 0.05$). **c** The recovery of cells in the KD group was significantly faster than the KD-NC group. The recovery of cells in the OE group was slower than the OE-NC group

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.