Prevention and Control of COVID-19 in Chronic Kidney Disease

Shipin Feng¹ · Min Xie¹ · Wei Luo¹ · Li Wang¹ · Limin Guo¹ · Ying Wu¹ · Jun Liu¹ · Qinwei Duan¹ · Qin Yang¹ · Jia Li¹ · Xi Liu¹ · Rong Zhu¹

Received: 25 February 2020 / Accepted: 13 April 2020 / Published online: 28 April 2020
© The Author(s) 2020

To the Editor: Since the outbreak of COVID-19 in Wuhan, China, the epidemic caused by COVID-19 has become a public health event of international concern, which is seriously threatening the health of people worldwide. As is known, children with chronic kidney disease (CKD) who are treated with long-term glucocorticoids and immunosuppressants belong to the high-risk and susceptible group of the 2019-nCoV. However, up to date, there is no case report of CKD complicated with COVID-19 in China. We here present our experience in the prevention and control of COVID-19 infection in this special group of patients, with a purpose to provide reference for the international clinical community. The general management comprises of proper rest, adequate nutrition support, and symptomatic treatment. For antiviral therapy, α - interferon, lopinavir / ritonavir, or abidol can be used.

Glucocorticoids in mild and common type COVID-19: For CKD children in the consolidation and maintenance stage of hormone treatment, the frequency of oral hormone use can be changed from every other day to every day for 7 consecutive days at the same dose in order to reduce the recurrence of the primary disease [1]; for those in the stage of hormone-induced remission, the dose of hormone can be tapered according to the patient's condition.

Glucocorticoids in COVID-19 severe and critical type: Hormone dosage should be adjusted according to the degree of systemic inflammatory reaction and whether the patient is complicated with acute respiratory distress syn-

drome (ARDS). It is recommended not to exceed 1–2 mg/kg per day used for methylprednisolone.

Immunosuppressants: Cyclosporine and mycophenolate mofetil were found to significantly inhibit the replication of Middle East respiratory syndrome coronavirus (MERS CoV) and severe acute respiratory syndrome coronavirus (SARS CoV) [2]. However, there is no report about the effect of these immunosuppressants on 2019-nCoV up-to-date. Therefore, the use of immunosuppressants in children with CKD should be determined according to the specific condition of the patient.

Management of blood pressure: The pathogenesis of 2019-nCoV is similar to that of SARS virus [3], which infects human cells by recognizing and binding to angiotensin-converting enzyme (ACE)-2 protein; the use of ACE inhibitors (ACEIs) and angiotensin receptor antagonists (ARBs) can increase the expression and activity of ACE-2 [4]. Therefore, it is suggested that ACEIs and ARBs should be discontinued in children with COVID-19 and other antihypertensive drugs should be used instead.

Compliance with Ethical Standards

Conflict of Interest None.

Source of Funding 1. Chengdu Municipal Health Commission 2018052; 2. Health Commission of Sichuan Province 12051.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

✉ Shipin Feng
fengshipin@sina.com

¹ Department of Pediatric Nephrology, Chengdu Women's and Children's Central Hospital, Chengdu, Sichuan, China

References

1. The Nephrology Group of the Pediatric Branch of the Chinese Medical Association. Evidence-based guidelines for the diagnosis and treatment of hormone-sensitive, relapsed / dependent nephrotic syndrome in children. *Chinese J Pediatr*. 2017;55:729–34.
2. Li HS, Kuok DIT, Cheung MC, et al. Effect of interferon alpha and cyclosporine treatment separately and in combination on Middle East respiratory syndrome coronavirus (MERS-CoV) replication in a human in-vitro and ex-vivo culture model. *Antivir Res*. 2018;155:89–96.
3. Zhou P, Yang XL, Wang XG, et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin [J/OL]. *bioRxiv*. 2020. <https://doi.org/10.1101/2020.01.22.914952>. Available at: <https://www.biorxiv.org/content/10.1101/2020.01.22.914952v2.full.pdf>.
4. ACE2 expression is induced by treatment with ACE-inhibitors (ACEIs) or angiotensin II AT1 receptor blockers (ARBs) in many rodent tissues. *Amino Acids*. 2015;47:693–705.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.