Thalassemic Child Presenting with Anosmia due to COVID-19

Wulandewi Marhaeni¹ · Andreas Budi Wijaya² · Prabandari Kusumaningtyas² · Rahmawan Sakup Mapianto²

Received: 19 May 2020 / Accepted: 28 May 2020 / Published online: 9 June 2020
© Dr. K C Chaudhuri Foundation 2020

To the Editor: Novel coronavirus (SARS-CoV-2) infection (COVID-19) has spread rapidly in Indonesia. It might also impact on patients with hemoglobin disorders, such as thalassemia [1]. Indonesia has one of the highest prevalence of thalassemia globally [2].

A 17-y-old girl with transfusion-dependent thalassemia (beta-thalassemia), presented with acute loss of sense of smell (anosmia) and diminished sense of taste (ageusia) since 8 d prior to admission, accompanied by 2 d history of sneezing. History inquiry revealed that she had contact with her father, who was positive for COVID-19. She was admitted to hospital due to pale appearance since 7 d prior to admission. Three days after admission, she also complained mild-moderate muscle aches, which were relieved by using Paracetamol. Lung examination showed normal result.

Laboratory studies showed low hemoglobin level (7.5 g/dl), normal white blood cell count ($9.4 \times 10^3/\mu\text{L}$), with 54.9% neutrophils and 34.2% lymphocytes, prothrombin time prolongation (PT 14.5 s; normal APTT 31.6 s), hyperuricemia (6.3 mg/dl), and slight increase in C-reactive protein level (6 mg/dl). Ferritin level was 980.51 ng/ml, accompanied by elevated transferrin saturation (88%). Oropharyngeal swab tested positive for SARS-CoV-2. Chest X-ray result was normal. Supportive and therapeutic care such as blood transfusion, antibiotic Azithromycin 500 mg/q24h and antiviral Oseltamivir 75 mg/q12h were given. Deferiprone has still been given, considering her good condition.

Anosmia and ageusia might become possible symptoms of COVID-19 in pediatric population with transfusion-dependent

thalassemia, as reported in adult cases [3]. Without full objective assessment and nasoendoscopy, precise pathomechanism is still hard to determine. The possible explanation is post-viral syndrome with direct infection of the olfactory mucous and impairment of olfactory sensory neurons [4].

Erythrocyte is strongly hypothesized for its involvement in COVID-19 pathophysiology, by which SARS-CoV-2 might attack beta chain of hemoglobin. Its turnover will be reduced, followed by cell lysis and development of hemolytic anemia [5]. This condition is interesting, because beta globin chain defect in beta-thalassemia might potentially interrupt the impact of SARS-CoV-2 infection. Therefore, beta-thalassemia patients would not have same risk of COVID-19 as other patients, considering its reduced severity. More studies are needed to have a better understanding of SARS-CoV-2 infection in pediatric population with beta-thalassemia.

Compliance with Ethical Standards

Conflict of Interest None.

References

1. Ji L, Chao S, Wang Y, et al. Clinical features of pediatric patients with COVID-19: a report of two family cluster cases. *World J Pediatr*. 2020. <https://doi.org/10.1007/s12519-020-00356-2>.
2. Grosso M, Sessa R, Puzone S, Storino MR, Izzo P. Molecular basis of thalassemia. In: Donald S, editors. *Anemia*. Israel: Intech Open; 2012. p. 341–60.
3. Hopkins C, Kumar N. Loss of sense of smell as marker of COVID-19 infection. Available at: <https://www.entuk.org/sites/default/files/files/Loss%20of%20sense%20of%20smell%20as%20marker%20of%20COVID.pdf>. Accessed 25 May 2020.
4. Gane SB, Kelly C, Hopkins C. Isolated sudden onset anosmia in COVID-19 infection. A novel syndrome? *Rhinology*. 2020. <https://doi.org/10.4193/Rhin20.114>.
5. Wenzhong L, Hualan L. COVID-19: attacks the 1-beta chain of hemoglobin and captures the porphyrin to inhibit human heme metabolism. *ChemRxiv Preprint*. <https://doi.org/10.26434/chemrxiv.11938173.v5>.

✉ Wulandewi Marhaeni
wdmarhaeni@yahoo.com

¹ Hematology-Oncology Division, Department of Pediatrics, Ulin General Hospital, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, South Kalimantan, Indonesia

² Department of Pediatrics, Ulin General Hospital, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, South Kalimantan, Indonesia