Correction to: Biotinidase Deficiency Presenting as Recurrent Laryngeal Stridor

Sruthi Alanghat¹ · John Matthai² · Vidhya Marimuthu¹ · R. Manivasakan³ · M. Ramaswamy¹

Published online: 16 April 2021

© Dr. K C Chaudhuri Foundation 2021

Correction to: The Indian Journal of Pediatrics.

<https://doi.org/10.1007/s12098-020-03635-y>

The image inside Fig. 1a went missing during the typesetting process. The correct Fig. 1 is shown below.

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s12098-020-03635-y>

✉ Sruthi Alanghat
sruthialanghat@gmail.com

¹ Department of Pediatrics, Masonic Medical Centre for Children, Coimbatore, Tamil Nadu, India

² Department of Gastroenterology, Masonic Medical Centre for Children, Coimbatore, Tamil Nadu, India

³ Department of Neurology, Masonic Medical Centre for Children, Coimbatore, Tamil Nadu, India

Fig. 1 Magnetic resonance imaging (MRI) of the brain and spinal cord done [Diffusion-weighted imaging (DWI) axial section **a-d**, T2- weighted image (T2W1) sagittal section **e**], which showed symmetrical white

matter hyperintensities in corpus callosum (**a**), bilateral cerebral white matter (**b**), medulla (**c**), parts of pons (**d**) with long segment myelopathy of spinal cord from cervicomedullary junction to D10 level (**e**)