



Oxygen Saturation Index to Predict Surfactant Requirement in Preterm Infants

Yasemin Ezgi Kostekci¹ · Emel Okulu¹ · Dilara Gungor Akin² · Omer Erdeve¹ · Begum Atasay¹ · Saadet Arsan¹

Received: 18 July 2022 / Accepted: 16 September 2022 / Published online: 29 September 2022
© The Author(s), under exclusive licence to Dr. K C Chaudhuri Foundation 2022

To the Editor: In the management of respiratory distress syndrome (RDS), which causes major mortality and morbidity in preterms, surfactant is suggested to avoid invasive mechanical ventilation. The guidelines recommend surfactant according to mean airway pressure (MAP)/positive end expiratory pressure (PEEP) and fraction of inspired oxygen - FiO₂ needs [1, 2]. The oxygenation index (OI) is calculated with MAP, FiO₂, and partial oxygen pressure - PaO₂ and used to assess the severity of respiratory failure [3]. The OSI uses oxygen saturation - SpO₂ instead of PaO₂, which is noninvasive and can be measured continuously. The studies found a strong correlation between OSI and OI in neonates [3, 4].

This single-center, retrospective study aimed to predict the surfactant requirement by evaluating OSI. Infants < 30 wk gestational age, who received nasal respiratory support but were not intubated were included. Baseline characteristics and the MAP, FiO₂, and OSI values at admission and before surfactant administration were recorded. For the nonsurfactant group, these values were also recorded at the postnatal sixth hour. The OSI was calculated as following: $OSI = [MAP (cmH_2O) \times FiO_2 (\%)] / SpO_2 (\%)$ [3].

Of the total 76 newborns, 28 required surfactant. The mean gestational age (GA) and birth weight (BW) were similar. The mean OSI values at admission and before surfactant administration were higher in the surfactant group (3.10 ± 1.10 vs. 1.66 ± 0.46 , and 3.98 ± 1.50 vs. 1.62 ± 0.4 , $p < 0.001$, respectively). The receiver operating characteristic (ROC) curves for OSI were statistically significant. The cut-off value of OSI at admission to predict surfactant requirement was 1.91 (89% sensitivity, 77% specificity). The

OSI value just before surfactant administration was found to be 2.33 with a sensitivity of 93%, and with a specificity of 98% (AUC 0.980; 95% CI: 0.952–1.000, $p < 0.001$). These findings highlight that using OSI could be helpful in making surfactant decisions.

To our knowledge, this is the first study about OSI for predicting surfactant requirements. Prospective validation of these cutoffs in a larger sample size can improve clinical relevance and identify potential differences in cutoffs.

Declarations

Conflict of Interest None.

References

1. Sweet DG, Carnielli V, Greisen G, et al. European consensus guidelines on the management of respiratory distress syndrome - 2019 update. *Neonatology*. 2019;115:432–50.
2. Ozkan H, Erdeve O, Kutman HGK. Turkish Neonatal Society guideline on the management of respiratory distress syndrome and surfactant treatment. *Turk Pediatri Ars*. 2018;53:S45–54.
3. Muniraman HK, Song AY, Ramanathan R, et al. Evaluation of oxygen saturation index compared with oxygenation index in neonates with hypoxemic respiratory failure. *JAMA Netw Open*. 2019;2:e191179.
4. Khalesi N, Choobdar FA, Khorasani M, et al. Accuracy of oxygen saturation index in determining the severity of respiratory failure among preterm infants with respiratory distress syndrome. *J Matern Fetal Neonatal Med*. 2021;34:2334–9.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

✉ Yasemin Ezgi Kostekci
ezgikostekci@gmail.com

¹ Division of Neonatology, Department of Pediatrics, Ankara University Faculty of Medicine, 06590 Mamak, Ankara, Turkey

² Department of Pediatrics, Ankara University Faculty of Medicine, Ankara, Turkey