




# Social support as a mediator in the relationship between perceived stress and nomophobia: An Investigation among Malaysian university students during the COVID-19 pandemic

Samantha Arielle Lai<sup>1</sup> · Khong Yun Pang<sup>3</sup> · Ching Sin Siau<sup>4</sup> · Caryn Mei Hsien Chan<sup>4</sup> · Yee Kee Tan<sup>4</sup> · Pei Boon Ooi<sup>3</sup> · Mohamad Ikhran Bin Mohamad Ridzuan<sup>1</sup> · Meng Chuan Ho<sup>1,2</sup> 

Accepted: 18 May 2022 / Published online: 2 June 2022

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2022

## Abstract

This study examined the mediating role of social support in the relationship between perceived stress and nomophobia among Malaysian university students during the COVID-19 pandemic. A cross-sectional study was conducted with  $N = 547$  university students. Participants answered a self-administered questionnaire measuring nomophobia, social support, and perceived stress. Exploratory analyses were conducted using partial least square structural equation modelling. We found that perceived stress was positively associated with nomophobia during the COVID-19 pandemic, whilst social support partially mediated the relationship between perceived stress and nomophobia. The results of this study indicated that stress may be buffered by social support in individuals with higher levels of nomophobia.

**Keywords** Nomophobia · Perceived stress · Social support · Partial least squares · Stress buffering hypothesis

Technological advancement has led to the worldwide adoption of smartphones (Yildirim & Correia, 2015) and with their affordable prices, the dependency on smartphones worldwide has increased (Dasgupta et al., 2017). This has concurrently given rise to the issue of internet addiction, which is significantly associated with a number of psychiatric conditions (Ho et al., 2014). One of the problems associated with this increased dependence is smartphone addiction, which is now a widespread phenomenon (Ahmed et al., 2019). The prevalence of smartphone addiction has been increasing. A study showed that the pooled prevalence of smartphone addiction globally is at 27.0%,

and disproportionately affects low and lower-and-middle income countries (Meng et al., 2022).

A condition associated with smartphone addiction is nomophobia. A number of studies have found a strong association between nomophobia, smartphone addiction, and digital addiction (Çobanoğlu et al., 2021). Nomophobia, which stands for “NoMobilePhone” phobia, refers to an individual’s fear of not having contact with their mobile phone (Gurbuz & Ozkan, 2020; King et al., 2010; Yildirim & Correia, 2015). The concept of nomophobia was constructed based on the criteria for specific phobia in the Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV; American Psychiatric Association, 1994; Battacharya et al., 2019). It has been used to describe pathological fear resulting in anxiety, discomfort, distress, and nervousness caused by the non-availability of a mobile phone (Bragazzi & del Puente, 2014; King et al., 2010).

Nomophobia develops as a result of an individual’s high reliance on mobile phones and frequent use of social media, as well as other activities such as access to entertainment and information gathering from the internet. These eventually may lead to different negative psychological, social, intellectual, and professional outcomes (Beard & Wolf, 2001; Caplan, 2002). Due to the nomophobic individual’s dependence on internet content when using mobile phones,

✉ Meng Chuan Ho  
homc@ucsiuniversity.edu.my

<sup>1</sup> Faculty of Social Sciences and Liberal Arts, UCSI University, No. 1, Jalan Menara Gading, UCSI Heights, 56000 Kuala Lumpur, Malaysia

<sup>2</sup> Faculty of Education, Universiti Malaya, Kuala Lumpur, Malaysia

<sup>3</sup> School of Medical & Life Sciences, Sunway University, Petaling Jaya, Malaysia

<sup>4</sup> Centre for Community Health Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Bangi, Malaysia

nomophobia may constitute a withdrawal symptom of internet addiction. People who have nomophobia are afraid of missing out on social media messages, events, and experiences (Yildirim and Correia, 2015). When they forget to bring their phones, the battery runs out, or there is no network coverage, they tend to feel excessively nervous and anxious (Gezgin et al., 2018). Individuals with nomophobia are prone to have access to their phone almost 24 h a day, and even taking it to bed with them (Bragazzi & Del Puente, 2014). Among university students, nomophobia affects the individuals' academic performance and motivational level during their learning processes (Dixit et al., 2010). Limone and Toto (2021) reported that dependency on the mobile phone has increased during the COVID-19 pandemic, and it is a critical risk factor leading to nomophobia.

A systematic review of nomophobia indicated that, especially among young people, nomophobia was significantly associated with a number of psychological disorders and symptomatology similar to internet addiction (Ho et al., 2014), such as depression, anxiety, stress, and sleep disorders (Bekaroğlu & Yılmaz, 2020; Bhattacharya et al., 2019; Mir & Akhtar, 2020; Rodríguez-García et al., 2020; Rosen et al., 2016). The relationship between nomophobia and stress was investigated in a few studies. For example, Bhuvaneswari and Emiline Joy (2021) found that nomophobia was more likely to occur among individuals who self-reported to be more stressed. Terzioğlu and Uğurlu (2021) found that perceived stress was closely associated with all dimensions of nomophobia, specifically the “Inability to communicate” had the strongest relationship, followed by “To give up comfort”, “Losing online connection,” and “Inability to access information”. On the other hand, social support has been shown to be a protective factor against stress. The findings of Besthorn et al. (2018) showed that perceived stress negatively influenced perceived social support.

Thus far there are no extant studies investigating the relationship between nomophobia and social support. Studies by Ergin and Ozer (2021) and Purnama and Sununianti (2020) found that nomophobia was significantly associated with social alienation, which may be inversely associated with social support. Tams and colleagues (Tams et al., 2018) had also found that nomophobia will lead to a higher level of perceived social threat. Based on a study by Ozdemir et al. (2018), loneliness was the strongest predictor of nomophobia among Turkish and Pakistani undergraduates. In Malaysia, a study by Ali et al. (2017) showed that nomophobia was associated with fear of connection loss and fear of loneliness. Individuals gain positive social support through their smartphones (Gökçearslan et al., 2018). This could have led to the development of the ‘losing connectedness’ dimension of nomophobia because people fear to lose the connectivity with others provided by smartphones (Yildirim & Correia, 2015). Therefore, this study will shed more light

on the role of social support in mediating the relationship between perceived stress and nomophobia, which may positively contribute to the formulation of evidence-based social interventions for nomophobia.

## Theoretical Consideration

We used the stress-buffering hypothesis (Cohen & Wills, 1985) as our theoretical framework to investigate the role of perceived social support in the relationship between perceived stress and nomophobia. The theory proposed that social support can act as a buffer to intervene in the perception of stress/stressful events and stressful reactions, which in turn could increase well-being and lower psychopathology risks (Cohen & Wills, 1985). This theory had been used in different contexts, for instance, in psychological health during the COVID-19 pandemic (Szkody et al., 2021) and smartphone addiction (Wang et al., 2018).

In light of the lack of studies investigating the relationship between social support and nomophobia, we aimed to examine the relationship based on the stress-buffering hypothesis and the relevance of social support as a mediator in the relationship between perceived stress and nomophobia. Social support has been heavily tested as a mediator throughout the literature (Li et al., 2018; Xie et al., 2018; Zhang et al., 2017). We therefore propose the hypotheses below:

*H1: Perceived stress will negatively predict social support*

*H2: Social support will negatively predict nomophobia*

*H3: Perceived stress will positively predict nomophobia*

*H4: Social support will mediate the relationship between perceived stress and nomophobia*

## Methods

### Study Design and Sample

This was a cross-sectional study. The a-priori sample size calculator (Soper, 2021) with an 80% power level and effect size of 0.15 showed that we need at least 545 participants for this study. Inclusion criteria were students who were actively enrolled in a university and owned a smartphone. Individuals unable and unwilling to provide informed consent were excluded.

### Measures

Nomophobia was assessed using the Nomophobia Questionnaire (NMP-Q; Yildirim & Correia, 2015). The NMP-Q consists of 20 items that measure nomophobia based on a

7-point Likert scale on how much participants agree or disagree on each statement. The NMP-Q has a 4-factor dimension, which are not being able to communicate, losing connectedness, not being able to access information, and giving up convenience. Higher scoring on the scale denoted a higher nomophobia level.

Perceived social support was assessed using the Multi-dimensional Scale of Perceived Social Support (MSPSS; Zimet et al., 1988). The MSPSS consists of 12 items that measure the sources of support based on a 7-point Likert scale on how much participants agree or disagree on each statement. The MSPSS had a 3-factor dimension, namely, family, friends, and significant others. Higher scoring on the scale denoted a higher social support level.

Perceived stress was assessed using the Perceived Stress Scale 4 (PSS-4; Cohen et al., 1983). The PSS-4 comprises 4 items that measure stress level based on a 5-point Likert scale, which were 0 = never, 1 = almost never, 2 = sometimes, 3 = fairly often and 4 = very often. The scale has two inversely scored items (item 2 and item 3), in which reverse coding is needed. Higher scoring on the scale denoted a higher stress level.

## Procedures

The survey was carried out using an online survey form. Convenience and snowballing sampling were used. All the questionnaires were self-administered in English.

## Data Analysis

We used partial least square structural equation modeling (PLS-SEM) in this study. According to Henseler et al. (2014), PLS-SEM was a prime candidate for exploratory analysis. Secondly, it was a better statistical method to access the mediation model compared to the more traditional methods such as the Sobel test and PROCESS Macro (Ramayah et al., 2018).

We used SmartPLS as the main statistical tool to assess the measurement and structural models in this study. We used IBM SPSS Statistics for Windows, version 26 to conduct the descriptive analysis.

## Results

Table 1 shows the descriptive information of the survey. In total, we recruited 547 participants. There were 249 male participants and 298 female participants in this study, the majority of whom were Buddhist (72.7%), and 292 of them (53.4%) were classified as the high change readiness group (Table 1). Bivariate correlation analyses were conducted

**Table 1** Socio-demographic Characteristics of Participants (N = 547)

Variables	Frequency	Percent (%)
Gender		
Male	249	45.5
Female	298	54.5
Religion		
Islam	15	2.7
Buddhist	392	72.7
Hindu	20	3.7
Others	120	21.9

prior to fitting the multivariate model. All variables were considered significantly correlated with each other at  $p < 0.05$ .

## Assessment of the Measurement Model

All constructs have higher value of factor loadings, AVE, and CR compared to the threshold value suggested by Hair et al. (2017), but some of the lower-order constructs (LOCs, e.g., not being able to communicate, losing connectedness, family, friends, and significant other) have an exceptionally high CR value which is more than 0.90; this suggests that most of the indicators are measuring the same phenomenon, therefore, decreasing the reliability of the constructs. Besides, one item (e.g., PS3) from the perceived stress construct is deleted to achieve optimum AVE and CR value.

To remedy this, we used the disjointed 2 stage approach (Sarstedt et al., 2019) to assign all the LOCs to their respective higher-order constructs (HOCs), for instance, nomophobia (LOCs: not being able to communicate, losing connectedness, not being able to access information, giving up convenience) and perceived social support (LOCs: family, friends, and significant other) are the 2 HOCs from the current study. Both of the HOCs showed significantly improved CR value.

All the Heterotrait-Monotrait ratio of correlations values (Henseler et al., 2015) fulfilled the criteria of being lower than 0.85 (Williams & Vogt, 2011) and 0.90 (Gold et al., 2001). Therefore, discriminant validity was justified.


## Assessment of the Structural Model

We evaluated the common method bias and multicollinearity issue by referring to the value of the Inner Variance Inflation Factor (VIF). All the Inner VIF values were lower than 3.3, therefore, no common method bias and multicollinearity issues in the structural model were detected (Kock, 2015). We used the segmentation approach to test the mediation model and also test the direct effect of the relationship between perceived stress and nomophobia (Rungtusanatham et al., 2014).

**Table 2** Hypotheses Testing

Hypothesis	Std Beta	Std Error	t-value	95%CI LL	95%CI UL	VAF
PS→PSS	-0.339	0.044	7.699***	-0.407	-0.259	0.02
PSS→N	0.326	0.048	6.811***	0.238	0.396	
PS→N	0.206	0.047	4.405***	0.124	0.279	
PS→PSS→N	-0.111	0.022	4.970***	-0.150	-0.077	

Note. N is nomophobia; PSS is perceived social support; PS is perceived stress. \*\*\* $p < .001$

**Fig. 1** Mediating role of perceived social support in perceived stress and nomophobia

We evaluated the model using bootstrap with 5000 resamples. Table 2 showed that all the proposed hypotheses were significant. Perceived stress negatively influenced social support ( $\beta = -0.339$ ,  $p < 0.001$ ) and positively influenced nomophobia ( $\beta = 0.206$ ,  $p < 0.001$ ), while perceived social support positively influenced nomophobia ( $\beta = 0.326$ ,  $p < 0.001$ ). The indirect effect of the relationship showed that perceived social support negatively mediated the relationship of perceived stress and nomophobia ( $\beta = -0.111$ ,  $p < 0.001$ ). As suggested by (Zhao et al., 2010), we concluded that the model was competitively partially mediated. (Table 2, Fig. 1).

In the model, 10% variance of nomophobia was explained by perceived stress and perceived social support. Both perceived stress ( $f^2 = 0.042$ ) and perceived social support ( $f^2 = 0.105$ ) had a small effect on nomophobia. We used the PLSpredict (Shmueli et al., 2016) to identify the prediction power of the models. In Table 3, half of the indicators (e.g., giving up convenience and not being able to access information) have lower RMSE in the PLS analysis. Therefore, the model is interpreted as having a medium predicting power.

**Table 3** PLSpredict Assessment of Research Model

Indicators	PLS-RMSE	Q <sup>2</sup> _predict	LM-RMSE	PLS-RMSE-LM-RMSE
NC	1.004	-0.004	1.006	-0.002
GV	<b>0.997</b>	<b>0.009</b>	<b>1.001</b>	<b>-0.004</b>
LC	0.999	0.006	0.996	0.003
NI	<b>0.997</b>	<b>0.009</b>	<b>1.002</b>	<b>-0.005</b>

Note. The bolded text and number indicate good predictors; NC is not being able to communicate; LC is losing connectedness; NI is not being able to access information; GC is giving up convenience; PS1 is “in the last month, how often have you felt that you were unable to control the important things in your life?”; PS2 is “in the last month, how often have you felt confident about your ability to handle your personal problems?”; PS4 is “in the last month, how often have you felt difficulties were piling up so high that you could not overcome them?”

Both indicators (e.g., not being able to communicate and losing connectedness) had a high prediction error, and both indicators were related to the social dimension (Table 3).

## Discussion

This study was aimed to examine mediating influence of social support in the relationship between perceived stress and nomophobia among university students during the COVID-19 pandemic. Based on relevant literature, four hypotheses were developed. The results suggested that all the hypotheses were accepted except the second hypothesis, in which we hypothesized that social support would negatively predict nomophobia.

With respect to our first hypothesis, the results from this study confirmed that individuals reporting higher perceived stress had lower social support. This finding was in line with many contemporary studies (Bonavita et al., 2021; Khodarahimi et al., 2016; Roohafza et al., 2016) which highlighted that social support was a key component in buffering the effect of stress, with persons receiving more social support reporting better ability to manage stress in comparison with those who had less social support. Positive social support may alleviate the harmful effects of stress due to the presence of other individuals who are able to provide objective

help (such as offering food and shelter) or subjective help (such as providing a listening ear) to those who are going through a stressful situation (Bender et al., 2019). Social support was also crucial in alleviating psychological issues like depression and anxiety (Guo et al., 2021; Roohafza et al., 2014).

The present study found a positive significant relationship between perceived social support and nomophobia, i.e., individuals experiencing higher levels of social support will also report higher levels of nomophobia. Hence the second hypothesis was rejected. Our finding was in contrast with previous studies which found that social support was negatively associated with nomophobia (Shirani, 2017; Zhou et al., 2021; Zwilling, 2022). During the COVID-19 pandemic lockdown, non-face-to-face contact through social media use was one of the methods to increase social support (Chen et al., 2021; Zhang et al., 2021). Past studies had found that the mobile phone was a means to receive social support through social media (Selkie et al., 2020), and individuals who were addicted to the internet were more open to interpersonal influences from their online friends (Tran et al., 2017). The increased importance of online social support sources, due to limitations to face-to-face social contact imposed by the pandemic, may possibly result in the tendency for higher nomophobia.

Our study found that higher perceived stress was positively associated with higher nomophobia; the results supported our third hypothesis. Our results were consistent with findings obtained by a number of researchers (e.g., Bano et al., 2021; Farchakh et al., 2021; Lelapary et al., 2020; Pavithra et al., 2015). According to Lelapary et al. (2020), university students with nomophobia may have used their mobile phone as a coping tool against study-related or relationship-related stress. However, this coping mechanism to alleviate stress may have become maladaptive, leading to nomophobia.

The fourth hypothesis examined in the present study was the mediating role of social support in the relationship between perceived stress and nomophobia. Regarding the partial mediation of social support in the relationship between perceived stress and nomophobia, our study confirmed that social support acted as a buffer or protective factor against the influence of stress on nomophobia, thus supporting the stress-buffering hypothesis (Cohen & Wills, 1985). Pavithra et al. (2015) suggested that young adults developed stress due to the need for social connection, in which 56% of young adults put social networking as the reason of daily phone usage. This may explain why higher social support was a protective factor against nomophobia, and partially mediated the relationship between stress and nomophobia. In a different context, Tams et al. (2018) examined the relationship between nomophobia and stress using perceived social threat, which saw that high levels

of perceived social threat were associated with increased stress level among nomophobes. A number of studies have also found that loneliness mediated the relationship between stress and nomophobia (e.g., Bano et al., 2021; Kara et al., 2021; Nguyen et al., 2022). People who had higher levels of stress often faced the risk of loneliness (Campagne, 2019); however, they may feel less lonely when social support was offered, especially during stressful situations (Zhang et al., 2018). This may indicate that people who lacked social support may rely on their mobile phones as a tool to seek affiliation with other socialization agents (Herrero et al., 2019). Therefore, social support may play an important role in alleviating perceived stress and nomophobia, and it needs to be considered in interventions for individuals with nomophobia.

## Implications

In clinical settings, when formulating the treatment for individuals with nomophobia, stress levels need to be addressed. Stress management skills and problem-focused solving could be taught to those facing nomophobia. Social support as a mitigating factor needs to be thoroughly assessed and comprehensive social support interventions could be applied to university students who experience social isolation, especially during and after the COVID-19 pandemic. Internet cognitive behavior therapy (I-CBT; Zhang & Ho, 2017) may be considered an amenable treatment modality to address nomophobia symptoms, especially due to the numerous restrictions imposed by the COVID-19 endemic.

This study has a few limitations. Being a cross-sectional study, we could not infer causation. Future cohort or intervention studies would be able to understand the contributions of perceived stress and social support towards the development and maintenance of nomophobia. Even though the study was carried out during the COVID-19 pandemic, we did not measure the impact of COVID-19 on stress level. Finally, being an online survey, we could not control for bias due to using convenience and snowball sampling. Future studies on nomophobia among college students may benefit from randomization and stratification in the sampling methods to ensure greater representativeness among the participants.

## Conclusions

This study found that higher perceived stress was associated with higher nomophobia, and social support partially mediated the relationship between perceived stress and nomophobia during the COVID-19 pandemic. The result highlighted the importance of perceived social support



in reducing nomophobia. This study carries clinical implications for the treatment of university students who suffer from nomophobia.

**Funding** This study was funded by the UCSI University Research Excellence and Innovation Grant (REIG-FOSSLA-2020–047).

**Data Availability** The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Declarations

**Competing Interests** The authors have no competing interests to declare that are relevant to the content of this article.

**Ethics Approval** The questionnaire and methodology for this study was approved by the UCSI University Institutional Ethics Committee (IEC-2020-FOSSLA-053).

**Consent to Participate** Informed consent was obtained from all individual participants included in the study.

## References

- Ahmed, S., Pokhrel, N., Roy, S., & Samuel, A. J. (2019). Impact of nomophobia: A nondrug addiction among students of physiotherapy course using an online cross-sectional survey. *Indian Journal of Psychiatry*, 61(1), 77. [https://doi.org/10.4103/psychiatry.IndianJPsychiatry\\_361\\_18](https://doi.org/10.4103/psychiatry.IndianJPsychiatry_361_18)
- Ali, A., Muda, M., Ridzuan, A. R., Nuji, M. N. N., Izzamuddin, M. H. M., & Latiff, D. I. A. (2017). The relationship between phone usage factors and nomophobia. *Advanced Science Letters*, 23(8), 7610–7613. <https://doi.org/10.1166/asl.2017.9534>
- American Psychiatric Association. (1994). *Diagnostic and Statistical Manual of Mental Disorders-IV*. Author.
- Bano, N., Khan, M. A., Asif, U., de Beer, J., & Rawass, H. (2021). Effects of nomophobia on anxiety, stress, and depression among Saudi medical students in Jeddah, Saudi Arabia. *JPMA The Journal of the Pakistan Medical Association*, 71(3), 1–11. <https://doi.org/10.47391/JPMA.983>
- Beard, K. W., & Wolf, E. M. (2001). Modification in the proposed diagnostic criteria for Internet addiction. *Cyberpsychology & Behavior*, 4(3), 377–383. <https://doi.org/10.1089/109493101300210286>
- Bekaroğlu, E., & Yılmaz, T. (2020). Nomophobia: Differential diagnosis and treatment. *Psikiyatride Guncel Yaklasimlar*, 12(1), 131–142. <https://doi.org/10.18863/pgy.528897>
- Bender, M., van Osch, Y., Slegers, W., & Ye, M. (2019). Social support benefits psychological adjustment of international students: Evidence from a meta-analysis. *Journal of Cross-Cultural Psychology*, 50(7), 827–847. <https://doi.org/10.1177/0022022119861151>
- Besthorn, F., Kalomo, E. N., Lightfoot, E., & Liao, M. (2018). The relationship between social support and anxiety amongst children living with HIV in rural northern Namibia. *African Journal of AIDS Research*, 17(4), 293–300. <https://doi.org/10.2989/16085906.2018.1534748>
- Bhattacharya, S., Bashar, M. A., Srivastava, A., & Singh, A. (2019). Nomophobia: No mobile phone phobia. *Journal of Family Medicine and Primary Care*, 8(4), 1297. [https://doi.org/10.4103/jfmpc.jfmpc\\_71\\_19](https://doi.org/10.4103/jfmpc.jfmpc_71_19)
- Bhuvanewari, G., & Emiline Joy, E. (2021). Assess the level of stress, sleep disturbance, depression with nomophobia among undergraduate students. *International Journal of Advanced Psychiatric Nursing*, 3(2), 24–27. <https://www.psychiatricjournal.net/archives/2021.v3.i2.A.59>
- Bonavita, S., Sparaco, M., Russo, A., Borriello, G., & Lavorgna, L. (2021). Perceived stress and social support in a large population of people with multiple sclerosis recruited online through the Covid-19 pandemic. *European Journal of Neurology*, 28(10), 3396–3402. <https://doi.org/10.1111/ene.14697>
- Bragazzi, N. L., & Del Puente, G. (2014). A proposal for including nomophobia in the new DSM-V. *Psychology Research and Behavior Management*, 7, 155–160. <https://doi.org/10.2147/PRBM.S41386>
- Campagne, D. M. (2019). Stress and perceived social isolation (loneliness). *Archives of Gerontology and Geriatrics*, 82, 192–199. <https://doi.org/10.1016/j.archger.2019.02.007>
- Caplan, S. E. (2002). Problematic internet use and psychosocial well-being: Development of a theory-based cognitive-behavioral measurement instrument. *Computers in Human Behavior*, 18(5), 553–575. [https://doi.org/10.1016/S0747-5632\(02\)00004-3](https://doi.org/10.1016/S0747-5632(02)00004-3)
- Chen, X., Zou, Y., & Gao, H. (2021). Role of neighborhood social support in stress coping and psychological wellbeing during the COVID-19 pandemic: Evidence from Hubei China. *Health & Place*, 69, 102532. <https://doi.org/10.1016/j.healthplace.2021.102532>
- Çobanoğlu, A., Bahadır Yılmaz, E., & Kızıltan, B. (2021). The relationship between nursing students' digital and smartphone addiction levels and nomophobia: A descriptive, correlational study. *Perspectives in Psychiatric Care*, 57, 1727–1734. <https://doi.org/10.1111/ppc.12742>
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385–396. <https://doi.org/10.2307/2136404>
- Cohen, S., & Wills, T. A. (1985). Stress, social support, and the buffering hypothesis. *Psychological Bulletin*, 98(2), 310. <https://doi.org/10.1037/0033-2909.98.2.310>
- Dasgupta, P., Bhattacharjee, S., Dasgupta, S., Roy, J. K., Mukherjee, A., & Biswas, R. (2017). Nomophobic behaviors among smartphone using medical and engineering students in two colleges of West Bengal. *Indian Journal of Public Health*, 61(3), 199–204.
- Dixit, S., Shukla, H., Bhagwat, A., Bindal, A., Goyal, A., Zaidi, A., & Shrivastava, A. (2010). A study to evaluate mobile phone dependence among students of a medical college and associated hospital of central India. *Indian Journal of Community Medicine*, 35(2), 339–341. <https://doi.org/10.4103/0970-0218.66878>
- Ergin, Z. O., & Ozer, A. (2021). Unravelling the relation between fear of missing out, time spent on the phone, sex, alienation, and nomophobia. *Psychological Reports*, 00332941211043456. <https://doi.org/10.1177/00332941211043456>
- Farchakh, Y., Hallit, R., Akel, M., Chalhoub, C., Hachem, M., Hallit, S., & Obeid, S. (2021). Nomophobia in Lebanon: Scale validation and association with psychological aspects. *PLoS ONE*, 16(4), e0249890. <https://doi.org/10.1371/journal.pone.0249890>
- Gezgin, D. M., Cakir, O., & Yildirim, S. (2018). The relationship between levels of nomophobia prevalence and internet addiction among high school students: The factors influencing nomophobia. *International Journal of Research in Education and Science*, 4(1), 215–225.
- Gökçearslan, Ş., Uluyol, Ç., & Şahin, S. (2018). Smartphone addiction, cyberloafing, stress and social support among university students: A path analysis. *Children and Youth Services Review*, 91, 47–54. <https://doi.org/10.1016/j.childyouth.2018.05.036>
- Gold, A. H., Malhotra, A., & Segars, A. H. (2001). Knowledge management: An organizational capabilities perspective. *Journal of Management Information Systems*, 18(1), 185–214. <https://doi.org/10.1080/07421222.2001.11045669>

- Guo, K., Zhang, X., Bai, S., Minhat, H. S., Nazan, A. I. N. M., Feng, J., Li, X., Luo, G., Zhang, X., Feng, J., Li, Y., Si, M., Qiao, Y., Ouyang, J., & Saliluddin, S. (2021). Assessing social support impact on depression, anxiety, and stress among undergraduate students in Shaanxi province during the COVID-19 pandemic of China. *PLoS ONE*, *16*(7), e0253891. <https://doi.org/10.1371/journal.pone.0253891>
- Gurbuz, I. B., & Ozkan, G. (2020). What is your level of Nomophobia? An investigation of prevalence and level of nomophobia among young people in Turkey. *Community Mental Health Journal*, *56*(5), 814–822. <https://doi.org/10.1007/s10597-019-00541-2>
- Hair, J. F., Jr., Sarstedt, M., Ringle, C. M., & Gudergan, S. P. (2017). *Advanced issues in partial least squares structural equation modeling*. SAGE Publications.
- Henseler, J., Dijkstra, T. K., Sarstedt, M., Ringle, C. M., Diamantopoulos, A., Straub, D. W., Ketchen, D. J., Hair, J. F., Hult, G. T. M., & Calantone, R. J. (2014). Common beliefs and reality about PLS: Comments on Rönkkö and Evermann (2013). *Organizational Research Methods*, *17*(2), 182–209. <https://doi.org/10.1177/1094428114526928>
- Henseler, J., Ringle, C. M., & Sarstedt, M. (2015). A new criterion for assessing discriminant validity in variance-based structural equation modeling. *Journal of the Academy of Marketing Science*, *43*(1), 115–135. <https://doi.org/10.1007/s11747-014-0403-8>
- Herrero, J., Uruña, A., Torres, A., & Hidalgo, A. (2019). Socially connected but still isolated: Smartphone addiction decreases social support over time. *Social Science Computer Review*, *37*(1), 73–88. <https://doi.org/10.1177/0894439317742611>
- Ho, R. C., Zhang, M. W., Tsang, T. Y., Toh, A. H., Pan, F., Lu, Y., et al. (2014). The association between internet addiction and psychiatric co-morbidity: A meta-analysis. *BMC Psychiatry*, *14*(1), 1–10. <https://doi.org/10.1186/1471-244X-14-183>
- Kara, M., Baytemir, K., & Inceman-Kara, F. (2021). Duration of daily smartphone usage as an antecedent of nomophobia: Exploring multiple mediation of loneliness and anxiety. *Behaviour & Information Technology*, *40*(1), 85–98. <https://doi.org/10.1080/0144929x.2019.1673485>
- Khodarahimi, S., Hashim, I. H., & Zaharim, N. M. (2016). Attachment styles, perceived stress and social support in a Malaysian young adults sample. *Psychologica Belgica*, *56*(1), 65–79. <https://doi.org/10.5334/pb.320>
- King, A. L. S., Valença, A. M., & Nardi, A. E. (2010). Nomophobia: The mobile phone in panic disorder with agoraphobia: Reducing phobias or worsening of dependence? *Cognitive and Behavioral Neurology*, *23*(1), 52–54. <https://doi.org/10.1097/WNN.0b013e3181b7eabc>
- Kock, N. (2015). Common method bias in PLS-SEM: A full collinearity assessment approach. *International Journal of e-Collaboration (IJec)*, *11*(4), 1–10. <https://doi.org/10.4018/ijec.2015100101>
- Lelapary, G. S., Mappedoki, S., & Nurafriani. (2020). Hubungan perilaku nomophobia dengan stress pada siswa SMA Negeri 21 Makassar. *Jurnal Ilmiah Kesehatan Diagnosis*, *15*(2), 189–193.
- Li, C., Jiang, S., Li, N., & Zhang, Q. (2018). Influence of social participation on life satisfaction and depression among Chinese elderly: Social support as a mediator. *Journal of Community Psychology*, *46*(3), 345–355. <https://doi.org/10.1002/jcop.21944>
- Limone, P., & Toto, G. A. (2021). Psychological and emotional effects of digital technology on children in Covid-19 Pandemic. *Brain Sciences*, *11*(9), 1126. <https://doi.org/10.3390/brainsci11091126>
- Meng, S. Q., Cheng, J. L., Li, Y. Y., Yang, X. Q., Zheng, J. W., Chang, X. W., Shi, Y., Chen, Y., Lu, L., Sun, Y., Bao, Y. P., & Shi, J. (2022). Global prevalence of digital addiction in general population: A systematic review and meta-analysis. *Clinical Psychology Review*, 102128. <https://doi.org/10.1016/j.cpr.2022.102128>
- Mir, R., & Akhtar, M. (2020). Effect of nomophobia on the anxiety levels of undergraduate students. *Journal of Pakistan Medical Association*, *70*(1492). <https://doi.org/10.5455/JPMA.31286>
- Nguyen, B. T. N., Nguyen, T. T., & Le, U. T. T. (2022). Nomophobia and stress among Vietnamese high school students in Covid-19 pandemic: A mediation model of loneliness. *Journal of Biochemical Technology*, *13*(1), 34–40. <https://doi.org/10.51847/zn5yxlp0nk>
- Ozdemir, B., Cakir, O., & Hussain, I. (2018). Prevalence of nomophobia among university students: A comparative study of Pakistani and Turkish undergraduate students. *Eurasia Journal of Mathematics, Science and Technology Education*, *14*(4), 1519–32. <https://doi.org/10.29333/ejmste/84839>
- Pavithra, M. B., Madhukumar, S., & Mahadeva, M. (2015). A study on nomophobia-mobile phone dependence, among students of a medical college in Bangalore. *National Journal of Community Medicine*, *6*(3), 340–344.
- Purnama, D. H., & Sununianti, V. V. (2020). Social alienation of students in Palembang. *Advances in Social Science, Education and Humanities Research*, *513*.
- Ramayah, T., Cheah, J., Chuah, F., Ting, H., & Memon, M. A. (2018). *Partial least squares structural equation modeling (PLS-SEM) using smartPLS 3.0: An updated guide and practical guide to statistical analysis* (2nd ed.). Pearson.
- Rodríguez-García, A. M., Moreno-Guerrero, A. J., & Lopez Belmonte, J. (2020). Nomophobia: An individual's growing fear of being without a smartphone—a systematic literature review. *International Journal of Environmental Research and Public Health*, *17*(2), 580. <https://doi.org/10.3390/ijerph17020580>
- Roohafza, H. R., Afshar, H., Keshteli, A. H., Mohammadi, N., Feizi, A., Taslimi, M., & Adibi, P. (2014). What's the role of perceived social support and coping styles in depression and anxiety? *Journal of Research in Medical Sciences: The Official Journal of Isfahan University of Medical Sciences*, *19*(10), 944.
- Roohafza, H., Feizi, A., Afshar, H., Mazaheri, M., Behnamfar, O., Hassanzadeh-Keshteli, A., & Adibi, P. (2016). Path analysis of relationship among personality, perceived stress, coping, social support, and psychological outcomes. *World Journal of Psychiatry*, *6*(2), 248. <https://doi.org/10.5498/wjp.v6.i2.248>
- Rosen, L., Carrier, L. M., Miller, A., Rökkum, J., & Ruiz, A. (2016). Sleeping with technology: Cognitive, affective, and technology usage predictors of sleep problems among college students. *Sleep Health*, *2*(1), 49–56. <https://doi.org/10.1016/j.sleh.2015.11.003>
- Rungtusanatham, M., Miller, J. W., & Boyer, K. K. (2014). Theorizing, testing, and concluding for mediation in SCM research: Tutorial and procedural recommendations. *Journal of Operations Management*, *32*(3), 99–113. <https://doi.org/10.1016/j.jom.2014.01.002>
- Sarstedt, M., Hair, J. F., Jr., Cheah, J. H., Becker, J. M., & Ringle, C. M. (2019). How to specify, estimate, and validate higher-order constructs in PLS-SEM. *Australasian Marketing Journal (AMJ)*, *27*(3), 197–211. <https://doi.org/10.1016/j.ausmj.2019.05.003>
- Selkie, E., Adkins, V., Masters, E., Bajpai, A., & Shumer, D. (2020). Transgender adolescents' uses of social media for social support. *Journal of Adolescent Health*, *66*(3), 275–280. <https://doi.org/10.1016/j.jadohealth.2019.08.011>
- Shirani, A. (2017). On the relationship between loneliness and social support and cell phone addiction among students. *Journal of School Psychology*, *5*(4), 7–30.
- Shmueli, G., Ray, S., Velasquez Estrada, J. M., & Chatla, S. B. (2016). The elephant in the room: Predictive performance of PLS models. *Journal of Business Research*, *69*(10), 4552–4564. <https://doi.org/10.1016/j.jbusres.2016.03.049>
- Soper, D. S. (2021). *A-priori Sample Size Calculator for Structural Equation Models [Software]*. <https://www.danielsoper.com/statcalc>

- Szkody, E., Stearns, M., Stanhope, L., & McKinney, C. (2021). Stress-buffering role of social support during COVID-19. *Family Process, 60*(3), 1002–1015. <https://doi.org/10.1111/famp.12618>
- Tams, S., Legoux, R., & Léger, P. M. (2018). Smartphone withdrawal creates stress: A moderated mediation model of nomophobia, social threat, and phone withdrawal context. *Computers in Human Behavior, 81*, 1–9. <https://doi.org/10.1016/j.chb.2017.11.026>
- Terzioğlu, M. A., & Uğurlu, T. T. (2021). Perceived stress and nomophobia in medical faculty students during COVID-19 pandemic. *Bağışıklık Dergisi, 22*(4), 474–482. <https://doi.org/10.51982/bagimli.953053>
- Tran, B. X., Huong, L. T., Hinh, N. D., Nguyen, L. H., Le, B. N., Nong, V. M., et al. (2017). A study on the influence of internet addiction and online interpersonal influences on health-related quality of life in young Vietnamese. *BMC Public Health, 17*(1), 138. <https://doi.org/10.1186/s12889-016-3983-z>
- Wang, P., Lei, L., Wang, X., Nie, J., Chu, X., & Jin, S. (2018). The exacerbating role of perceived social support and the “buffering” role of depression in the relation between sensation seeking and adolescent smartphone addiction. *Personality and Individual Differences, 130*, 129–134. <https://doi.org/10.1016/j.paid.2018.04.009>
- Williams, M., & Vogt, W. P. (2011). The SAGE handbook of innovation in social research methods. *SAGE Publications*. <https://doi.org/10.4135/9781446268261>
- Xie, H., Peng, W., Yang, Y., Zhang, D., Sun, Y., Wu, M., Zhang, J., Jia, J., & Su, Y. (2018). Social support as a mediator of physical disability and depressive symptoms in Chinese elderly. *Archives of Psychiatric Nursing, 32*(2), 256–262. <https://doi.org/10.1016/j.apnu.2017.11.012>
- Yildirim, C., & Correia, A. P. (2015). Exploring the dimensions of nomophobia: Development and validation of a self-reported questionnaire. *Computers in Human Behavior, 49*, 130–137. <https://doi.org/10.1016/j.chb.2015.02.059>
- Zhang, H., Zhao, Q., Cao, P., & Ren, G. (2017). Resilience and quality of life: Exploring the mediator role of social support in patients with breast cancer. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research, 23*, 5969–5979. <https://doi.org/10.12659/msm.907730>
- Zhang, K., Kim, K., Silverstein, N. M., Song, Q., & Burr, J. A. (2021). Social media communication and loneliness among older adults: The mediating roles of social support and social contact. *The Gerontologist, 61*(6), 888–896. <https://doi.org/10.1093/geront/gnaa197>
- Zhang, M. W., & Ho, R. (2017). Moodle: The cost effective solution for internet cognitive behavioral therapy (I-CBT) interventions. *Technology and Health Care, 25*(1), 163–165. <https://doi.org/10.3233/THC-161261>
- Zhang, S., Tian, Y., Sui, Y., Zhang, D., Shi, J., Wang, P., Meng, W., & Si, Y. (2018). Relationships between social support, loneliness, and internet addiction in Chinese postsecondary students: A longitudinal cross-lagged analysis. *Frontiers in Psychology, 9*, 1707. <https://doi.org/10.3389/fpsyg.2018.01707>
- Zhou, H. L., Jiang, H. B., Zhang, B., & Liang, H. Y. (2021). Social anxiety, maladaptive cognition, mobile phone addiction, and perceived social support: A moderated mediation model. *Journal of Psychology in Africa, 31*(3), 248–253. <https://doi.org/10.1080/14330237.2021.1927354>
- Zhao, X., Lynch, J. G., Jr., & Chen, Q. (2010). Reconsidering Baron and Kenny: Myths and truths about mediation analysis. *Journal of Consumer Research, 37*(2), 197–206. <https://doi.org/10.1086/651257>
- Zimet, G. D., Dahlem, N. W., Zimet, S. G., & Farley, G. K. (1988). The multidimensional scale of perceived social support. *Journal of Personality Assessment, 52*(1), 30–41. [https://doi.org/10.1207/s15327752jpa5201\\_2](https://doi.org/10.1207/s15327752jpa5201_2)
- Zwilling, M. (2022). The impact of nomophobia, stress, and loneliness on smartphone addiction among young adults during and after the COVID-19 pandemic: An Israeli case analysis. *Sustainability, 14*(6), 3229. <https://doi.org/10.3390/su14063229>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.