Correction to: Effects of eculizumab treatment on quality of life in patients with paroxysmal nocturnal hemoglobinuria in Japan

Yasutaka Ueda^{1,2} · Naoshi Obara^{2,3} · Yuji Yonemura^{2,4} · Hideyoshi Noji^{2,5} · Masayoshi Masuko⁶ · Yoshinobu Seki^{7,8} · Katsuya Wada⁹ · Takahisa Matsuda¹⁰ · Hirozumi Akiyama¹⁰ · Takayuki Ikezoe^{2,5} · Shigeru Chiba^{2,11} · Yoshinobu Kanda^{2,12} · Tatsuya Kawaguchi^{2,13} · Tsutomu Shichishima^{2,5} · Hideki Nakakuma^{2,14} · Shinichiro Okamoto^{2,15} · Jun-ichi Nishimura^{1,2} · Yuzuru Kanakura^{1,2} · Haruhiko Ninomiya^{2,16}

Published online: 6 July 2018
© The Japanese Society of Hematology 2018

Correction to:
International Journal of Hematology (2018) 107:656–665
<https://doi.org/10.1007/s12185-018-2409-3>

In the original publication of this article, Tables 2, 3 and 4 were published incorrectly. The corrected Tables 2, 3 and 4 are given in the following pages.

The original article can be found online at <https://doi.org/10.1007/s12185-018-2409-3>.

✉ Haruhiko Ninomiya
ninomiya.haruhiko.ft@u.tsukuba.ac.jp

¹ Department of Hematology and Oncology, Osaka University Graduate School of Medicine, Suita, Osaka, Japan

² Japan PNH Study Group, Tokyo, Japan

³ Department of Hematology, University of Tsukuba, Tsukuba, Japan

⁴ Department of Transfusion Medicine and Cell Therapy, Kumamoto University Hospital, Kumamoto, Japan

⁵ Department of Hematology, Fukushima Medical University, Fukushima, Japan

⁶ Department of Stem Cell Transplantation, Niigata University Hospital, Niigata, Japan

⁷ Department of Hematology, Niigata Prefectural Shibata Hospital, Shibata, Japan

⁸ Department of Hematology, Uonuma Institute of Community Medicine, Niigata University Medical and Dental Hospital, Minami-Uonuma, Japan

⁹ Department of Hematology, Matsushita Memorial Hospital, Moriguchi, Japan

¹⁰ Alexion Pharma G.K, Tokyo, Japan

¹¹ Department of Hematology, Faculty of Medicine, University of Tsukuba, Tsukuba, Japan

¹² Division of Hematology, Jichi Medical University, Shimotsuke, Japan

¹³ Departments of Hematology and Infectious Diseases, Kumamoto University, Kumamoto, Japan

¹⁴ Kagoshima Tokusyukai Hospital, Kagoshima, Japan

¹⁵ Division of Hematology, Department of Medicine, Keio University School of Medicine, Tokyo, Japan

¹⁶ Department of Medical Sciences, Faculty of Medicine, University of Tsukuba, Tsukuba, Japan

Table 2 Disease characteristics at baseline

	Post-marketing	QOL	BMF (+)	BMF (−)
LDH (U/L)				
<i>n</i>	477	51	30	21
Mean (SD)	1478.3 (776.3)	1435.3 (922.6)	1161.8 (591.4)	1826.0 (1161.2)
Median (min, max)	1340.0 (157, 6082)	1131.0 (447, 6082)	1049.0 (447, 2870)	1638.0 (618, 6082)
Hb (g/dL)				
<i>n</i>	480	52	31	21
Mean (SD)	8.04 (1.70)	8.22 (1.92)	8.17 (2.19)	8.29 (1.47)
Median (min, max)	7.90 (3.8, 15.0)	8.20 (4.5, 15.0)	8.20 (4.5, 15.0)	8.20 (5.7, 11.5)
PLT ($\times 10^3/\mu\text{L}$)				
<i>n</i>	451	51	30	21
Mean (SD)	133.3 (80.6)	129.1 (69.3)	103.0 (59.3)	166.4 (66.6)
Median (min, max)	126.0 (4, 574)	128.0 (12, 281)	93.0 (12, 228)	172.0 (60, 281)
PNH clone size (%)				
<i>n</i>	241	26	14	12
Mean (SD)	42.18 (23.90)	34.94 (26.66)	32.93 (29.62)	37.28 (23.81)
Median (min, max)	40.00 (0.0, 99.0)	23.35 (1.7, 97.0)	19.20 (10.2, 97.0)	36.70 (1.7, 83.5)

PNH clone size (%) was evaluated by erythrocytes

Table 3 Changes in laboratory data from baseline to 1-year

	Baseline	1 year	Paired <i>t</i> test
LDH (U/L) (<i>n</i> = 51)			
Mean (SD)	1435.3 (922.6)	292.2 (170.7)	<i>P</i> < 0.01
Median (min, max)	1131.0 (447, 6082)	242.0 (148, 1045)	
Hb (g/dL) (<i>n</i> = 52)			
Mean (SD)	8.22 (1.92)	9.37 (1.97)	<i>P</i> < 0.01
Median (min, max)	8.20 (4.5, 15.0)	9.35 (5.1, 14.3)	
PLT ($\times 10^3/\mu\text{L}$) (<i>n</i> = 51)			
Mean (SD)	129.1 (69.3)	133.8 (64.8)	N.S.
Median (min, max)	128.0 (12, 281)	137.0 (11, 306)	
eGFR (mL/min/1.73 m ²) (<i>n</i> = 29)			
Mean (SD)	73.75 (37.87)	75.57 (38.47)	N.S.
Median (min, max)	58.60 (27.0, 161.3)	64.80 (25.2, 171.1)	
PNH clone size (%) (<i>n</i> = 5)			
Mean (SD)	28.88 (30.81)	33.60 (16.41)	N.S.
Median (min, max)	18.80 (11.0, 83.5)	23.50 (19.3, 52.0)	

PNH clone size (%) was evaluated by erythrocytes

Table 4 Canonical correlation analysis between changes in laboratory data and changes in QOL scores

	Canonical correlation coefficient
FACIT-fatigue	0.74
EORTC-QLQ-C30	
Global health status/QoL	0.69
Physical functioning	0.72
Role functioning	0.64
Emotional functioning	0.38
Cognitive functioning	0.20
Social functioning	0.59
Fatigue	− 0.87
Nausea and vomiting	− 0.44
Pain	− 0.27
Dyspnea	− 0.70
Insomnia	− 0.58
Appetite loss	− 0.55
Constipation	− 0.12
Diarrhea	− 0.16
Financial difficulties	− 0.51
Laboratory value	
LDH	− 0.91
Hb	0.76
PLT	0.03
Statistics	
Canonical <i>R</i>	0.8393
<i>F</i> value	1.9166
<i>P</i> value	0.0042