Correction to: Comparison of minimal residual disease detection in multiple myeloma by SRL 8-color single-tube and EuroFlow 8-color 2-tube multiparameter flow cytometry

Hiroyuki Takamatsu¹ · Takeshi Yoroidaka¹ · Momoko Fujisawa¹ · Kazuya Kobori² · Masako Hanawa² · Takeshi Yamashita³ · Ryoichi Murata³ · Mikio Ueda³ · Shinji Nakao¹ · Kosei Matsue⁴

Published online: 21 March 2019
© Japanese Society of Hematology 2019

Correction to: International Journal of Hematology
<https://doi.org/10.1007/s12185-019-02615-z>

The authors would like to correct the error in the publication of the original article. The corrected detail is given below for your reading.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements We would like to acknowledge Dr Juan Flores-Montero, Dr Luzalba Sanoja-Flores and Prof Alberto Orfao of Universidad de Salamanca, Dr Brian Durie of International Myeloma Foundation, and Prof Hirokazu Murakami, Dr Kenshi Suzuki and Dr Kazuyuki Shimizu of Japanese Society of Myeloma for the EuroFlow-NGF assessments.

The original article can be found online at <https://doi.org/10.1007/s12185-019-02615-z>.

✉ Hiroyuki Takamatsu
takamaz@staff.kanazawa-u.ac.jp

- ¹ Department of Hematology and Respiriology, Faculty of Medicine, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, 13-1 Takaramachi, Kanazawa, Ishikawa 920-8641, Japan
- ² Cellular Immunology Analysis Section, Genetic and Chromosome Analysis Department, SRL, Inc, Tokyo, Japan
- ³ Division of Internal Medicine, Keiju Kanazawa Hospital, Kanazawa, Japan
- ⁴ Department of Hematology/Oncology, Kameda Medical Center, Kamogawa, Japan