Correction to: Treatment of aggressive adult T-cell leukemia/lymphoma: a retrospective study in a hospital located in HTLV-1 highly endemic area

Daisuke Nakamura^{1,2} · Makoto Yoshimitsu² · Tomohisa Tabuchi^{1,2} · Naosuke Arima^{1,2} · Maiko Hayashida^{1,2} · Hirosaka Inoue^{1,2} · Kakushi Matsushita^{1,2} · Tadashi Matsumoto¹ · Naomichi Arima² · Kenji Ishitsuka²

Published online: 6 March 2020
© Japanese Society of Hematology 2020

Correction to:
International Journal of Hematology (2020) 111:234–240
<https://doi.org/10.1007/s12185-019-02769-w>

In the original publication of the article, Table 5 was published with incorrect contents. The correct Table 5 is given in this correction.

Table 5 Relative dose intensity (RDI) of younger and elderly patients treated by OPEC/MPEC or VCAP-AMP-VECP chemotherapy

	RDI (%) ± SD	<i>p</i> value
Patients aged < 70 years		
OPEC/MPEC (<i>n</i> = 58)	89.5 ± 17.1	0.175
VCAP-AMP-VECP (<i>n</i> = 20)	83.7 ± 12.9	
Patients aged ≥ 70 years		
OPEC/MPEC (<i>n</i> = 80)	88.6 ± 18	0.002
VCAP-AMP-VECP (<i>n</i> = 13)	71.6 ± 14.8	

The original article can be found online at <https://doi.org/10.1007/s12185-019-02769-w>.

✉ Daisuke Nakamura
d5039@m3.kufm.kagoshima-u.ac.jp
Makoto Yoshimitsu
myoshimi@m.kufm.kagoshima-u.ac.jp
Tomohisa Tabuchi
tomohisatabuchi1116@gmail.com
Naosuke Arima
aritouma@m2.kufm.kagoshima-u.ac.jp
Maiko Hayashida
smarf@m3.kufm.kagoshima-u.ac.jp
Hirosaka Inoue
hirsino@gmail.com
Kakushi Matsushita
kks.matsushita@jiaikai.jp

Tadashi Matsumoto
tmatsumoto@jiaikai.jp
Naomichi Arima
nao@m2.kufm.kagoshima-u.ac.jp
Kenji Ishitsuka
kenji-i@m.kufm.kagoshima-u.ac.jp

¹ Department of Hematology, Izuro Imamura Hospital, 17-1 Horie, Kagoshima, Japan

² Department of Hematology and Rheumatology, Kagoshima University Hospital, 8-35-1 Sakuragaoka, Kagoshima, Japan