CORRECTION



Correction to: Intravenous ferric derisomaltose versus saccharated ferric oxide for iron deficiency anemia associated with menorrhagia: a randomized, open-label, active-controlled, noninferiority study

Hiroshi Kawabata 10 · Takeshi Tamura 2 · Soichiro Tamai 3 · Akiko Fujibayashi 2 · Motoi Sugimura 4 · Study Group

Published online: 8 November 2022 © The Author(s) 2022

Correction to:

International Journal of Hematology (2022) 116:647-658 https://doi.org/10.1007/s12185-022-03401-0

In the original publication of the article, the Fig. 3 was published with errors. The correct Fig. 3 is given in this correction.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long

as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at https://doi.org/10.1007/s12185-022-03401-0.

- ☐ Hiroshi Kawabata hkawabat@kuhp.kyoto-u.ac.jp
- Department of Hematology, National Hospital Organization Kyoto Medical Center, Kyoto, Japan
- ² Clinical Development Department, Nippon Shinyaku Co., Ltd, Kyoto, Japan
- Data Science Department, Nippon Shinyaku Co., Ltd, Kyoto, Janan
- Department of Obstetrics, Gynecology and Family Medicine, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan






Fig. 3 Laboratory value changes over the study period (mean \pm SD) in (a) hemoglobin concentration (g/dL), (b) mean corpuscular volume (fL), (c) reticulocyte ratio (% $_{e}$), (d) transferrin saturation (%),

(e) serum ferritin concentration (ng/mL), and (f) serum hepcidin concentration (ng/mL). *FDI* ferric derisomaltose, *SD* standard deviation, *SFO* saccharated ferric oxide.

