



A rapidly converging domain decomposition algorithm for a time delayed parabolic problem with mixed type boundary conditions exhibiting boundary layers

Aakansha¹ · Sunil Kumar¹ · Higinio Ramos^{2,3} 

Received: 1 November 2023 / Revised: 28 December 2023 / Accepted: 1 January 2024 /

Published online: 2 February 2024

© The Author(s) 2024

Abstract

A rapidly converging domain decomposition algorithm is introduced for a time delayed parabolic problem with mixed type boundary conditions exhibiting boundary layers. Firstly, a space-time decomposition of the original problem is considered. Subsequently, an iterative process is proposed, wherein the exchange of information to neighboring subdomains is accomplished through the utilization of piecewise-linear interpolation. It is shown that the algorithm provides uniformly convergent numerical approximations to the solution. Our analysis utilizes some novel auxiliary problems, barrier functions, and a new maximum principle result. More importantly, we showed that only one iteration is needed for small values of the perturbation parameter. Some numerical results supporting the theory and demonstrating the effectiveness of the algorithm are presented.

Keywords Delay differential equations · Singularly perturbed problems · Uniformly convergent methods · Domain decomposition methods · Parabolic problems

Mathematics Subject Classification 65M06 · 65M55 · 65M15 · 65M12

✉ Higinio Ramos
higra@usal.es

Aakansha
aakansha.chauhan03@gmail.com

Sunil Kumar
skumar.mat@iitbhu.ac.in

¹ Department of Mathematical Sciences, Indian Institute of Technology (BHU) Varanasi, Varanasi, India

² Scientific Computing Group, Universidad de Salamanca, Plaza de la Merced, 37008 Salamanca, Spain

³ Escuela Politécnica Superior de Zamora, Campus Viriato, 49029 Zamora, Spain

1 Introduction

In the mathematical modeling of various physical phenomena in bioscience and control theory, delay differential equations arise frequently, for instance, in population dynamics [12], immunology [5], physiology [6], and neural networks [11]. The presence of the delay term in these models is due to feedback control because a finite time is required to sense and react to the data like the immune period, the time between a cell is infected and the new virus is produced, the stages of the life cycle, the duration of the infectious period, etc. [28]. In delay models, the evolution of the solution and its derivatives involve the past history also. Singularly perturbed delay differential equations form a significant category among these equations. When the highest order derivative term is multiplied with a small positive parameter, the delay differential equations are called as singularly perturbed delay differential equations. Some applications include modeling of the human pupil-light reflex [17], variational problems in control theory [8], the study of bistable devices [7], and chemical processes [2]. The following Hutchinson’s equation [3]

$$u_t(y, t) = \epsilon u_{yy}(y, t) + ru(y, t)[1 - u(y, t - \tau)], \tag{1}$$

is a singularly perturbed differential equation in mathematical ecology, where τ represents the time lag and ϵ is the perturbation parameter. This equation serves as a fundamental model for the evolution of a population with density $u(y, t)$. The following singularly perturbed partial differential equation is used to model the processing of metal sheets [3]

$$u_t = \epsilon u_{yy} + v(f(u(y, t - \tau)))u_y + a[g(u(y, t - \tau)) - u(y, t)], \tag{2}$$

where u denotes the temperature distribution in the metal sheet with velocity v and heated by a source g . The fixed delay of length τ is introduced by the controller’s speed.

Consider the singularly perturbed time delayed parabolic problem

$$\mathcal{L}u(y, t) := Lu(y, t) + b(y, t)u(y, t - \tau) = f(y, t), \quad (y, t) \in \mathcal{Q}, \tag{3}$$

with initial condition

$$u(y, t) = g_b(y, t), \quad (y, t) \in \gamma_b = [0, 1] \times [-\tau, 0],$$

and mixed boundary conditions

$$\begin{cases} \Gamma_\ell u(y, t) := \alpha_1 u(y, t) - \delta_1 \sqrt{\epsilon} u_y(y, t) = g_\ell(t), \text{ on } \gamma_\ell = \{(0, t) : 0 \leq t \leq T\}, \\ \Gamma_r u(y, t) := \alpha_2 u(y, t) + \delta_2 \sqrt{\epsilon} u_y(y, t) = g_r(t), \text{ on } \gamma_r = \{(1, t) : 0 \leq t \leq T\}, \end{cases} \tag{4}$$

where

$$Lu(y, t) = u_t(y, t) - \epsilon u_{yy}(y, t) + a(y, t)u(y, t).$$

We define $\mathcal{Q} := \mathcal{D} \times (0, T]$, $\mathcal{D} = (0, 1)$. Suppose $\gamma = \gamma_\ell \cup \gamma_b \cup \gamma_r$ and $\bar{\gamma} = \gamma_\ell \cup \gamma_o \cup \gamma_r$, where $\gamma_o = [0, 1] \times \{0\}$. Also, assume that $0 < \epsilon \leq 1$, $a(y, t) + b(y, t) \geq \beta > 0$, with $0 < \alpha \leq a(y, t)$, $b(y, t) \leq 0$, and $\tau > 0$. Further, the terminal time T is assumed to be $T = \sigma\tau$, for some integer $\sigma > 1$. We assume that the coefficients of the mixed boundary conditions verify $\alpha_1 > 0$, $\delta_1 \geq 0$, $\alpha_1 \geq \delta_1$, and $\alpha_2, \delta_2 \geq 0$ with $\alpha_2 + \delta_2 > 0$. The data of problem (3)-(4) is supposed to be enough smooth and satisfy appropriate compatibility conditions [31]. With these assumptions, there is a unique solution of (3)-(4) exhibiting boundary layers near $y = 0$ and $y = 1$.

It is well known that the classical discretization methods are unable to provide satisfactory results for problems with boundary layers, as they require a large number of mesh points to resolve the boundary layers and leads to a high computational cost. To address this issue, a variety of robust/uniform numerical methods have been considered in the literature, see, e.g. [3, 4, 9, 13, 32] for delay differential equations with Dirichlet boundary conditions and [29, 31] for delay differential equations with Robin boundary conditions. Additionally, some new developments addressing singularly perturbed delay differential equations can be observed in [19–25].

Domain decomposition-based numerical methods have gained significant popularity for solving partial differential equations. Schwarz's groundbreaking study on domain decomposition in 1870 [30] is acknowledged as the initial work in this field. Domain decomposition enables the application of distinct treatments to each subdomain, allowing the use of different discretization techniques. This approach enables localized handling of singularities within specific subdomains. Additionally, domain decomposition methods offer the advantage of avoiding non-uniform meshes. Furthermore, these methods can be effectively implemented in parallel to enhance the computational efficiency of numerical computations. Most work on domain decomposition for singularly perturbed partial differential equations is restricted to Dirichlet boundary conditions (see [1, 14, 15, 27, 32] and the references therein). For an in-depth exploration of domain decomposition methods, it is recommended to refer to [26, 33]. The existence of a time delay in the differential model and the inclusion of mixed boundary conditions do, indeed, set the theoretical aspect apart. Consequently, problem (3)-(4) has received less attention in the literature.

As far as we are aware, there is no study on domain decomposition for singularly perturbed partial differential equations with a time delay and mixed boundary conditions. So, the purpose of this paper is two-fold: firstly, to develop a novel domain decomposition method for solving problem (3)-(4), and secondly, to provide a comprehensive error analysis of the developed method. We split the domain into three overlapping subdomains (a regular subdomain and two layer subdomains) with the help of a subdomain parameter. Subsequently, an iterative process is proposed, wherein the exchange of information to neighboring subdomains is accomplished through the utilization of piecewise-linear interpolation. It is shown that the algorithm provides uniformly convergent numerical approximations to the solution. More importantly, we showed that only one iteration is needed for very small values of the perturbation parameter ϵ . Numerical results are presented that support the theory and demonstrate the effectiveness of the algorithm. The current research introduces several novel aspects in terms of analysis. Our analysis utilizes some novel auxiliary problems, barrier functions, and a new maximum principle result.

The work is arranged as follows. The continuous maximum principle and the bounds of the derivatives of the solution are discussed in Sect. 2. In Sect. 3 we introduce an algorithm which is analyzed in Sect. 4. Numerical results are presented in Sect. 5 for two test problems and some concluding remarks are included in Sect. 6.

Notation: C is assumed to be a positive constant independent of the discretization parameters $N_y, \Delta t$, the iteration parameter k and the perturbation parameter ϵ . We define a function $h_{i,n} = h(y_i, t_n)$, for $h \in C([0, 1] \times [0, T])$. The norm $\|\cdot\|_{\mathcal{D}, N_y, M_t}$ is defined as discrete maximum norm corresponding to the maximum norm $\|\cdot\|_{\mathcal{D}}$ on a bounded and closed subset \mathcal{D} of $[0, 1] \times [0, T]$. And for a mesh function Z , $\mathcal{I}_n Z$ denotes a piecewise linear interpolation at time t_n .

2 A priori estimates

We obtain some bounds on the derivatives of the solution of (3)-(4). For this purpose, first we introduce a continuous maximum principle for the operator \mathcal{L} .

Lemma 1 *Let w be a sufficiently differentiable function in $\overline{\mathcal{Q}}$. Suppose $\Gamma_\ell w(0, t) \geq 0$, $\Gamma_r w(1, t) \geq 0$ on $(0, T]$ and $w(y, t) \geq 0$ on γ_b . Then $Lw(y, t) \geq 0$ on \mathcal{Q} implies that $w(y, t) \geq 0$ on $\overline{\mathcal{Q}}$.*

Proof The proof can be obtained using the arguments in [21, 31]. □

Lemma 2 *Suppose z is a sufficiently differentiable function satisfying $\Gamma_\ell z(0, t) \geq 0$, $\Gamma_r z(1, t) \geq 0$ on $(0, T]$ and $z(y, t) \geq 0$ on γ_b . Then $\mathcal{L}z(y, t) \geq 0$ on \mathcal{Q} implies that $z(y, t) \geq 0$ on $\overline{\mathcal{Q}}$.*

Proof We assume that $w = z$ for $(y, t) \in \overline{\mathcal{D}} \times [-\tau, \tau]$. Therefore, we have $w(0, t) \geq 0$, $w(1, t) \geq 0$ on $(0, \tau]$, and $w(y, t) \geq 0$ on γ_b . Further,

$$Lw(y, t) \geq -b(y, t)w(y, t - \tau) \geq 0, \quad (y, t) \in \mathcal{D} \times (0, \tau],$$

as $b \leq 0$ and $w \geq 0$ for $(y, t) \in \overline{\mathcal{D}} \times [-\tau, 0]$. Therefore, $z = w \geq 0$ for $(y, t) \in \overline{\mathcal{D}} \times [0, \tau]$ by Lemma 1. Repeating the above arguments and using the fact that $z \geq 0$ for $(y, t) \in \overline{\mathcal{D}} \times [(n - 1)\tau, n\tau]$, we can show that $z \geq 0, \forall (y, t) \in \overline{\mathcal{D}} \times [n\tau, (n + 1)\tau], n \geq 1$. □

Using the aforementioned maximum principle, we can readily derive the following stability estimate.

Lemma 3 *The solution u of problem (3)-(4) satisfies*

$$\|u\|_{\overline{\mathcal{Q}}} \leq \max \left\{ \frac{1}{\alpha} \|Lu\|_{\mathcal{Q}}, \|\Gamma_\ell u\|_{\gamma_\ell}, \|\Gamma_r u\|_{\gamma_r}, \|u\|_{\gamma_b} \right\}. \tag{5}$$

Furthermore, the continuous solution u of (3)-(4) satisfies the following estimates.

Lemma 4 Suppose $a, b, f \in C^{(2+\mu, 1+\mu/2)}(\bar{\mathcal{Q}})$, $g_\ell, g_r \in C^{\frac{3+\mu}{2}}([0, T])$, $g_b \in C^{(4+\mu, 2+\mu/2)}(\gamma_b)$, $\mu \in (0, 1)$. If the compatibility conditions are satisfied up to the second order, then there exists a unique solution to problem (3)-(4) such that $u \in C^{(4+\mu, 2+\mu/2)}(\bar{\mathcal{Q}})$. Furthermore, it is true that

$$\left\| \frac{\partial^{s_1+s_2} u}{\partial y^{s_1} \partial t^{s_2}} \right\|_{\bar{\mathcal{Q}}} \leq C \epsilon^{-s_1/2} \text{ for } 0 \leq s_1 + 2s_2 \leq 4. \tag{6}$$

Proof The first part is proved by using [16]. Now, by using the transformation $\tilde{y} = y/\sqrt{\epsilon}$, we transform problem (3)-(4) as follows

$$\begin{aligned} \tilde{u}_t - \epsilon \tilde{u}_{\tilde{y}\tilde{y}} + \tilde{a}\tilde{u} &= -\tilde{b}\tilde{u}(\tilde{y}, t - \tau) + \tilde{f}, & \text{in } \mathcal{Q}_\epsilon, \\ \Gamma_\ell \tilde{u}(\tilde{y}, t) &= g_\ell(t), & \text{in } \gamma_{\ell, \epsilon}, \\ \Gamma_r \tilde{u}(\tilde{y}, t) &= g_r(t), & \text{in } \gamma_{r, \epsilon}, \\ \tilde{u}(\tilde{y}, t) &= g_{b, \epsilon}(\tilde{y}, t), & \text{in } \gamma_{b, \epsilon}, \end{aligned} \tag{7}$$

where $\mathcal{Q}_\epsilon = (0, 1/\sqrt{\epsilon}) \times (0, T]$, $\gamma_{\ell, \epsilon} = \gamma_\ell$, $\gamma_{r, \epsilon} = \{1/\sqrt{\epsilon}, t) : t \in (0, T]\}$, $\gamma_{b, \epsilon} = [0, 1/\sqrt{\epsilon}] \times [-\tau, 0]$. Now, we use the result from [16, p. 352] to obtain

$$\left\| \frac{\partial^{s_1+s_2} \tilde{u}}{\partial \tilde{y}^{s_1} \partial t^{s_2}} \right\|_{\tilde{D}_\delta} \leq C(1 + \|\tilde{u}\|_{\tilde{D}_{2\delta}}),$$

for all \tilde{D}_δ in $\tilde{\mathcal{Q}}_\epsilon$. Here, $0 \leq s_1 + 2s_2 \leq 4$, $s_1, s_2 \geq 0$, and \tilde{D}_δ is a δ -neighbourhood in $\tilde{\mathcal{Q}}_\epsilon$ of diameter δ .

Transitioning back to the original variable we get

$$\left\| \frac{\partial^{s_1+s_2} u}{\partial y^{s_1} \partial t^{s_2}} \right\|_{\bar{\mathcal{Q}}} \leq C \epsilon^{-s_1/2} (1 + \|u\|_{\bar{\mathcal{Q}}}), \quad 0 \leq s_1 + 2s_2 \leq 4, \quad s_1, s_2 \geq 0.$$

Therefore, the proof is completed using Lemma 3. □

A further decomposition of the solution u is required to prove the convergence analysis of the numerical algorithm. The actual solution u is decomposed as $u = v + w$. The component v is further decomposed as $v = v_0 + \epsilon v_1$, where v_0 and v_1 are the solutions of the problems

$$\begin{aligned} \partial_t v_0 + a v_0 &= -b v_0(y, t - \tau) + f, & \text{in } \mathcal{Q}, \\ v_0 &= g_b, & \text{in } \gamma_b \end{aligned} \tag{8}$$

and

$$\begin{aligned} L v_1 &= -b v_1(y, t - \tau) + \frac{\partial^2 v_0}{\partial x^2}, & \text{in } \mathcal{Q}, \\ \Gamma_\ell v_1 &= 0, & \text{in } \gamma_\ell, \quad \Gamma_r v_1 = 0, & \text{in } \gamma_r, \\ v &= 0, & \text{in } \gamma_b. \end{aligned} \tag{9}$$

Thus, the component v solves

$$\begin{aligned}Lv &= -bv(y, t - \tau) + f, \quad \text{in } \mathcal{Q}, \\ \Gamma_\ell v &= \Gamma_\ell v_0, \quad \text{in } \gamma_\ell, \quad \Gamma_r v = \Gamma_r v_0, \quad \text{in } \gamma_r, \\ v &= g_b, \quad \text{in } \gamma_b.\end{aligned}$$

The component w solves

$$\begin{aligned}Lw &= -bw(y, t - \tau), \quad \text{in } \mathcal{Q}, \\ \Gamma_\ell w &= g_\ell - \Gamma_\ell v_0, \quad \text{in } \gamma_\ell \quad \Gamma_r w = g_r - \Gamma_r v_0, \quad \text{in } \gamma_r, \\ w &= 0, \quad \text{in } \gamma_b.\end{aligned} \tag{10}$$

Lemma 5 Suppose $a, b, f \in C^{(4+\mu, 2+\mu/2)}(\overline{\mathcal{Q}})$, $g_\ell, g_r \in C^{\frac{5+\mu}{2}}([0, T])$, $g_b \in C^{(6+\mu, 3+\mu/2)}(\gamma_b)$, $\mu \in (0, 1)$, and appropriate compatibility conditions hold. Then, one can split u as $u = v + w$, where v and w are the regular and singular components, respectively, satisfying

$$\left\| \frac{\partial^{s_1+s_2} v}{\partial y^{s_1} \partial t^{s_2}} \right\|_{\overline{\mathcal{Q}}} \leq C(1 + \epsilon^{(2-s_1)/2}), \tag{11}$$

$$\left| \frac{\partial^{s_1+s_2} w}{\partial y^{s_1} \partial t^{s_2}} \right| \leq C\epsilon^{-s_1/2} \left(e^{-y\sqrt{1/\epsilon}} + e^{-(1-y)\sqrt{1/\epsilon}} \right), \tag{12}$$

for $(y, t) \in \overline{\mathcal{Q}}$, $0 \leq s_1 + 2s_2 \leq 4$, $s_1, s_2 \geq 0$.

Proof The bounds are derived as follows. Note that v_0 is the solution of problem (8), which is a problem that does not involve ϵ . Hence, using a classical argument, we get

$$\left\| \frac{\partial^{s_1+s_2} v_0}{\partial y^{s_1} \partial t^{s_2}} \right\|_{\overline{\mathcal{Q}}} \leq C.$$

Moreover, v_1 is the solution to a problem in a form for which Lemma 4 is applicable. Consequently, it holds

$$\left\| \frac{\partial^{s_1+s_2} v_1}{\partial y^{s_1} \partial t^{s_2}} \right\|_{\overline{\mathcal{Q}}} \leq C\epsilon^{-s_1/2}.$$

Since $v = v_0 + \epsilon v_1$, the necessary bounds for the component v and its derivatives follow from the above two bounds. The bounds for the component w and its derivatives can be proved using the arguments in [10]. \square

3 The numerical algorithm

The computational domain \mathcal{Q} is divided into three overlapping subdomains \mathcal{Q}_p , $p = \ell, m, r$, where $\mathcal{Q}_\ell = (0, 2\rho) \times (0, T]$ and $\mathcal{Q}_r = (1 - 2\rho, 1) \times (0, T]$ are the layer subdomains, and $\mathcal{Q}_m = (\rho, 1 - \rho) \times (0, T]$ is the regular subdomain with

$$\rho = \min \left\{ \frac{1}{4}, 2\sqrt{\frac{\epsilon}{\beta}} \ln N_y \right\}. \tag{13}$$

Here, ρ is the subdomain parameter, which is chosen as the Shishkin transition point. Each subdomain $\mathcal{Q}_p = \mathcal{D}_p \times (0, T]$, where $\mathcal{D}_p = (c, d)$, is discretized using a rectangular mesh which is formed by uniform meshes in both directions. We define $\gamma_{b,p} = \overline{\mathcal{D}}_p \times \overline{\Omega}_o$, where $\overline{\Omega}_o = [-\tau, 0]$. The uniform mesh $\overline{\mathcal{D}}_p^{N_y}$ in the spatial direction is defined as $\overline{\mathcal{D}}_p^{N_y} = \{y_i = c + ih_p, h_p = (d - c)/N_y\}_{i=0}^{N_y}$, while $\overline{\Omega}_o^{m_\tau}, \overline{\Omega}^{M_t}$ are the meshes in time direction with uniform length $\Delta t = T/M_t$, which are obtained by dividing $[-\tau, 0]$ and $[0, T]$ in a number of subintervals m_τ and $M_t = \sigma m_\tau$, respectively. Defining $\mathcal{D}_p^{N_y} = \overline{\mathcal{D}}_p^{N_y} \cap \mathcal{D}_p$ and $\Omega^{M_t} = \overline{\Omega}^{M_t} \cap (0, T]$, then on each subdomain, the meshes $\mathcal{Q}_p^{N_y, M_t}$ and $\gamma_{b,p}^{N_y, m_\tau}$ are defined as $\mathcal{Q}_p^{N_y, M_t} = \mathcal{D}_p^{N_y} \times \Omega^{M_t}$ and $\gamma_{b,p}^{N_y, m_\tau} = \overline{\mathcal{D}}_p^{N_y} \times \overline{\Omega}_o^{m_\tau}$ where N_y and M_t are the discretization parameters in space and time directions respectively.

On each subdomain $\mathcal{Q}_p^{N_y, M_t}$, $p = \ell, m, r$, the backward Euler scheme and central differencing are employed to discretize in time and space respectively. Thus, we have

$$L_p^{N_y, M_t} \mathcal{U}_p(y_i, t_n) + b(y_i, t_n) \mathcal{U}_p(y_i, t_{n-m_\tau}) = f(y_i, t_n), \tag{14}$$

where

$$L_p^{N_y, M_t} \mathcal{U}_p(y_i, t_n) = F_t^- \mathcal{U}_p(y_i, t_n) - \epsilon D_y \mathcal{U}_p(y_i, t_n) + a(y_i, t_n) \mathcal{U}_p(y_i, t_n).$$

The boundary conditions are discretized with a special second order scheme as follows:

$$\begin{cases} \Gamma_\ell^{N_y, M_t} \mathcal{U}_p(0, t_n) \equiv \alpha_1 \mathcal{U}(0, t_n) - \delta_1 \sqrt{\epsilon} F_y^+ \mathcal{U}(0, t_n) + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (a \mathcal{U} + F_t^- \mathcal{U})(0, t_n) \\ \quad = g_\ell(t_n) + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (-b(0, t_n) \mathcal{U}(0, t_{n-m_\tau}) + f(0, t_n)), \\ \Gamma_r^{N_y, M_t} \mathcal{U}_p(1, t_n) \equiv \alpha_2 \mathcal{U}(1, t_n) + \delta_2 \sqrt{\epsilon} F_y^- \mathcal{U}(1, t_n) + \frac{h_r \delta_2}{2\sqrt{\epsilon}} (a \mathcal{U} + F_t^- \mathcal{U})(1, t_n) \\ \quad = g_r(t_n) + \frac{h_r \delta_2}{2\sqrt{\epsilon}} (-b(1, t_n) \mathcal{U}(1, t_{n-m_\tau}) + f(1, t_n)), \\ \mathcal{U}_p(y_i, t_n) = g_b(y_i, t_n). \end{cases} \tag{15}$$

Here,

$$[D_y \mathcal{Y}]_{i,n} = \frac{\mathcal{Y}_{i+1,n} - 2\mathcal{Y}_{i,n} + \mathcal{Y}_{i-1,n}}{h_p^2},$$

$$[F_t^- \mathcal{Y}]_{i,n} := \frac{\mathcal{Y}_{i,n} - \mathcal{Y}_{i,n-1}}{\Delta t}, [F_y^+ \mathcal{Y}]_{i,n} := \frac{\mathcal{Y}_{i+1,n} - \mathcal{Y}_{i,n}}{h_p}, [F_y^- \mathcal{Y}]_{i,n} := \frac{\mathcal{Y}_{i,n} - \mathcal{Y}_{i-1,n}}{h_p}.$$

In this approach, we discretized the boundary conditions using a combination of spatial and temporal discretizations. It is important to observe that employing the typical upwind discretization for mixed boundary conditions results in linear accuracy. However, with the proposed methodology, we have the possibility to increase this accuracy to a quadratic order by refining the truncation error in both space and time.

Setting $\overline{\mathcal{Q}}^{N_y, M_t} := (\overline{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m) \cup \overline{\mathcal{Q}}_m^{N_y, M_t} \cup (\overline{\mathcal{Q}}_r^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m)$, the numerical solution of (3)-(4) is computed using the step by step process defined as follows. Suppose the initial approximation $\mathcal{U}^{[0]}$ is

$$\mathcal{U}^{[0]}(y_i, t_n) = \begin{cases} 0, & (y_i, t_n) \in (0, 1) \times (0, T], \\ u(y_i, t_n), & (y_i, t_n) \in [0, 1] \times [-\tau, 0], \end{cases}$$

and compute $\mathcal{U}_p^{[k]}$, $p = \ell, m, r$, for $k \geq 1$, by solving the following set of equations

$$\begin{cases} [L_\ell^{N_y, M_t} \mathcal{U}_\ell^{[k]}]_{i,n} + b_{i,n} \mathcal{U}_{\ell; i, n - m_\tau}^{[k]} = f_{i,n}, & (y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t}, \\ \mathcal{U}_\ell^{[k]}(y_i, t_n) = g_b(y_i, t_n), & (y_i, t_n) \in \gamma_{b, \ell}^{N_y, m_\tau}, \\ \Gamma_\ell^{N_y, M_t} \mathcal{U}_\ell^{[k]}(0, t_n) \\ = g_\ell(t_n) + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (-b(0, t_n) \mathcal{U}(0, t_n - m_\tau) + f(0, t_n)), & t_n \in \Omega^{M_t}, \\ \mathcal{U}_\ell^{[k]}(2\rho, t_n) = \mathcal{I}_n \mathcal{U}^{[k-1]}(2\rho, t_n), & t_n \in \Omega^{M_t}, \end{cases} \tag{16}$$

$$\begin{cases} [L_r^{N_y, M_t} \mathcal{U}_r^{[k]}]_{i,n} + b_{i,n} \mathcal{U}_{r; i, n - m_\tau}^{[k]} = f_{i,n}, & (y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t}, \\ \mathcal{U}_r^{[k]}(y_i, t_n) = g_b(y_i, t_n), & (y_i, t_n) \in \gamma_{b, r}^{N_y, m_\tau}, \\ \mathcal{U}_r^{[k]}(1 - 2\rho, t_n) = \mathcal{I}_n \mathcal{U}^{[k-1]}(1 - 2\rho, t_n), & t_n \in \Omega^{M_t}, \\ \Gamma_r^{N_y, M_t} \mathcal{U}_r^{[k]}(1, t_n) \\ = g_r(t_n) + \frac{h_r \delta_2}{2\sqrt{\epsilon}} (-b(1, t_n) \mathcal{U}(1, t_n - m_\tau) + f(1, t_n)), & t_n \in \Omega^{M_t}, \end{cases} \tag{17}$$

$$\begin{cases} [L_m^{N_y, M_t} \mathcal{U}_m^{[k]}]_{i,n} + b_{i,n} \mathcal{U}_{m; i, n - m_\tau}^{[k]} = f_{i,n}, & (y_i, t_n) \in \overline{\mathcal{Q}}_m^{N_y, M_t}, \\ \mathcal{U}_m^{[k]}(y_i, t_n) = g_b(y_i, t_n), & (y_i, t_n) \in \gamma_{b, m}^{N_y, m_\tau}, \\ \mathcal{U}_m^{[k]}(\rho, t_n) = \mathcal{I}_n \mathcal{U}_\ell^{[k]}(\rho, t_n), & t_n \in \Omega^{M_t}, \\ \mathcal{U}_m^{[k]}(1 - \rho, t_n) = \mathcal{I}_n \mathcal{U}_r^{[k]}(1 - \rho, t_n), & t_n \in \Omega^{M_t}. \end{cases} \tag{18}$$

The aforementioned process is repeated until the condition $\|\mathcal{U}^{[k+1]} - \mathcal{U}^{[k]}\|_{\overline{\mathcal{Q}}^{N_y, M_t}} \leq \delta$ is not met, where δ represents a tolerance specified by the user and the approximate solution $\mathcal{U}^{[k]}$ of (3)-(4) is defined by

$$\mathcal{U}^{[k]}(y_i, t_n) = \begin{cases} \mathcal{U}_\ell^{[k]}(y_i, t_n), & (y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m, \\ \mathcal{U}_m^{[k]}(y_i, t_n), & (y_i, t_n) \in \overline{\mathcal{Q}}_m^{N_y, M_t}, \\ \mathcal{U}_r^{[k]}(y_i, t_n), & (y_i, t_n) \in \overline{\mathcal{Q}}_r^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m. \end{cases} \tag{19}$$

4 Convergence analysis

To prove the convergence of the proposed method we first introduce the maximum principle for the discrete operators $[L_p^{N_y, M_t} \mathcal{U}_p]_{i,n}$ and $[\mathcal{L}_p^{N_y, M_t} \mathcal{U}_p]_{i,n} = [L_p^{N_y, M_t} \mathcal{U}_p]_{i,n} +$

$b_{i,n} \mathcal{U}_{p;i,n-m\tau}$ for $(y_i, t_n) \in \mathcal{Q}_p^{N_y, M_t}$. Suppose the operators $G_p^{N_y, M_t}$, $p = \ell, m, r$, are defined as $G_\ell^{N_y, M_t} Z(0, t_n) = \Gamma_\ell^{N_y, M_t} Z(0, t_n)$, $G_r^{N_y, M_t} Z(1, t_n) = \Gamma_r^{N_y, M_t} Z(1, t_n)$ and $G_p^{N_y, M_t} Z(a, t_n) = Z(a, t_n)$ at the points $a = \rho, 1 - \rho, 2\rho, 1 - 2\rho$ for $t_n \in \Omega^{M_t}$.

Lemma 6 *Let Z_p , $p = \ell, m, r$, be a mesh function satisfying $G_p^{N_y, M_t} Z_p(y_0, t_n) \geq 0$, $G_p^{N_y, M_t} Z_p(y_{N_y}, t_n) \geq 0$, $t_n \in \Omega^{M_t}$ and $Z_p(y_i, t_n) \geq 0$, $(y_i, t_n) \in \gamma_{b,p}^{N_y, m\tau}$. Then $L_p^{N_y, M_t} Z_p(y_i, t_n) \geq 0$, $(y_i, t_n) \in \mathcal{Q}_p^{N_y, M_t}$ implies that $Z_p(y_i, t_n) \geq 0$ for $(y_i, t_n) \in \overline{\mathcal{Q}}_p^{N_y, M_t}$.*

Proof We prove the result for $p = \ell$. A similar argument can be used for $p = m, r$. Suppose the mesh function Z_ℓ attains its minimum at (\bar{y}_i, \bar{t}_n) , that is,

$$Z_\ell(\bar{y}_i, \bar{t}_n) = \min_{(y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t}} Z_\ell(y_i, t_n) < 0.$$

Clearly, (\bar{y}_i, \bar{t}_n) is not a member of $\gamma_{b,\ell}^{N_y, M_t}$. If $(\bar{y}_i, \bar{t}_n) \in \mathcal{Q}_\ell^{N_y, M_t}$, then

$$L_\ell^{N_y, M_t} Z_\ell(\bar{y}_i, \bar{t}_n) = (F_t^- Z_\ell - \epsilon D_y Z_\ell + a Z_\ell)(\bar{y}_i, \bar{t}_n) < 0,$$

since $F_t^- Z_\ell(\bar{y}_i, \bar{t}_n) \leq 0$ and $D_y Z_\ell(\bar{y}_i, \bar{t}_n) \geq 0$, which contradicts the assumption.

Now, suppose $(\bar{y}_i, \bar{t}_n) \in \gamma_\ell^{N_y, M_t}$, then

$$G_\ell^{N_y, M_t} Z_\ell(\bar{y}_i, \bar{t}_n) = \left(\alpha_1 Z_\ell - \delta_1 \sqrt{\epsilon} F_y^+ Z_\ell + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (a Z_\ell + F_t^- Z_\ell) \right) (\bar{y}_i, \bar{t}_n) < 0,$$

since $F_t^- Z_\ell(\bar{y}_i, \bar{t}_n) \leq 0$ and $F_y^+ Z_\ell(\bar{y}_i, \bar{t}_n) \geq 0$, which contradicts the assumption.

Hence, the proof is complete. \square

Lemma 7 *Let Y_p , $p = \ell, m, r$, be a mesh function satisfying $G_p^{N_y, M_t} Y_p(y_0, t_n) \geq 0$, $G_p^{N_y, M_t} Y_p(y_{N_y}, t_n) \geq 0$, $t_n \in \Omega^{M_t}$ and $Y_p(y_i, t_n) \geq 0$, $(y_i, t_n) \in \gamma_{b,p}^{N_y, m\tau}$. Then $\mathcal{L}_p^{N_y, M_t} Y_p(y_i, t_n) \geq 0$, $(y_i, t_n) \in \mathcal{Q}_p^{N_y, M_t}$ implies that $Y_p(y_i, t_n) \geq 0$ for $(y_i, t_n) \in \overline{\mathcal{Q}}_p^{N_y, M_t}$.*

Proof Define $\overline{\mathcal{Q}}_{p,q_1}^{N_y, m\tau} = \overline{\mathcal{D}}_p^{N_y} \times \overline{\Omega}_{q_1}^{m\tau}$, $p = \ell, m, r$, $q_1 = 0, 1, \dots, n$, where $\overline{\Omega}_{q_1}^{m\tau}$ is defined by splitting $[(q_1 - 1)\tau, q_1\tau]$ into $m\tau$ intervals of equal length. Further, we introduce $\overline{\mathcal{Q}}_{p,q_1,q_2}^{N_y, m\tau} = \overline{\mathcal{D}}_p^{N_y} \times \overline{\Omega}_{q_1,q_2}^{m\tau}$, where $\overline{\Omega}_{q_1,q_2}^{m\tau}$ is obtained by splitting the interval $[(q_1 - 1)\tau, q_2\tau]$ into $(q_2 - q_1 + 1)m\tau$ intervals of equal length.

Assume that $Z_p(y_i, t_n) = Y_p(y_i, t_n)$, for $(y_i, t_n) \in \overline{\mathcal{Q}}_{p;0,1}^{N_y, m\tau}$. Therefore, we have

$$Z_p(y_i, t_n) \geq 0, (y_i, t_n) \in \gamma_{b,p}^{N_y, m\tau} \text{ and } G_p^{N_y, M_t} Z_p(y_0, t_n) \geq 0, G_p^{N_y, M_t} Z_p(y_{N_y}, t_n) \geq 0, t_n \in \Omega_1^{m\tau},$$

and

$$\begin{aligned} [L_p^{N_y, M_t} Z_p]_{i,n} &\geq -b_{i,n} Z_p(y_i, t_{n-m_\tau}) \geq 0, \\ (y_i, t_n) &\in \mathcal{Q}_{p,1}^{N_y, m_\tau}. \end{aligned}$$

Hence, Lemma 6 implies that $Y_p(y_i, t_n) = Z_p(y_i, t_n) \geq 0$, for $(y_i, t_n) \in \mathcal{Q}_{p,1}^{N_y, m_\tau}$. Similarly, for $q_1 \geq 2$, we can establish $Y_p(y_i, t_n) \geq 0$, for $(y_i, t_n) \in \mathcal{Q}_{p,q_1}^{N_y, m_\tau}$ using the fact $Y_p(y_i, t_n) \geq 0$ for $(y_i, t_n) \in \mathcal{Q}_{p,q_1-1}^{N_y, m_\tau}$. \square

Next, we define some values that will be used in further analysis.

$$\begin{aligned} \chi_\rho &= \max \left\{ \max_{t_n \in \Omega^{M_t}} |(\overline{\mathcal{U}}_m - \overline{\mathcal{U}}_\ell)(\rho, t_n)|, \max_{t_n \in \Omega^{M_t}} |(\overline{\mathcal{U}}_m - \overline{\mathcal{U}}_r)(1 - \rho, t_n)| \right\}, \\ \chi_{2\rho} &= \max \left\{ \max_{t_n \in \Omega^{M_t}} |(\overline{\mathcal{U}}_\ell - \overline{\mathcal{U}}_m)(2\rho, t_n)|, \max_{t_n \in \Omega^{M_t}} |(\overline{\mathcal{U}}_r - \overline{\mathcal{U}}_m)(1 - 2\rho, t_n)| \right\}, \\ \chi^{[k]} &= \max \left\{ \max_{t_n \in \Omega^{M_t}} |(\overline{\mathcal{U}}_\ell - \mathcal{I}_n \mathcal{U}^{[k-1]})(2\rho, t_n)|, \max_{t_n \in \Omega^{M_t}} |(\overline{\mathcal{U}}_r - \mathcal{I}_n \mathcal{U}^{[k-1]})(1 - 2\rho, t_n)| \right\}, \\ \xi^{[k]} &= \max \left\{ \|\overline{\mathcal{U}}_\ell - \mathcal{U}^{[k]}\|_{\mathcal{Q}_\ell^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m}, \|\overline{\mathcal{U}}_m - \mathcal{U}^{[k]}\|_{\overline{\mathcal{Q}}_m^{N_y, M_t}}, \|\overline{\mathcal{U}}_r - \mathcal{U}^{[k]}\|_{\overline{\mathcal{Q}}_r^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m} \right\}. \end{aligned}$$

Further, we define the following auxiliary problems that will be used to prove the convergence of the method.

$$\begin{cases} [L_\ell^{N_y, M_t} \overline{\mathcal{U}}_\ell]_{i,n} + b_{i,n} \overline{\mathcal{U}}_{\ell; i, n - m_\tau} = f_{i,n}, & (y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}, \\ \overline{\mathcal{U}}_\ell(y_i, t_n) = u(y_i, t_n), & (y_i, t_n) \in \mathcal{Y}_{b,\ell}^{N_y, m_\tau}, \\ \Gamma_\ell^{N_y, M_t} \overline{\mathcal{U}}_\ell(0, t_n) = \Gamma_\ell^{N_y, M_t} \mathcal{U}^{[k]}(0, t_n), \quad \overline{\mathcal{U}}_\ell(2\rho, t_n) \\ \quad = u(2\rho, t_n), & t_n \in \Omega^{M_t}, \end{cases}$$

$$\begin{cases} [L_m^{N_y, M_t} \overline{\mathcal{U}}_m]_{i,n} + b_{i,n} \overline{\mathcal{U}}_{m; i, n - m_\tau} = f_{i,n}, & (y_i, t_n) \in \mathcal{Q}_m^{N_y, M_t}, \\ \overline{\mathcal{U}}_m(y_i, t_n) = u(y_i, t_n), & (y_i, t_n) \in \mathcal{Y}_{b,m}^{N_y, m_\tau}, \\ \overline{\mathcal{U}}_m(\rho, t_n) = u(\rho, t_n), \quad \overline{\mathcal{U}}_m(1 - \rho, t_n) = u(1 - \rho, t_n), & t_n \in \Omega^{M_t}, \end{cases}$$

$$\begin{cases} [L_r^{N_y, M_t} \overline{\mathcal{U}}_r]_{i,n} + b_{i,n} \overline{\mathcal{U}}_{r; i, n - m_\tau} = f_{i,n}, & (y_i, t_n) \in \mathcal{Q}_r^{N_y, M_t}, \\ \overline{\mathcal{U}}_r(y_i, t_n) = u(y_i, t_n), & (y_i, t_n) \in \mathcal{Y}_{b,r}^{N_y, m_\tau}, \\ \overline{\mathcal{U}}_r(1 - 2\rho, t_n) = u(1 - 2\rho, t_n), \quad \Gamma_r^{N_y, M_t} \overline{\mathcal{U}}_r(1, t_n) \\ \quad = \Gamma_r^{N_y, M_t} \mathcal{U}^{[k]}(1, t_n), & t_n \in \Omega^{M_t}. \end{cases}$$

Lemma 8 Let u be the exact solution of (3)-(4) and $\overline{\mathcal{U}}_p$ the approximate solution provided by the proposed algorithm. Then, it holds that

$$\|u - \overline{\mathcal{U}}_p\|_{\mathcal{Q}_p^{N_y, M_t}} \leq C(\Delta t + N_y^{-2} \ln^2 N_y). \quad (20)$$

Proof For $(y_i, t_n) \in \mathcal{Q}_p^{N_y, M_t}$, $p = \ell, m, r$,

$$\begin{aligned} \mathcal{L}_p^{N_y, M_t}(u - \bar{\mathcal{U}}_p)(y_i, t_n) &= \mathcal{L}_p^{N_y, M_t} u(y_i, t_n) - \mathcal{L}u(y_i, t_n) \\ &= (F_t^- u - u_t)(y_i, t_n) - \epsilon (D_y u - u_{yy})(y_i, t_n). \end{aligned} \tag{21}$$

Applying Taylor expansions and using the bounds in (6) with $h_\ell \leq C\sqrt{\epsilon}N_y^{-1} \ln N_y$ we obtain

$$\begin{aligned} \left| [\mathcal{L}_\ell^{N_y, M_t}(u - \bar{\mathcal{U}}_\ell)]_{i,n} \right| &\leq \frac{1}{2}(t_n - t_{n-1}) \|u_{tt}(y, \cdot)\|_{[t_{n-1}, t_n]} \\ &\quad + \frac{\epsilon}{12} h_\ell^2 \|u_{yyyy}(\cdot, t_n)\|_{[y_{i-1}, y_{i+1}]} \\ &\leq C(\Delta t + N_y^{-2} \ln^2 N_y). \end{aligned} \tag{22}$$

Next, for $t_n \in \Omega^{M_t}$ we have $(u - \bar{\mathcal{U}}_\ell)(\rho, t_n) = 0$ and equations (4) and (15) give

$$\begin{aligned} \Gamma_\ell^{N_y, M_t}(u - \bar{\mathcal{U}}_\ell)(0, t_n) &= (\alpha_1 u - \delta_1 \sqrt{\epsilon} F_y^+ u + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}}(au + F_t^- u) - (\alpha_1 u - \delta_1 \sqrt{\epsilon} u_y))(0, t_n) \\ &\quad - \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (-b(0, t_n)u(0, t_{n-m_\tau}) + f(0, t_n)) \\ &= \delta_1 \sqrt{\epsilon} (u_y - F_y^+ u)(0, t_n) \\ &\quad + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} ((au - f + u_t)(0, t_n) + b(0, t_n)u(0, t_{n-m_\tau})) \\ &\quad + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (F_t^- u - u_t)(0, t_n) \\ &= \delta_1 \sqrt{\epsilon} \left(u_y - F_y^+ u + \frac{h_\ell}{2} u_{yy} \right) (0, t_n) + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (F_t^- u - u_t)(0, t_n). \end{aligned}$$

Now, using Taylor expansions and the bounds in (6), we get

$$\begin{aligned} \left| \Gamma_\ell^{N_y, M_t}(u - \bar{\mathcal{U}}_\ell)(0, t_n) \right| &\leq \frac{\delta_1 \sqrt{\epsilon}}{6} h_\ell^2 \|u_{yyy}(\cdot, t_n)\|_{[y_{i-1}, y_{i+1}]} \\ &\quad + \frac{h_\ell \delta_1}{4\sqrt{\epsilon}} (t_n - t_{n-1}) \|u_{tt}(y_i, \cdot)\|_{[t_{n-1}, t_n]} \\ &\leq C(\Delta t + N_y^{-2} \ln^2 N_y). \end{aligned}$$

Further, applying Lemma 7 to the mesh function $\varphi^\pm(y_i, t_n) = C(\Delta t + N_y^{-2} \ln^2 N_y) \pm (u - \bar{\mathcal{U}}_\ell)(y_i, t_n)$, we get

$$\|u - \bar{\mathcal{U}}_\ell\|_{\mathcal{Q}_\ell^{N_y, M_t}} \leq C(\Delta t + N_y^{-2} \ln^2 N_y).$$

Similarly,

$$\|u - \bar{\mathcal{U}}_r\|_{\mathcal{Q}_r^{N_y, M_t}} \leq C(\Delta t + N_y^{-2} \ln^2 N_y).$$

Now, we consider two cases to find an estimate for $\|u - \overline{\mathcal{U}}_m\|_{\overline{\mathcal{Q}}_m^{N_y, M_t}}$.

Case 1: If $\rho = 1/4$, then $h_m = 1/(2N_y)$ and $\epsilon^{-1} \leq C \ln^2 N_y$. Now, using (6) and equation (21), from Taylor expansions we get

$$\left| [\mathcal{L}_m^{N_y, M_t}(u - \overline{\mathcal{U}}_m)]_{i,n} \right| \leq C(\Delta t + N_y^{-2} \ln^2 N_y), \quad (y_i, t_n) \in \overline{\mathcal{Q}}_m^{N_y, M_t}.$$

Case 2: If $\rho = 2\sqrt{\frac{\epsilon}{\alpha}} \ln N_y$, then $h_m \leq CN_y^{-1}$. So by using Taylor expansion and (6), we get

$$\left| (F_t^- u - u_t)(y_i, t_n) \right| \leq C\Delta t \text{ for } (y_i, t_n) \in \overline{\mathcal{Q}}_m^{N_y, M_t}.$$

To get a bound of the term $\epsilon |(D_y u - u_{yy})(y_i, t_n)|$ we use the solution decomposition $u = v + w$ such that

$$\epsilon |(D_y u - u_{yy})(y_i, t_n)| \leq \epsilon |(D_y v - v_{yy})(y_i, t_n)| + \epsilon |(D_y w - w_{yy})(y_i, t_n)|.$$

Now, by using Taylor expansions and the bounds in (11), (12), we get

$$\begin{aligned} \epsilon |(D_y u - u_{yy})u(y_i, t_n)| &\leq C\epsilon h_m^2 \|v_{yyyy}(\cdot, t_n)\|_{[y_{i-1}, y_{i+1}]} + C\epsilon \|w_{yy}(\cdot, t_n)\|_{[y_{i-1}, y_{i+1}]} \\ &\leq CN_y^{-2}. \end{aligned}$$

Thus, for $(y_i, t_n) \in \overline{\mathcal{Q}}_m^{N_y, M_t}$, we have

$$\left| [\mathcal{L}_m^{N_y, M_t}(u - \overline{\mathcal{U}}_m)]_{i,n} \right| \leq C(\Delta t + N_y^{-2} \ln^2 N_y).$$

Now, applying Lemma 7 to the mesh function $C(\Delta t + N_y^{-2} \ln^2 N_y) \pm (u - \overline{\mathcal{U}}_m)(y_i, t_n)$, we get

$$\|u - \overline{\mathcal{U}}_m\|_{\overline{\mathcal{Q}}_m^{N_y, M_t}} \leq C(\Delta t + N_y^{-2} \ln^2 N_y).$$

Therefore, by combining the aforementioned bounds, the desired result can be obtained. □

In the underlying theorem we have shown that the method converges uniformly for $\rho = 2\sqrt{\frac{\epsilon}{\beta}} \ln N_y$.

Theorem 1 *The solution of (3)-(4) and the first iterate of the proposed scheme satisfy*

$$\|u - \mathcal{U}^{[1]}\|_{\overline{\mathcal{Q}}_m^{N_y, M_t}} \leq C(\Delta t + N_y^{-2} \ln^2 N_y).$$

Proof Suppose

$$\Psi^\pm(y_i, t_n) = \Psi_\ell(y_i, t_n) \pm (\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n),$$

is the mesh function, where Ψ_ℓ is the solution of following problem

$$\begin{cases} F_t^- \Psi_\ell(x_i, t_n) - \epsilon D_y \Psi_\ell(x_i, t_n) + \frac{\beta}{2} \Psi_\ell(x_i, t_n) \\ \quad + \frac{\beta}{2} \Psi_\ell(x_i, t_n - m_\tau) = 0, & (y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}, \\ \Psi_\ell(y_i, t_n) = \chi^{[1]} \frac{\varphi_2 \zeta_1^i - \varphi_1 \zeta_2^i}{\varphi_2 \zeta_1^{N_y} - \varphi_1 \zeta_2^{N_y}}, & (y_i, t_n) \in \gamma_{b,\ell}^{N_y, m_\tau}, \\ \tilde{\Gamma}_\ell^{N_y, M_t} \Psi_\ell(0, t_n) = 0, & t_n \in \Omega^{M_t}, \\ \Psi_\ell(2\rho, t_n) = \chi^{[1]}, & t_n \in \Omega^{M_t}, \end{cases} \quad (23)$$

where the operator $\tilde{\Gamma}_\ell^{N_y, M_t}$ is defined as

$$\tilde{\Gamma}_\ell^{N_y, M_t} \Psi_\ell(0, t_n) = \alpha_1 \Psi_\ell(0, t_n) - \delta_1 \sqrt{\epsilon} F_y^+ \Psi_\ell(0, t_n) + \frac{h_\ell \delta_1}{2\sqrt{\epsilon}} (\beta \Psi_\ell + F_t^- \Psi_\ell)(0, t_n),$$

$\varphi_1 = 2\alpha_1 \sqrt{\epsilon} h_\ell - 2\delta_1 \epsilon (\zeta_1 - 1) + \delta_1 h_\ell^2 a$, $\varphi_2 = 2\alpha_1 \sqrt{\epsilon} h_\ell - 2\delta_1 \epsilon (\zeta_2 - 1) + \delta_1 h_\ell^2 a$ and $\zeta_1 = \beta_1 + \beta_2$, $\zeta_2 = \beta_1 - \beta_2$ with

$$\beta_1 = 1 + \left(\frac{\rho}{N_y} \sqrt{\frac{\beta}{\epsilon}} \right)^2, \quad \beta_2 = 2 \left(\frac{\rho}{N_y} \sqrt{\frac{\beta}{\epsilon}} \right) \sqrt{1 + \left(\frac{\rho}{N_y} \sqrt{\frac{\beta}{\epsilon}} \right)^2},$$

and $\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]}$ satisfies

$$\begin{aligned} \mathcal{L}_\ell^{N_y, M_t} (\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n) &= 0, \quad (y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}, \\ (\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n) &= 0, \quad (y_i, t_n) \in \gamma_{b,\ell}^{N_y, m_\tau} \\ \text{and } \Gamma_\ell^{N_y, M_t} (\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(0, t_n) &= 0, \quad |(\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(2\rho, t_n)| \leq \chi^{[1]}, \quad t_n \in \Omega^{M_t}. \end{aligned}$$

The solution of problem (23) is

$$\Psi_\ell(y_i, t_n) = \chi^{[1]} \frac{\varphi_2 \zeta_1^i - \varphi_1 \zeta_2^i}{\varphi_2 \zeta_1^{N_y} - \varphi_1 \zeta_2^{N_y}}, \quad (24)$$

which satisfies $\Psi_\ell(y_i, t_n) \geq 0$, $(y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}$ and it is monotonically increasing. Also, we have $\Psi^\pm(y_i, t_n) \geq 0$ for $(y_i, t_n) \in \gamma_{b,\ell}^{N_y, m_\tau}$, $\Gamma_\ell^{N_y, M_t} \Psi^\pm(0, t_n) \geq 0$, $\Psi^\pm(2\rho, t_n) \geq 0$ for $t_n \in \Omega^{M_t}$, and $\mathcal{L}_\ell^{N_y, M_t} \Psi^\pm(y_i, t_n) \geq 0$, for $(y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}$.

Thus, by applying Lemma 7 to the mesh function $\Psi^\pm(y_i, t_n)$, we get

$$|(\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n)| \leq \Psi_\ell(y_i, t_n) \quad \text{for } (y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t}.$$

Since $y_i \leq \rho$ for $(y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m$, equation (24) implies that

$$\begin{aligned} \Psi_\ell(y_i, t_n) &\leq \chi^{[1]} \frac{\varphi_2 \zeta_1^{N_y/2} - \varphi_1 \zeta_2^{N_y/2}}{\varphi_2 \zeta_1^{N_y} - \varphi_1 \zeta_2^{N_y}} = \chi^{[1]} \frac{\lambda \zeta_1^{N_y/2} - \zeta_2^{N_y/2}}{\lambda \zeta_1^{N_y} - \zeta_2^{N_y}}, \text{ where } \lambda = \varphi_2/\varphi_1, \\ &= \frac{\chi^{[1]}}{\lambda \zeta_1^{N_y/2} + \zeta_2^{N_y/2}} \left[\frac{\lambda^2 \zeta_1^{N_y} - \zeta_2^{N_y}}{\lambda \zeta_1^{N_y} - \zeta_2^{N_y}} \right]. \end{aligned}$$

Therefore, we have

$$\Psi_\ell(y_i, t_n) \leq \frac{\chi^{[1]}}{\lambda \zeta_1^{N_y/2} + \zeta_2^{N_y/2}} \left[\frac{\lambda^2 - \zeta_2^{N_y}/\zeta_1^{N_y}}{\lambda - \zeta_2^{N_y}/\zeta_1^{N_y}} \right] \leq \chi^{[1]} \frac{C\lambda}{\lambda \zeta_1^{N_y/2} + \zeta_2^{N_y/2}},$$

since $\lambda > 1$ and $\zeta_2/\zeta_1 < 1$, and hence we get $\left[\frac{\lambda^2 - \zeta_2^{N_y}/\zeta_1^{N_y}}{\lambda - \zeta_2^{N_y}/\zeta_1^{N_y}} \right] \leq C\lambda$. Thus, we have

$$\Psi_\ell(y_i, t_n) \leq \frac{\chi^{[1]}C}{\zeta_1^{N_y/2}}.$$

Further, employing the arguments presented in [18, Lemma 5.1] and utilizing the inequality $\chi^{[1]} \leq C$, we can establish that $\Psi_\ell(y_i, t_n) \leq CN_y^{-2}$ for $(y_i, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m$. Thus, it is

$$\|\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]}\|_{\overline{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m} \leq CN_y^{-2}. \tag{25}$$

Proceeding in a similar way we can arrive at

$$\|\overline{\mathcal{U}}_r - \mathcal{U}_r^{[1]}\|_{\overline{\mathcal{Q}}_r^{N_y, M_t} \setminus \overline{\mathcal{Q}}_m} \leq CN_y^{-2}. \tag{26}$$

Now, take $\overline{\mathcal{U}}_m - \mathcal{U}_m^{[1]}$, which satisfies the following

$$\begin{aligned} [\mathcal{L}_m^{N_y, M_t}(\overline{\mathcal{U}}_m - \mathcal{U}_m^{[1]})]_{i,n} &= 0, \quad (y_i, t_n) \in \mathcal{Q}_m^{N_y, M_t}, \\ (\overline{\mathcal{U}}_m - \mathcal{U}_m^{[1]})(y_i, t_n) &= 0, \quad (y_i, t_n) \in \gamma_{b,m}^{N_y, m\tau}, \\ |(\overline{\mathcal{U}}_m - \mathcal{U}_m^{[1]})(\eta, t_n)| &\leq \chi\rho + CN_y^{-2}, \quad (\eta, t_n) \in \{\rho, 1 - \rho\} \times \Omega^{M_t}. \end{aligned}$$

Thus, Lemma 7 gives

$$\|\overline{\mathcal{U}}_m - \mathcal{U}_m^{[1]}\|_{\overline{\mathcal{Q}}_m^{N_y, M_t}} \leq \chi\rho + CN_y^{-2}, \tag{27}$$

and hence

$$\xi^{[1]} \leq \chi_\rho + CN_y^{-2}. \tag{28}$$

Also, we note that $\chi_\rho \leq C(\Delta t + N_y^{-2} \ln^2 N_y)$ using Lemma 8 and $(\rho, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t}$, $(1 - \rho, t_n) \in \overline{\mathcal{Q}}_r^{N_y, M_t}$. Thus, we get the result by combining (28) and Lemma 8. \square

In the following theorem, for $\rho = 1/4$, we find a bound on $\|\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[k]}\|_{\overline{\mathcal{Q}}^{N_y, M_t}}$ and then combine it with Lemma 8 to establish a bound on $\|u - \mathcal{U}^{[k]}\|_{\overline{\mathcal{Q}}^{N_y, M_t}}$.

Theorem 2 *The exact solution of problem (3)-(4) and the approximate solution $\mathcal{U}^{[k]}$ defined by (19) satisfy*

$$\|u - \mathcal{U}^{[k]}\|_{\overline{\mathcal{Q}}^{N_y, M_t}} \leq C(\Delta t + N_y^{-2} \ln^2 N_y) + C\left(\frac{5}{6}\right)^k. \tag{29}$$

Proof Let consider the mesh function

$$\Phi^\pm(y_i, t_n) = \frac{y_i + \sqrt{\epsilon}}{2\rho + \sqrt{\epsilon}} \chi^{[1]} \pm (\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n),$$

where $\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]}$ satisfies

$$\begin{aligned} \mathcal{L}_\ell^{N_y, M_t}(\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n) &= 0, \quad (y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}, \\ (\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n) &= 0, \quad (y_i, t_n) \in \gamma_{b, \ell}^{N_y, m_\tau}, \\ \Gamma_\ell^{N_y, M_t}(\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(0, t_n) &= 0, \quad |(\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(2\rho, t_n)| \leq \chi^{[1]}, \quad t_n \in \Omega^{M_t}. \end{aligned}$$

For $t_n \in \Omega^{M_t}$, we have

$$\begin{aligned} \Gamma_\ell^{N_y, M_t} \Phi^\pm(0, t_n) &= \Gamma_\ell^{N_y, M_t} \left[\frac{y_i + \sqrt{\epsilon}}{2\rho + \sqrt{\epsilon}} \chi^{[1]} \right] \pm \Gamma_\ell^{N_y, M_t} (\overline{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(0, t_n) \\ &= \frac{\chi^{[1]}}{2\rho + \sqrt{\epsilon}} \left[\alpha_1(y_0 + \sqrt{\epsilon}) - \delta_1 \sqrt{\epsilon} \left(\frac{y_1 - y_0}{h_\ell} \right) \right. \\ &\quad \left. + \frac{h_\ell(a_{i,n} + b_{i,n})\delta_1}{2\sqrt{\epsilon}} (y_0 + \sqrt{\epsilon}) \right] \\ &= \frac{\chi^{[1]}}{2\rho + \sqrt{\epsilon}} \left(\alpha_1 y_0 + (\alpha_1 - \delta_1)\sqrt{\epsilon} + \frac{h_\ell(a_{i,n} + b_{i,n})\delta_1}{2\sqrt{\epsilon}} (y_0 + \sqrt{\epsilon}) \right) \\ &\geq 0, \end{aligned}$$

i.e. $\Phi^\pm(y_i, t_n) \geq 0$, $(y_i, t_n) \in \gamma_{b, \ell}^{N_y, m_\tau}$ and $\Gamma_\ell^{N_y, M_t} \Phi^\pm(0, t_n) \geq 0$, $\Phi^\pm(2\rho, t_n) \geq 0$, $t_n \in \Omega^{M_t}$.

For $(y_i, t_n) \in \mathcal{Q}_\ell^{N_y, M_t}$, it is

$$\mathcal{L}_\ell^{N_y, M_t} \Phi^\pm(y_i, t_n) = (a_{i,n} + b_{i,n}) \left(\frac{y_i + \sqrt{\epsilon}}{2\rho + \sqrt{\epsilon}} \right) \chi^{[1]} \geq 0.$$

Then, Lemma 7 leads to

$$|(\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]})(y_i, t_n)| \leq \left(\frac{y_i + \sqrt{\epsilon}}{2\rho + \sqrt{\epsilon}} \right) \chi^{[1]} \quad \text{for } (y_i, t_n) \in \bar{\mathcal{Q}}_\ell^{N_y, M_t}.$$

Hence, for $(y_i, t_n) \in \bar{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \bar{\mathcal{Q}}_m$, we get

$$|\bar{\mathcal{U}}_\ell - \mathcal{U}_\ell^{[1]}|_{\bar{\mathcal{Q}}_\ell^{N_y, M_t} \setminus \bar{\mathcal{Q}}_m} \leq 5\chi^{[1]}/6, \quad \text{since } y_i \leq \rho. \tag{30}$$

Similarly, we can show that

$$|\bar{\mathcal{U}}_r - \mathcal{U}_r^{[1]}|_{\bar{\mathcal{Q}}_r^{N_y, M_t} \setminus \bar{\mathcal{Q}}_m} \leq 5\chi^{[1]}/6. \tag{31}$$

Since $(\rho, t_n) \in \bar{\mathcal{Q}}_\ell^{N_y, M_t}$ and $(1 - \rho, t_n) \in \bar{\mathcal{Q}}_r^{N_y, M_t}$, we have the following

$$\begin{aligned} \mathcal{L}_m^{N_y, M_t} (\bar{\mathcal{U}}_m - \mathcal{U}_m^{[1]})(y_i, t_n) &= 0, \quad (y, t_n) \in \bar{\mathcal{Q}}_m^{N_y, M_t}, \\ (\bar{\mathcal{U}}_m - \mathcal{U}_m^{[1]})(y_i, t_n) &= 0, \quad (y_i, t_n) \in \gamma_{b,m}^{N_y, M_t}, \\ |(\bar{\mathcal{U}}_m - \mathcal{U}_m^{[1]})(\eta, t_n)| &\leq \chi_\rho + 5\chi^{[1]}/6, \quad (\eta, t_n) \in \{\rho, 1 - \rho\} \times \Omega^{M_t}. \end{aligned}$$

So, applying Lemma 7, we get

$$|\bar{\mathcal{U}}_m - \mathcal{U}_m^{[1]}|_{\bar{\mathcal{Q}}_m^{N_y, M_t}} \leq \chi_\rho + 5\chi^{[1]}/6. \tag{32}$$

Next, we calculate a bound of $\xi^{[2]}$ for which we need to estimate $\chi^{[2]}$ first. We note that $(2\rho, t_n), (1 - 2\rho, t_n) \in \bar{\mathcal{Q}}_m^{N_y, M_t}$. Consequently

$$\begin{aligned} |(\bar{\mathcal{U}}_\ell - \mathcal{J}_n \mathcal{U}^{[1]})(2\rho, t_n)| &\leq \chi_{2\rho} + \chi_\rho + 5\chi^{[1]}/6, \\ |(\bar{\mathcal{U}}_r - \mathcal{J}_n \mathcal{U}^{[1]})(1 - 2\rho, t_n)| &\leq \chi_{2\rho} + \chi_\rho + 5\chi^{[1]}/6. \end{aligned}$$

Therefore, we have $\chi^{[2]} \leq \chi_{2\rho} + \chi_\rho + 5\chi^{[1]}/6$. Hence, it is

$$\max\{\xi^{[1]}, \chi^{[2]}\} \leq \Upsilon + 5\chi^{[1]}/6, \quad \Upsilon = \chi_{2\rho} + \chi_\rho.$$

By performing the aforementioned procedure, we have

$$\max\{\xi^{[k]}, \chi^{[k+1]}\} \leq \Upsilon + 5\chi^{[k]}/6.$$

Simplifying the above expression we obtain

$$\chi^{[k]} \leq 6\Upsilon + 5^{k-1}\chi^{[1]}/6^{k-1},$$

and therefore

$$\xi^{[k]} \leq 6\Upsilon + 5^k\chi^{[1]}/6^k. \tag{33}$$

Further, Lemma 8 gives $\Upsilon \leq C(\Delta t + N_y^{-2} \ln^2 N_y)$, since $(2\rho, t_n), (1 - 2\rho, t_n) \in \overline{\mathcal{Q}}_m^{N_y, M_t}$, and $(\rho, t_n) \in \overline{\mathcal{Q}}_\ell^{N_y, M_t}$ and $(1 - \rho, t_n) \in \overline{\mathcal{Q}}_r^{N_y, M_t}$. Also, we have $\chi^{[1]} \leq C$. Hence, we get the proof by combining (33) and Lemma 8. \square

5 Numerical results

Here, two test problems are considered to illustrate the numerical scheme discussed earlier. We compute the uniform error and convergence rate as $\mathbb{E}^{N_y, M_t} = \max_{\epsilon} \mathbb{E}_{\epsilon}^{N_y, M_t}$ and $\mathbb{R}^{N_y, M_t} = \log_2 \left(\frac{\mathbb{E}^{N_y, M_t}}{\mathbb{E}^{2N_y, 4M_t}} \right)$, taking $\epsilon \in \{10^{-1}, 10^{-2}, \dots, 10^{-8}\}$ and different values of the discretization parameters N_y, M_t . The formula for computation of errors $\mathbb{E}_{\epsilon}^{N_y, M_t}$ is given below. The approximate solutions are computed using Algorithm 1. The stopping criterion of the algorithm for the considered problems is $\|\mathcal{U}^{[k+1]} - \mathcal{U}^{[k]}\|_{\overline{\mathcal{Q}}^{N_y, M_t}} \leq N_y^{-2}$.

Algorithm 1: Algorithm

1. Initialize $\mathcal{U}^{[0]}$ such that $\mathcal{U}^{[0]}(y_i, t_n) = 0, (y_i, t_n) \in (0, 1) \times (0, T]$ and $\mathcal{U}^{[0]}(y_i, t_n) = 0, (y_i, t_n) \in [0, 1] \times [-\tau, 0]$. Set $k = 1$.
 2. Compute $\mathcal{U}_\ell^{[k]}$ and $\mathcal{U}_r^{[k]}$ by solving (16) and (17), respectively. Then compute $\mathcal{U}_m^{[k]}$ by solving (18).
 3. Update $\mathcal{U}^{[k]}$ by (19).
 4. The final solution $\mathcal{U}^{[k]}$ is obtained if the stopping condition is fulfilled; if not, set $k = k + 1$ and go to Step 2.
-

Example 1 Consider

$$\begin{cases} \frac{\partial u(y,t)}{\partial t} - \epsilon \frac{\partial^2 u(y,t)}{\partial y^2} + (1 + ye^{-t})u(y,t) - 0.9u(y,t - 1) \\ \quad = f(y,t) & (y,t) \in \mathcal{Q} := \mathcal{D} \times (0, 2], \\ u(y,t) - \sqrt{\epsilon} \frac{\partial u}{\partial y}(y,t) = g_\ell(t) & (y,t) \in \{0\} \times (0, 2], \\ u(y,t) + \sqrt{\epsilon} \frac{\partial u}{\partial y}(y,t) = g_r(t) & (y,t) \in \{1\} \times (0, 2], \\ u(y,t) = g_b(y,t) & (y,t) \in [0, 1] \times [-1, 0]. \end{cases}$$

where g_b, f, g_ℓ and g_r are determined with the help of the exact solution

$$u(y,t) = t \left[\frac{e^{-y/\sqrt{\epsilon}} + e^{(y-1)/\sqrt{\epsilon}}}{1 + e^{-1/\sqrt{\epsilon}}} - \cos^2 \pi y \right].$$

Table 1 Maximum pointwise errors $\mathbb{E}_\epsilon^{N_y, M_t}$, uniform errors \mathbb{E}^{N_y, M_t} and uniform convergence rate \mathbb{R}^{N_y, M_t} for Example 1

| $\epsilon = 10^{-p}$ | $N_y = 2^6$ $M_t = 4^2$ | $N_y = 2^7$ $M_t = 4^3$ | $N_y = 2^8$ $M_t = 4^4$ | $N_y = 2^9$ $M_t = 4^5$ | $N_y = 2^{10}$ $M_t = 4^6$ |
|-------------------------|----------------------------|----------------------------|----------------------------|----------------------------|-------------------------------|
| $p = 1$ | 1.683E-04 | 4.279E-05 | 1.067E-05 | 2.675E-06 | 6.686E-07 |
| 2 | 7.490E-04 | 1.872E-04 | 4.679E-05 | 1.169E-05 | 2.924E-06 |
| 3 | 7.205E-03 | 1.811E-03 | 4.535E-04 | 1.134E-04 | 2.835E-05 |
| 4 | 6.669E-02 | 1.776E-02 | 4.511E-03 | 1.132E-03 | 2.834E-04 |
| 5 | 7.316E-02 | 2.647E-02 | 8.832E-03 | 2.815E-03 | 8.708E-04 |
| 6 | 7.316E-02 | 2.647E-02 | 8.832E-03 | 2.815E-03 | 8.708E-04 |
| 7 | 7.316E-02 | 2.647E-02 | 8.833E-03 | 2.815E-03 | 8.708E-04 |
| 8 | 7.316E-02 | 2.647E-02 | 8.833E-03 | 2.815E-03 | 8.708E-04 |
| \mathbb{E}^{N_y, M_t} | 7.316E-02 | 2.647E-02 | 8.833E-03 | 2.815E-03 | 8.708E-04 |
| \mathbb{R}^{N_y, M_t} | 1.466 | 1.583 | 1.649 | 1.692 | |

Table 2 Required iteration counts to attain the convergence for Example 1

| $\epsilon = 10^{-p}$ | $N_y = 2^6$ $M_t = 4^2$ | $N_y = 2^7$ $M_t = 4^3$ | $N_y = 2^8$ $M_t = 4^4$ | $N_y = 2^9$ $M_t = 4^5$ | $N_y = 2^{10}$ $M_t = 4^6$ |
|----------------------|----------------------------|----------------------------|----------------------------|----------------------------|-------------------------------|
| $p = 1$ | 5 | 6 | 6 | 7 | 7 |
| 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 1 | 1 | 1 | 1 | 1 |
| 4 | 1 | 1 | 1 | 1 | 1 |
| 5 | 1 | 1 | 1 | 1 | 1 |
| 6 | 1 | 1 | 1 | 1 | 1 |
| 7 | 1 | 1 | 1 | 1 | 1 |
| 8 | 1 | 1 | 1 | 1 | 1 |

Table 3 Uniform error \mathbb{E}^{N_y, M_t} and uniform convergence rate \mathbb{R}^{N_y, M_t} for Example 1

| Methods | | $N_y = 2^6$ $M_t = 4^2$ | $N_y = 2^7$ $M_t = 4^3$ | $N_y = 2^8$ $M_t = 4^4$ | $N_y = 2^9$ $M_t = 4^5$ | $N_y = 2^{10}$ $M_t = 4^6$ |
|-----------------|-------------------------|----------------------------|----------------------------|----------------------------|----------------------------|-------------------------------|
| Proposed method | \mathbb{E}^{N_y, M_t} | 7.316E-02 | 2.647E-02 | 8.833E-03 | 2.815E-03 | 8.708E-04 |
| | \mathbb{R}^{N_y, M_t} | 1.466 | 1.583 | 1.649 | 1.692 | |
| Method [31] | \mathbb{E}^{N_y, M_t} | 7.082E-02 | 2.803E-02 | 9.918E-03 | 3.284E-03 | 1.440E-03 |
| | \mathbb{R}^{N_y, M_t} | 1.337 | 1.4999 | 1.594 | 1.653 | |

The maximum pointwise errors for this example are evaluated as $\mathbb{E}_\epsilon^{N_y, M_t} = \|u - \mathcal{U}^{N_y, M_t}\|_{\mathcal{Q}^{N_y, M_t}}$, where u is the exact solution and \mathcal{U}^{N_y, M_t} is the approximate solution with the proposed scheme.

Table 4 Maximum pointwise errors $\mathbb{E}_\epsilon^{N_y, M_t}$, uniform errors \mathbb{E}^{N_y, M_t} and uniform convergence rate \mathbb{R}^{N_y, M_t} for Example 2

| $\epsilon = 10^{-p}$ | $N_y = 2^6$ $M_t = 4^2$ | $N_y = 2^7$ $M_t = 4^3$ | $N_y = 2^8$ $M_t = 4^4$ | $N_y = 2^9$ $M_t = 4^5$ | $N_y = 2^{10}$ $M_t = 4^6$ |
|-------------------------|----------------------------|----------------------------|----------------------------|----------------------------|-------------------------------|
| $p = 1$ | 1.7501E-03 | 4.4629E-04 | 1.1214E-04 | 2.8072E-05 | 7.0201E-06 |
| | 5.0496E-03 | 1.2733E-03 | 3.1903E-04 | 7.9800E-05 | 1.9953E-05 |
| | 1.0580E-02 | 2.6743E-03 | 6.7038E-04 | 1.6771E-04 | 4.1935E-05 |
| | 2.8278E-02 | 7.3827E-03 | 1.8647E-03 | 4.6736E-04 | 1.1691E-04 |
| | 3.1430E-02 | 1.1480E-02 | 3.8707E-03 | 1.2453E-03 | 3.8822E-04 |
| | 3.1426E-02 | 1.1479E-02 | 3.8705E-03 | 1.2453E-03 | 3.8822E-04 |
| | 3.1425E-02 | 1.1479E-02 | 3.8705E-03 | 1.2453E-03 | 3.8822E-04 |
| | 3.1425E-02 | 1.1479E-02 | 3.8705E-03 | 1.2453E-03 | 3.8822E-04 |
| \mathbb{E}^{N_y, M_t} | 3.1425E-02 | 1.1479E-02 | 3.8705E-03 | 1.2453E-03 | 3.8822E-04 |
| \mathbb{R}^{N_y, M_t} | 1.453 | 1.568 | 1.636 | 1.682 | |

Table 5 Required iteration counts to attain the convergence rate \mathbb{R}^{N_y, M_t} for Example 2

| $\epsilon = 10^{-p}$ | $N_y = 2^6$ $M_t = 4^2$ | $N_y = 2^7$ $M_t = 4^3$ | $N_y = 2^8$ $M_t = 4^4$ | $N_y = 2^9$ $M_t = 4^5$ | $N_y = 2^{10}$ $M_t = 4^6$ |
|----------------------|----------------------------|----------------------------|----------------------------|----------------------------|-------------------------------|
| $p = 1$ | 5 | 6 | 6 | 6 | 7 |
| 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 1 | 1 | 1 | 1 | 1 |
| 4 | 1 | 1 | 1 | 1 | 1 |
| 5 | 1 | 1 | 1 | 1 | 1 |
| 6 | 1 | 1 | 1 | 1 | 1 |
| 7 | 1 | 1 | 1 | 1 | 1 |
| 8 | 1 | 1 | 1 | 1 | 1 |

Table 6 Maximum pointwise errors $\mathbb{E}_\epsilon^{N_y, M_t}$, uniform errors \mathbb{E}^{N_y, M_t} and uniform convergence rate $\mathbb{R}_*^{N_y, M_t}$ for Example 2

| $\epsilon = 10^{-p}$ | $N_y = 2^5$ $M_t = 2^3$ | $N_y = 2^6$ $M_t = 2^4$ | $N_y = 2^7$ $M_t = 2^5$ | $N_y = 2^8$ $M_t = 2^6$ | $N_y = 2^9$ $M_t = 2^7$ |
|--------------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| $p = 1$ | 3.4408E-03 | 1.7690E-03 | 8.9756E-04 | 4.5203E-04 | 2.2684E-04 |
| | 9.9801E-03 | 5.0470E-03 | 2.5377E-03 | 1.2725E-03 | 6.3713E-04 |
| | 2.0719E-02 | 1.0478E-02 | 5.2684E-03 | 2.6416E-03 | 1.3226E-03 |
| | 2.5421E-02 | 1.2764E-02 | 6.4047E-03 | 3.2086E-03 | 1.6060E-03 |
| | 2.8453E-02 | 1.3283E-02 | 6.6825E-03 | 3.3515E-03 | 1.6783E-03 |
| | 2.8455E-02 | 1.3394E-02 | 6.7437E-03 | 3.3836E-03 | 1.6948E-03 |
| | 2.8456E-02 | 1.3423E-02 | 6.7598E-03 | 3.3920E-03 | 1.6990E-03 |
| | 2.8456E-02 | 1.3432E-02 | 6.7646E-03 | 3.3945E-03 | 1.7003E-03 |
| \mathbb{E}^{N_y, M_t} | 2.8456E-02 | 1.3432E-02 | 6.7646E-03 | 3.3945E-03 | 1.7003E-03 |
| $\mathbb{R}_*^{N_y, \Delta t}$ | 1.083 | 0.989 | 0.995 | 0.997 | |

Table 7 Required iteration counts to attain the convergence rate $\mathbb{R}_*^{N_y, \Delta t}$ for Example 2

| $\epsilon = 10^{-p}$ | $N_y = 2^5$ $M_t = 2^3$ | $N_y = 2^6$ $M_t = 2^4$ | $N_y = 2^7$ $M_t = 2^5$ | $N_y = 2^8$ $M_t = 2^6$ | $N_y = 2^9$ $M_t = 2^7$ |
|----------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| $p = 1$ | 5 | 6 | 6 | 7 | 7 |
| 2 | 2 | 2 | 2 | 2 | 2 |
| 3 | 1 | 1 | 1 | 1 | 1 |
| 4 | 1 | 1 | 1 | 1 | 1 |
| 5 | 1 | 1 | 1 | 1 | 1 |
| 6 | 1 | 1 | 1 | 1 | 1 |
| 7 | 1 | 1 | 1 | 1 | 1 |
| 8 | 1 | 1 | 1 | 1 | 1 |

In Table 1 we present the maximum pointwise errors $\mathbb{E}_\epsilon^{N_y, M_t}$, uniform errors \mathbb{E}^{N_y, M_t} and uniform convergence rate \mathbb{R}^{N_y, M_t} for Example 1, which verify the theoretical findings that we have proved earlier in Theorems 1 and 2. The iteration counts mentioned in Table 2 display the number of iteration needed to satisfy the stopping criterion for the algorithm. From Table 2 we can notice that only one iterate is required to stop the iterative procedure for small values of the perturbation parameters, i.e. we can get the desired approximate solution only after one iteration of the algorithm. Further, a comparison between our method and method [31] is given in Table 3. Notably, our method demonstrates lower errors than [31].

Example 2 Consider

$$\begin{cases} \frac{\partial u(y,t)}{\partial t} - \epsilon \frac{\partial^2 u(y,t)}{\partial y^2} + (1 + ye^{-t})u(y,t) - 0.9u(y,t-1) \\ \quad = f(y,t) & (y,t) \in \mathcal{Q} := \mathcal{D} \times (0,2], \\ u(y,t) - \sqrt{\epsilon} \frac{\partial u}{\partial y}(y,t) = g_\ell(t) & (y,t) \in \{0\} \times (0,2], \\ u(y,t) + \sqrt{\epsilon} \frac{\partial u}{\partial y}(y,t) = g_r(t) & (y,t) \in \{1\} \times (0,2], \\ u(y,t) = g_b(y,t) & (y,t) \in [0,1] \times [-1,0]. \end{cases}$$

where g_b, f, g_ℓ and g_r are calculated by the following exact solution

$$u(y,t) = (1 - e^{-t}) \left[\frac{e^{-y/\sqrt{\epsilon}} + e^{(y-1)/\sqrt{\epsilon}}}{1 + e^{-1/\sqrt{\epsilon}}} - \cos^2 \pi y \right].$$

The maximum pointwise errors, uniform errors and uniform rates of convergence are also computed for this example. Table 4 displays the numerical results for Example 2, which agree with the theoretical findings established in Theorems 1 and 2. From this, we can conclude that the rate of convergence of the proposed scheme is almost two. This is evident by the fact that in this case, the space discretization errors dominate the global errors. To indicate the contribution of time discretization errors to global errors the following formula is used

$$\mathbb{R}_*^{N_y, M_t} = \log_2 \left(\frac{\mathbb{E}^{N_y, M_t}}{\mathbb{E}^{2N_y, 2M_t}} \right).$$




Fig. 1 Solution plot for Example 1 with $\epsilon = 10^{-7}$ and $N_y = 64, M_t = 32$




Fig. 2 Solution plot for Example 2 with $\epsilon = 10^{-7}$ and $N_y = 64, M_t = 32$




Fig. 3 Error plot corresponding to Example 1




Fig. 4 Error plot corresponding to Example 2

The results are collected in Table 6. From this data, we can observe that the rate of convergence is one. Tables 5 and 7 show how many iterations the iterative procedure requires to reach the required accuracy.

The numerical solutions and corresponding loglog plots for Examples 1 and 2 are represented in Figs. 1, 2 and 3, 4 respectively. From Figs. 1 and 2 we can clearly observe the multiscale character of the solutions near the boundary points and that the slope obtained in the error plots validates the convergence order.

6 Conclusion

We developed a rapidly converging domain decomposition algorithm for solving singularly perturbed time delayed parabolic problems with mixed type boundary conditions. The numerical algorithm is proved to be robust convergent, with an accuracy of order one in time and almost two in space. Also, we observed that the numerical results obtained by the proposed numerical algorithm corresponding to the test problems validate the theoretical findings. The data displayed in Tables 2 and 5 illustrate that a single iteration is adequate to meet the specified threshold for low magnitudes of the perturbation parameter.

Author contributions All the authors have equivalently contributed to the work.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature. Not applicable.

Availability of data and materials Not applicable.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

Ethical approval Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Aakansha, Singh, J., Kumar, S.: Additive schemes based domain decomposition algorithm for solving singularly perturbed parabolic reaction-diffusion systems. *Comput. Appl. Math.* **40**, 1–15 (2021)
2. Adomian, G., Rach, R.: Nonlinear stochastic differential delay equations. *J. Math. Anal. Appl.* **91**(1), 94–101 (1983)
3. Ansari, A., Bakr, S., Shishkin, G.: A parameter-robust finite difference method for singularly perturbed delay parabolic partial differential equations. *J. Comput. Appl. Math.* **205**(1), 552–566 (2007)

4. Bashier, E.B.M., Patidar, K.C.: A novel fitted operator finite difference method for a singularly perturbed delay parabolic partial differential equation. *Appl. Math. Comput.* **217**(9), 4728–4739 (2011)
5. Burić, N., Todorović, D.: Dynamics of delay-differential equations modelling immunology of tumor growth. *Chaos Solitons Fractals* **13**(4), 645–655 (2002)
6. Chattopadhyay, J., Sarkar, R.R., El Abdllaoui, A.: A delay differential equation model on harmful algal blooms in the presence of toxic substances. *Math. Med. Biol. J. IMA* **19**(2), 137–161 (2002)
7. Derstine, M., Gibbs, H., Hopf, F., Kaplan, D.: Bifurcation gap in a hybrid optically bistable system. *Phys. Rev. A* **26**(6), 3720 (1982)
8. Glizer, V.: Asymptotic analysis and solution of a finite-horizon H_∞ control problem for singularly-perturbed linear systems with small state delay. *J. Optim. Theory Appl.* **117**(2), 295–325 (2003)
9. Govindarao, L., Mohapatra, J., Das, A.: A fourth-order numerical scheme for singularly perturbed delay parabolic problem arising in population dynamics. *J. Appl. Math. Comput.* **63**(1), 171–195 (2020)
10. Gracia, J., Lisbona, F.: A uniformly convergent scheme for a system of reaction-diffusion equations. *J. Comput. Appl. Math.* **206**(1), 1–16 (2007)
11. Han, Q.L.: Robust stability of uncertain delay-differential systems of neutral type. *Automatica* **38**(4), 719–723 (2002)
12. Kuang, Y.: *Delay Differential Equations: With Applications in Population Dynamics*. Academic Press (1993)
13. Kumar, S., Kumar, M.: High order parameter-uniform discretization for singularly perturbed parabolic partial differential equations with time delay. *Comput. Math. Appl.* **68**(10), 1355–1367 (2014)
14. Kumar, S., Rao, S.C.S.: A robust overlapping Schwarz domain decomposition algorithm for time-dependent singularly perturbed reaction-diffusion problems. *J. Comput. Appl. Math.* **261**, 127–138 (2014)
15. Kumar, S., Singh, J., Kumar, M.: A robust domain decomposition method for singularly perturbed parabolic reaction-diffusion systems. *J. Math. Chem.* **57**(5), 1557–1578 (2019)
16. Ladyzhenskaya, O., Solonnikov, V., Ural'tseva, N.: *Linear and quasilinear equations of parabolic type*, transl. *Math. Monographs*. Am. Math. Soc. **23** (1968)
17. Longtin, A., Milton, J.G.: Complex oscillations in the human pupil light reflex with “mixed” and delayed feedback. *Math. Biosci.* **90**(1–2), 183–199 (1988)
18. Miller, J.J.H., O’Riordan, E., Shishkin, G.I.: *Fitted Numerical Methods for Singular Perturbation Problems*. World Scientific, Singapore (1996)
19. Mohapatra, J., Priyadarshana, S., Raji Reddy, N.: Uniformly convergent computational method for singularly perturbed time delayed parabolic differential-difference equations. *Eng. Comput.* **40**(3), 694–717 (2023)
20. Priyadarshana, S.: Monotone hybrid numerical method for singularly perturbed time-lagged semilinear parabolic problems. *Natl. Acad. Sci. Lett.* **46**, 1–4 (2023)
21. Priyadarshana, S., Mohapatra, J.: An efficient numerical approximation for mixed singularly perturbed parabolic problems involving large time-lag. *Indian J. Pure Appl. Math.* 1–20 (2023) <https://doi.org/10.1007/s13226-023-00445-8>
22. Priyadarshana, S., Mohapatra, J., Govindrao, L.: An efficient uniformly convergent numerical scheme for singularly perturbed semilinear parabolic problems with large delay in time. *J. Appl. Math. Comput.* **68**, 2617–2639 (2022)
23. Priyadarshana, S., Mohapatra, J., Pattanaik, S.: Parameter uniform optimal order numerical approximations for time-delayed parabolic convection diffusion problems involving two small parameters. *Comput. Appl. Math.* **41**(6), 233 (2022)
24. Priyadarshana, S., Mohapatra, J., Pattanaik, S.: An improved time accurate numerical estimation for singularly perturbed semilinear parabolic differential equations with small space shifts and a large time lag. *Math. Comput. Simul.* **214**, 183–203 (2023)
25. Priyadarshana, S., Mohapatra, J., Pattanaik, S.: A second order fractional step hybrid numerical algorithm for time delayed singularly perturbed 2d convection-diffusion problems. *Appl. Numer. Math.* **189**, 107–129 (2023)
26. Quarteroni, A., Valli, A.: *Domain Decomposition Methods for Partial Differential Equations*. Oxford University Press (1999)
27. Rao, S.C.S., Kumar, S.: Robust high order convergence of an overlapping Schwarz method for singularly perturbed semilinear reaction-diffusion problems. *J. Comput. Math.* **31**, 509–521 (2013)

28. Rihan, F.A., Rahman, D.H.A., Lakshmanan, S., Alkhajeh, A.S.: A time delay model of tumour-immune system interactions: global dynamics, parameter estimation, sensitivity analysis. *Appl. Math. Comput.* **232**, 606–623 (2014)
29. Saini, S., Das, P., Kumar, S.: Parameter uniform higher order numerical treatment for singularly perturbed Robin type parabolic reaction diffusion multiple scale problems with large delay in time. *Appl. Numer. Math.* **196**, 1–21 (2024)
30. Schwarz, H.A.: Über einen Grenzübergang durch alternierendes Verfahren. *Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich* **15**, 272–286 (1870)
31. Selvi, P.A., Ramanujam, N.: A parameter uniform difference scheme for singularly perturbed parabolic delay differential equation with Robin type boundary condition. *Appl. Math. Comput.* **296**, 101–115 (2017)
32. Singh, J., Kumar, S., Kumar, M.: A domain decomposition method for solving singularly perturbed parabolic reaction-diffusion problems with time delay. *Numer. Methods Partial Differ. Equ.* **34**(5), 1849–1866 (2018)
33. Toselli, A., Widlund, O.: *Domain Decomposition Methods-Algorithms and Theory*, vol. 34. Springer (2004)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.