Elliptic Measures and Square Function Estimates on 1-Sided Chord-Arc Domains

Mingming Cao¹ · José María Martell¹ · Andrea Olivo²

Received: 26 July 2021 / Accepted: 15 October 2021 / Published online: 12 January 2022
© The Author(s) 2021

Abstract

In nice environments, such as Lipschitz or chord-arc domains, it is well-known that the solvability of the Dirichlet problem for an elliptic operator in L^p , for some finite p , is equivalent to the fact that the associated elliptic measure belongs to the Muckenhoupt class A_∞ . In turn, any of these conditions occurs if and only if the gradient of every bounded null solution satisfies a Carleson measure estimate. This has been recently extended to much rougher settings such as those of 1-sided chord-arc domains, that is, sets which are quantitatively open and connected with a boundary which is Ahlfors–David regular. In this paper, we work in the same environment and consider a qualitative analog of the latter equivalence showing that one can characterize the absolute continuity of the surface measure with respect to the elliptic measure in terms of the finiteness almost everywhere of the truncated conical square function for any bounded null solution. As a consequence of our main result particularized to the Laplace operator and some previous results, we show that the boundary of the domain is rectifiable if and only if the truncated conical square function is finite almost everywhere for any bounded harmonic function. In addition, we obtain that for two given elliptic operators L_1 and L_2 , the absolute continuity of the surface measure with

The first author is supported by Grant FJC2018-038526-I funded by MCIN/AEI/10.13039/501100011033. The first and second authors acknowledge financial support from MCIN/AEI/10.13039/501100011033 grants CEX2019-000904-S and PID2019-107914GB-I00, and from the Spanish National Research Council, through the “Ayuda extraordinaria a Centros de Excelencia Severo Ochoa” (2020SCX001). The second author also acknowledges that the research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ ERC agreement no. 615112 HAPDEGMT. The last author was supported by PIP 112201501003553 (CONICET) and has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 777822. The last author would like to express her gratitude to the first two authors and the Instituto de Ciencias Matemáticas (ICMAT), for their support and hospitality. The authors wish to thank the anonymous referee for his/her suggestions and, in particular, for pointing out a simpler argument to derive (a) $\implies$ (b) in Theorem 1.3.

✉ José María Martell
chema.martell@icmat.es

Extended author information available on the last page of the article

respect to the elliptic measure of L_1 is equivalent to the same property for L_2 provided the disagreement of the coefficients satisfy some quadratic estimate in truncated cones for almost everywhere vertex. Finally, for the case on which L_2 is either the transpose of L_1 or its symmetric part we show the equivalence of the corresponding absolute continuity upon assuming that the antisymmetric part of the coefficients has some controlled oscillation in truncated cones for almost every vertex.

Keywords Elliptic measure · Surface measure · Truncated conical square function · Rectifiability · Poisson kernel · 1-sided chord-arc domains · Absolute continuity · A_∞ Muckenhoupt weights

Mathematics Subject Classification 42B37 · 28A75 · 28A78 · 31A15 · 31B05 · 35J25 · 42B25 · 42B35

1 Introduction

A classical theorem in [41] states that

$$\omega \ll \mathcal{H}^1|_{\partial\Omega} \ll \omega \text{ on } \partial\Omega \text{ for any simply connected domain } \Omega \subset \mathbb{R}^2 \text{ with a rectifiable boundary,} \quad (1.1)$$

where ω denotes the harmonic measure relative to the domain Ω . A quantitative version of this result was obtained later by Lavrentiev [36] who showed that in a chord-arc domain in the plane, harmonic measure is quantitatively absolutely continuous with respect to the arc-length measure, that is, harmonic measure is an A_∞ weight with respect to surface measure. After these two fundamental results there has been many authors seeking to find necessary and sufficient geometric criteria for the absolute continuity, or its quantitative version, of harmonic measure with respect to surface measure on the boundary of a domain in higher dimensions. In general, those can be divided into two categories: quantitative and qualitative.

In the quantitative category it has been recently established that if $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, is a 1-sided CAD (chord-arc domain, cf. Definition 2.4), then the following are equivalent:

- (a) $\partial\Omega$ is uniformly rectifiable,
- (b) Ω satisfies the exterior corkscrew condition, hence it is a CAD, (1.2)
- (c) $\omega \in A_\infty(\sigma)$.

Here, $\sigma = \mathcal{H}^n|_{\partial\Omega}$ denotes the surface measure and $A_\infty(\sigma)$ is as mentioned above the scale-invariant version of absolute continuity. The direction (a) implies (b) was shown by Azzam et al. [7]. That (b) implies (c) was proved by David and Jerison [17], and independently by Semmes [42]. Also, (a) implies (c) was proved by Hofmann and Martell [23]. These two authors jointly with Uriarte-Tuero [27] also established that (c) implies (a). The equivalent statements in (1.2) reveal the close connection between the regularity of the boundary of a domain and the good behavior of harmonic measure

with respect to surface measure. In addition, (1.2) connects several known results, including the extension of [41] on Lipschitz domain [15], L_1^p domain [29] and BMO_1 domain [30].

For divergence form elliptic operators $Lu = -\operatorname{div}(A\nabla u)$ with real variable coefficients, that (b) implies (c) (with the elliptic measure ω_L in place of ω) was proved by Kenig and Pipher [33] under some Carleson measure estimate assumption for the matrix of coefficients A . The converse, that is, the fact that (c) implies (b) on a 1-sided CAD for the Kenig-Pipher class has been recently obtained by Hofmann et al. [24] (see also [25] for a previous result in a smaller class of operators). In another direction, it was shown in [12] that for any real (not necessarily non-symmetric) elliptic operator L , $\omega_L \in A_\infty(\sigma)$ is equivalent to the so-called Carleson measure estimates, that is, every bounded weak null solution of L satisfies Carleson measure estimates.

On the other hand, the qualitative version of (1.2) has been also studied extensively. In contrast with (1.1), some counterexamples have been presented to show how the absolute continuity of harmonic measure is indeed affected by the topology/geometry of the domain and its boundary.

- Example 1. Lavrentiev constructed in [36] a simply connected domain $\Omega \subset \mathbb{R}^2$ and a set $E \subset \partial\Omega$ such that E has zero arclength, but $\omega(E) > 0$.
- Example 2. Bishop and Jones [10] found a uniformly rectifiable set E on the plane and some subset of E with zero arc-length which carries positive harmonic measure relative to the domain $\mathbb{R}^2 \setminus E$.
- Example 3. Wu proved in [44] that there exists a topological ball $\Omega \subset \mathbb{R}^3$ and a set $E \subset \partial\Omega$ lying on a 2-dimensional hyperplane so that Hausdorff dimension of E is 1 (which implies $\sigma(E) = 0$) but $\omega(E) > 0$.
- Example 4. In [8], Azzam et al. obtained that for all $n \geq 2$, there is a Reifenberg flat domain $\Omega \subset \mathbb{R}^{n+1}$ and there is a set $E \subset \partial\Omega$ such that $\omega(E) > 0 = \sigma(E)$.

Compared with (1.1), Examples 1 and 2 indicate that both the regularity of the boundary and the connectivity of the domain seem to be necessary for absolute continuity to occur. However, Examples 3 and 4 say that $\omega \ll \sigma$ fails in the presence of some connectivity assumption. Indeed, a quantitative form of path connectedness is contained in Example 4 since Reifenberg flat domains, which are sufficiently flat, are in fact NTA domains (cf. Definition 2.4), see [34, Theorem 3.1]. Taking into consideration these, it is natural to investigate what extra mild assumptions are necessary to obtain the absolute continuity of harmonic measure.

It was shown by McMillan [37, Theorem 2] that for bounded simply connected domains $\Omega \subset \mathbb{C}$, $\omega \ll \sigma \ll \omega$ on the set of cone points. Later, Bishop and Jones [10] obtained that for any simply connected domain $\Omega \subset \mathbb{R}^2$ and curve Γ of finite length, $\omega \ll \sigma$ on $\partial\Omega \cap \Gamma$. That result refined the conclusions in [40, p. 471] and [35, Theorem 3] where Γ was a line and a quasi-smooth curve, respectively. Beyond that, in a Wiener regular domain with large complement (cf. [1, Definition 1.5]), Akman et al. [1] gave a characterization of sets of absolute continuity in terms of the cone point condition and the rectifiable structure of elliptic measure. Let us point out that in all of the just mentioned results, the absolute continuity happens locally. In the case of the whole boundary, for every Lipschitz domain Dahlberg [14] proved that harmonic measure belongs to the reverse Hölder class with exponent 2 with respect

to surface measure, this, in turn, yields $\omega \ll \sigma \ll \omega$. This was extended to the setting of CAD domains in [17, 42]. For general NTA domains $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 1$, Badger [9] proved that $\sigma \ll \omega$ if the boundary $\partial\Omega$ has finite surface measure. When Ω is a 1-sided CAD, Akman et al. established in [2] that $\partial\Omega$ is rectifiable if and only if $\sigma \ll \omega$ on $\partial\Omega$, which is also equivalent to the fact that $\partial\Omega$ possesses exterior corkscrew points in a qualitative way and that $\partial\Omega$ can be covered σ -a.e. by a countable union of portions of boundaries of bounded chord-arc subdomains of Ω . Based on a qualitative Carleson measure condition, they also got that the same conclusions hold for some class of elliptic operators with regular coefficients. The remarkable result in [6] proved that, in any dimension and in the absence of any connectivity condition, any piece of the boundary with finite surface measure is rectifiable, provided surface measure is absolutely continuous with respect to harmonic measure on that piece. The converse was treated in [3] by Akman et al. assuming that the boundary has locally finite surface measure and satisfies some weak lower Ahlfors–David regular condition.

Motivated by the previous work, the purpose of this article is to find characterizations of the absolute continuity of surface measure with respect to elliptic measure for real second order divergence form uniformly elliptic operators. Our main goal is to establish the equivalence between the absolute continuity and the finiteness almost everywhere of the conical square function applied to any bounded weak solution. To set the stage let us give few definitions (see Sect. 2 for more definitions and notation). The conical square function is defined as

$$S_\alpha u(x) := \left(\iint_{\Gamma_\alpha(x)} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY \right)^{\frac{1}{2}}, \quad x \in \partial\Omega,$$

where $\delta(\cdot) = \text{dist}(\cdot, \partial\Omega)$ and the cone $\Gamma_\alpha(x)$ with vertex at $x \in \partial\Omega$ and aperture $\alpha > 0$ is given by

$$\Gamma_\alpha(x) = \{Y \in \Omega : |Y - x| < (1 + \alpha)\delta(Y)\}.$$

Similarly, we define the truncated square function S_α^r by integrating over the truncated cone $\Gamma_\alpha^r(x) := \Gamma_\alpha(x) \cap B(x, r)$ for any $r > 0$.

Our main result is a qualitative analog of [31] and [12, Theorem 1.1]. More precisely, condition (a) is a qualitative analog of $\omega_L \in A_\infty(\sigma)$ —or equivalently $\sigma \in A_\infty(\omega_L)$ —while condition (c), or (d), or (e) is a qualitative version of the so-called Carleson measure condition, which is in turn equivalent to some local scale-invariant L^2 estimate for the truncated conical square function.

Theorem 1.3 *Let $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, be a 1-sided CAD (cf. Definition 2.4) and write $\sigma := \mathcal{H}^n|_{\partial\Omega}$. There exists $\alpha_0 > 0$ (depending only on the 1-sided CAD constants) such that for each fixed $\alpha \geq \alpha_0$ and for every real (not necessarily symmetric) elliptic operator $Lu = -\text{div}(A\nabla u)$ the following statements are equivalent:*

- (a) $\sigma \ll \omega_L$ on $\partial\Omega$.

- (b) $\partial\Omega = \bigcup_{N \geq 0} F_N$, where $\sigma(F_0) = 0$ and for each $N \geq 1$ there exists $C_N > 1$ such that

$$C_N^{-1}\sigma(F) \leq \omega_L(F) \leq C_N\sigma(F), \quad \forall F \subset F_N.$$

- (c) $\partial\Omega = \bigcup_{N \geq 0} F_N$, where $\sigma(F_0) = 0$, for each $N \geq 1$, $F_N = \partial\Omega \cap \partial\Omega_N$ for some bounded 1-sided CAD $\Omega_N \subset \Omega$, and $S_\alpha^r u \in L^2(F_N, \sigma)$ for every weak solution $u \in W_{\text{loc}}^{1,2}(\Omega) \cap L^\infty(\Omega)$ of $Lu = 0$ in Ω and for all (or for some) $r > 0$.
- (d) $S_\alpha^r u(x) < \infty$ for σ -a.e. $x \in \partial\Omega$ for every weak solution $u \in W_{\text{loc}}^{1,2}(\Omega) \cap L^\infty(\Omega)$ of $Lu = 0$ in Ω and for all (or for some) $r > 0$.
- (e) For every weak solution $u \in W_{\text{loc}}^{1,2}(\Omega) \cap L^\infty(\Omega)$ of $Lu = 0$ in Ω and for σ -a.e. $x \in \partial\Omega$ there exists $r_x > 0$ such that $S_\alpha^{r_x} u(x) < \infty$.

Remark 1.4 We would like to make the following observation regarding the parameter α in the previous statement. Note first that if one of the conditions (c), (d), or (e) holds for some $\alpha > 0$, then the same condition is automatically true for all $\alpha' \leq \alpha$. Thus, (a) or (b) implies (c), (d), or (e) holds for all $\alpha > 0$. On the other hand, for the converse implications we need to make sure that α does not get too small to prevent having empty cones, in which case the corresponding assumption trivially holds.

When turning to the harmonic measure, we obtain the following connection between the rectifiability of the boundary, the absolute continuity of surface measure with respect to harmonic measure, and the square functions estimates for harmonic functions.

Theorem 1.5 Let $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, be a 1-sided CAD and write $\sigma := \mathcal{H}^n|_{\partial\Omega}$. There exists $\alpha_0 > 0$ (depending only on the 1-sided CAD constants) such that for each fixed $\alpha \geq \alpha_0$ if we write ω to denote the harmonic measure for Ω then the following statements are equivalent:

- (a) $\partial\Omega$ is rectifiable, that is, σ -almost all of $\partial\Omega$ can be covered by a countable union of n -dimensional (possibly rotated) Lipschitz graphs.
- (b) $\sigma \ll \omega$ on $\partial\Omega$.
- (c) $S_\alpha^r u(x) < \infty$ for σ -a.e. $x \in \partial\Omega$ for every bounded harmonic function $u \in W_{\text{loc}}^{1,2}(\Omega)$ and for all (or for some) $r > 0$.

The equivalence of (a) and (b) was established in [2], while Theorem 1.3 readily gives that (b) is equivalent to (c).

As an application of Theorem 1.3, we can obtain some additional results. The first deals with perturbations (see [4,5,11,12,16,20–22,38,39]) and should be compared with its quantitative version in the 1-sided CAD setting [12, Theorem 1.3]. We note that our next result provides also a qualitative version of the work by Fefferman [21] who showed that in the unit ball if the right hand side of (1.7) is an essentially bounded function (rather than knowing that is finite almost everywhere) then one has $\omega_{L_0} \in A_\infty(\sigma)$ if and only if $\omega_{L_1} \in A_\infty(\sigma)$.

Theorem 1.6 Let $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, be a 1-sided CAD and write $\sigma := \mathcal{H}^n|_{\partial\Omega}$. There exists $\alpha_0 > 0$ (depending only on the 1-sided CAD constants) such that if the

real (not necessarily symmetric) elliptic operators $L_0u = -\operatorname{div}(A_0\nabla u)$ and $L_1u = -\operatorname{div}(A_1\nabla u)$ satisfy for some $\alpha \geq \alpha_0$ and for some $r > 0$

$$\iint_{\Gamma_\alpha^r(x)} \frac{\varrho(A_0, A_1)(X)^2}{\delta(X)^{n+1}} dX < \infty, \quad \sigma\text{-a.e. } x \in \partial\Omega, \quad (1.7)$$

where

$$\varrho(A_0, A_1)(X) := \sup_{Y \in B(X, \delta(X)/2)} |A_0(Y) - A_1(Y)|, \quad X \in \Omega,$$

then $\sigma \ll \omega_{L_0}$ if and only if $\sigma \ll \omega_{L_1}$.

Our second application of Theorem 1.3 allows us to establish a connection between the absolute continuity properties of the elliptic measures of an operator, its adjoint and/or its symmetric part. Given $Lu = -\operatorname{div}(A\nabla u)$ a real (not necessarily symmetric) elliptic operator, we let $L^\top$ denote the transpose of L , and let $L^{\operatorname{sym}} = \frac{L+L^\top}{2}$ be the symmetric part of L . These are respectively the divergence form elliptic operators with associated matrices $A^\top$ (the transpose of A) and $A^{\operatorname{sym}} = \frac{A+A^\top}{2}$. In this case, the following result is a qualitative version of [12, Theorem 1.6].

Theorem 1.8 *Let $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, be a 1-sided CAD and write $\sigma := \mathcal{H}^n|_{\partial\Omega}$. There exists $\alpha_0 > 0$ (depending only on the 1-sided CAD constants) such that if $Lu = -\operatorname{div}(A\nabla u)$ is a real (not necessarily symmetric) elliptic operator; and we assume that $(A - A^\top) \in \operatorname{Lip}_{\operatorname{loc}}(\Omega)$ and that for some $\alpha \geq \alpha_0$ and for some $r > 0$ one has*

$$\iint_{\Gamma_\alpha^r(x)} \left| \operatorname{div}_C(A - A^\top)(X) \right|^2 \delta(X)^{1-n} dX < \infty, \quad \sigma\text{-a.e. } x \in \partial\Omega, \quad (1.9)$$

where div_C stands for the column divergence, that is,

$$\operatorname{div}_C(A - A^\top)(X) = \left(\sum_{i=1}^{n+1} \partial_i (a_{i,j} - a_{j,i})(X) \right)_{1 \leq j \leq n+1}, \quad X \in \Omega;$$

then $\sigma \ll \omega_L$ if and only if $\sigma \ll \omega_{L^\top}$ if and only if $\sigma \ll \omega_{L^{\operatorname{sym}}}$.

The paper is organized as follows: In Sect. 2, we present some preliminaries, definitions, and some background results that will be used throughout the paper. Section 3 is devoted to showing Theorem 1.3. Finally, in Sect. 4, applying Theorem 1.3 (a) $\Leftrightarrow$ (d), we obtain a more general perturbation result about the absolute continuity of surface measure with respect to elliptic measure and then prove Theorems 1.6 and 1.8.

2 Preliminaries

2.1 Notation and Conventions

- Our ambient space is $\mathbb{R}^{n+1}$, $n \geq 2$.
- We use the letters c , C to denote harmless positive constants, not necessarily the same at each occurrence, which depend only on dimension and the constants appearing in the hypotheses of the theorems (which we refer to as the “allowable parameters”). We shall also sometimes write $a \lesssim b$ and $a \approx b$ to mean, respectively, that $a \leq Cb$ and $0 < c \leq a/b \leq C$, where the constants c and C are as above, unless explicitly noted to the contrary. Moreover, if c and C depend on some given parameter η , which is somehow relevant, we write $a \lesssim_\eta b$ and $a \approx_\eta b$. At times, we shall designate by M a particular constant whose value will remain unchanged throughout the proof of a given lemma or proposition, but which may have a different value during the proof of a different lemma or proposition.
- Given $E \subset \mathbb{R}^{n+1}$ we write $\text{diam}(E) = \sup_{x,y \in E} |x - y|$ to denote its diameter.
- Given a domain (i.e., open and connected) $\Omega \subset \mathbb{R}^{n+1}$, we shall use lower case letters x, y, z , etc., to denote points on $\partial\Omega$, and capital letters X, Y, Z , etc., to denote generic points in $\mathbb{R}^{n+1}$ (especially those in Ω).
- The open $(n + 1)$ -dimensional Euclidean ball of radius r will be denoted $B(x, r)$ when the center x lies on $\partial\Omega$, or $B(X, r)$ when the center $X \in \mathbb{R}^{n+1} \setminus \partial\Omega$. A “surface ball” is denoted $\Delta(x, r) := B(x, r) \cap \partial\Omega$, and unless otherwise specified it is implicitly assumed that $x \in \partial\Omega$. Also if $\partial\Omega$ is bounded, we typically assume that $0 < r \lesssim \text{diam}(\partial\Omega)$, so that $\Delta = \partial\Omega$ if $\text{diam}(\partial\Omega) < r \lesssim \text{diam}(\partial\Omega)$.
- Given a Euclidean ball B or surface ball Δ , its radius will be denoted r_B or r_Δ respectively.
- Given a Euclidean ball $B = B(X, r)$ or a surface ball $\Delta = \Delta(x, r)$, its concentric dilate by a factor of $\kappa > 0$ will be denoted by $\kappa B = B(X, \kappa r)$ or $\kappa \Delta = \Delta(x, \kappa r)$.
- For $X \in \mathbb{R}^{n+1}$, we set $\delta(X) := \text{dist}(X, \partial\Omega)$.
- We let $\mathcal{H}^n$ denote the n -dimensional Hausdorff measure, and let $\sigma := \mathcal{H}^n|_{\partial\Omega}$ denote the surface measure on $\partial\Omega$.
- For a Borel set $A \subset \mathbb{R}^{n+1}$, we let $\text{int}(A)$ denote the interior of A , and $\bar{A}$ denote the closure of A . If $A \subset \partial\Omega$, $\text{int}(A)$ will denote the relative interior, i.e., the largest relatively open set in $\partial\Omega$ contained in A . Thus, for $A \subset \partial\Omega$, the boundary is then well defined by $\partial A := \bar{A} \setminus \text{int}(A)$.
- For a Borel set $A \subset \partial\Omega$ with $0 < \sigma(A) < \infty$, we write $\int_A f d\sigma := \sigma(A)^{-1} \int_A f d\sigma$.
- We shall use the letter I (and sometimes J) to denote a closed $(n + 1)$ -dimensional Euclidean cube with sides parallel to the coordinate axes, and we let $\ell(I)$ denote the side length of I . We use Q to denote a dyadic “cube” on $\partial\Omega$. The latter exist, given that $\partial\Omega$ is Ahlfors–David regular (see [18], [13], and enjoy certain properties which we enumerate in Lemma 2.5 below).

2.2 Some Definitions

Definition 2.1 (*Ahlfors–David regular*) We say that a closed set $E \subset \mathbb{R}^{n+1}$ is n -dimensional Ahlfors–David regular (or simply ADR) if there is some uniform constant $C \geq 1$ such that

$$C^{-1}r^n \leq \mathcal{H}^n(E \cap B(x, r)) \leq Cr^n, \quad \forall x \in E, r \in (0, 2 \operatorname{diam}(E)).$$

Definition 2.2 (*Corkscrew condition*) We say that an open set $\Omega \subset \mathbb{R}^{n+1}$ satisfies the Corkscrew condition if for some uniform constant $c \in (0, 1)$, and for every surface ball $\Delta := \Delta(x, r)$ with $x \in \partial\Omega$ and $0 < r < \operatorname{diam}(\partial\Omega)$, there is a ball $B(X_\Delta, cr) \subset B(x, r) \cap \Omega$. The point $X_\Delta \in \Omega$ is called a “Corkscrew point” relative to Δ . We note that we may allow $r < C \operatorname{diam}(\partial\Omega)$ for any fixed C , simply by adjusting the constant c .

Definition 2.3 (*Harnack Chain condition*) We say that an open set Ω satisfies the Harnack Chain condition if there is a uniform constant C such that for every $\rho > 0$, $\Lambda \geq 1$, and every pair of points $X, X' \in \Omega$ with $\min\{\delta(X), \delta(X')\} \geq \rho$ and $|X - X'| < \Lambda\rho$, there is a chain of open balls $B_1, \dots, B_N \subset \Omega$, $N \leq C(\Lambda)$, with $X \in B_1$, $X' \in B_N$, $B_k \cap B_{k+1} \neq \emptyset$, $C^{-1} \operatorname{diam}(B_k) \leq \operatorname{dist}(B_k, \partial\Omega) \leq C \operatorname{diam}(B_k)$. Such a sequence is called a “Harnack Chain”.

We remark that the Corkscrew condition is a quantitative, scale invariant version of openness, and the Harnack Chain condition is a scale invariant version of path connectedness.

Definition 2.4 (*1-sided NTA domains, 1-sided CAD, NTA domains, CAD*). We say that Ω is a 1-sided NTA (non-tangentially accessible) domain if Ω satisfies both the Corkscrew and Harnack Chain conditions. Furthermore, we say that Ω is an NTA domain if it is a 1-sided NTA domain and if, in addition, $\mathbb{R}^{n+1} \setminus \overline{\Omega}$ satisfies the Corkscrew condition. If a 1-sided NTA domain, or an NTA domain, has an ADR boundary, then it is called a 1-sided CAD (chord-arc domain) or a CAD, respectively.

2.3 Dyadic Grids and Sawtooths

We give a lemma concerning the existence of a “dyadic grid”, which was proved in [13, 18, 19].

Lemma 2.5 *Suppose that $E \subset \mathbb{R}^{n+1}$ is an n -dimensional ADR set. Then there exist constants $a_0 > 0$, $\gamma > 0$, and $C_1 < 1$ depending only on n and the ADR constant such that, for each $k \in \mathbb{Z}$, there is a collection of Borel sets (cubes)*

$$\mathbb{D}_k = \left\{ Q_j^k \subset E : j \in \mathfrak{J}_k \right\}$$

where $\mathfrak{J}_k$ denotes some (possibly finite) index set depending on k , satisfying:

(a) $E = \bigcup_j Q_j^k$, for each $k \in \mathbb{Z}$.

- (b) If $m \geq k$, then either $Q_i^m \subset Q_j^k$ or $Q_i^m \cap Q_j^k = \emptyset$.
- (c) For each (j, k) and each $m < k$, there is a unique i such that $Q_j^k \subset Q_i^m$.
- (d) $\text{diam}(Q_j^k) \leq C_1 2^{-k}$.
- (e) Each Q_j^k contains some surface ball $\Delta(x_j^k, a_0 2^{-k}) := B(x_j^k, a_0 2^{-k}) \cap E$.
- (f) $\mathcal{H}^n(\{x \in Q_j^k : \text{dist}(x, E \setminus Q_j^k) \leq 2^{-k} a\}) \leq C_1 a^\gamma \mathcal{H}^n(Q_j^k)$ for all k, j and $a \in (0, a_0)$.

A few remarks are in order concerning this lemma.

- In the setting of a general space of homogeneous type, this lemma has been proved by Christ [13], with the dyadic parameter $1/2$ replaced by some constant $\delta \in (0, 1)$. In fact, one may always take $\delta = 1/2$ (cf. [28, Proof of Proposition 2.12]). In the presence of the Ahlfors-David property, the result already appears in [18, 19].
- For our purposes, we may ignore those $k \in \mathbb{Z}$ such that $2^{-k} \gtrsim \text{diam}(E)$, in the case that the latter is finite.
- We shall denote by $\mathbb{D}(E)$ the collection of all relevant Q_j^k , i.e.,

$$\mathbb{D}(E) := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_k,$$

where, if $\text{diam}(E)$ is finite, the union runs over those k such that $2^{-k} \lesssim \text{diam}(E)$.

- For a dyadic cube $Q \in \mathbb{D}_k$, we shall set $\ell(Q) = 2^{-k}$, and we shall refer to this quantity as the “length” of Q . Evidently, $\ell(Q) \simeq \text{diam}(Q)$. We set $k(Q) = k$ to be the dyadic generation to which Q belongs if $Q \in \mathbb{D}_k$; thus, $\ell(Q) = 2^{-k(Q)}$.
- Properties (d) and (e) imply that for each cube $Q \in \mathbb{D}_k$, there is a point $x_Q \in E$, a Euclidean ball $B(x_Q, r_Q)$ and a surface ball $\Delta(x_Q, r_Q) := B(x_Q, r_Q) \cap E$ such that $c\ell(Q) \leq r_Q \leq \ell(Q)$, for some uniform constant $c > 0$, and

$$\Delta(x_Q, 2r_Q) \subset Q \subset \Delta(x_Q, Cr_Q), \quad (2.6)$$

for some uniform constant $C > 1$. We shall write

$$B_Q := B(x_Q, r_Q), \quad \Delta_Q := \Delta(x_Q, r_Q), \quad \tilde{\Delta}_Q := \Delta(x_Q, Cr_Q), \quad (2.7)$$

and we shall refer to the point x_Q as the “center” of Q .

- Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set satisfying the corkscrew condition and such that $\partial\Omega$ is ADR. Given $Q \in \mathbb{D}(\partial\Omega)$, we define the “corkscrew point relative to Q ” as $X_Q := X_{\Delta_Q}$. We note that

$$\delta(X_Q) \simeq \text{dist}(X_Q, Q) \simeq \text{diam}(Q).$$

We next introduce the notation of “Carleson region” and “discretized sawtooth” from [23, Sect. 3]. Given a dyadic cube $Q \in \mathbb{D}(E)$, the “discretized Carleson region” $\mathbb{D}_Q$ relative to Q is defined by

$$\mathbb{D}_Q := \{Q' \in \mathbb{D}(E) : Q' \subset Q\}.$$

Let $\mathcal{F} = \{Q_j\} \subset \mathbb{D}(E)$ be a pairwise family of disjoint cubes. The “global discretized sawtooth” relative to $\mathcal{F}$ is the collection of cubes $Q \in \mathbb{D}(E)$ that are not contained in any $Q_j \in \mathcal{F}$, that is,

$$\mathbb{D}_{\mathcal{F}} := \mathbb{D}(E) \setminus \bigcup_{Q_j \in \mathcal{F}} \mathbb{D}_{Q_j}.$$

For a given cube $Q \in \mathbb{D}(E)$, we define the “local discretized sawtooth” relative to $\mathcal{F}$ is the collection of cubes in $\mathbb{D}_Q$ that are not contained in any $Q_j \in \mathcal{F}$ of, equivalently,

$$\mathbb{D}_{\mathcal{F}, Q} := \mathbb{D}_Q \setminus \bigcup_{Q_j \in \mathcal{F}} \mathbb{D}_{Q_j} = \mathbb{D}_{\mathcal{F}} \cap \mathbb{D}_Q.$$

We also introduce the “geometric” Carleson regions and sawtooths. In the sequel, $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, is a 1-sided CAD. Given $Q \in \mathbb{D} := \mathbb{D}(\partial\Omega)$ we want to define some associated regions which inherit the good properties of Ω . Let $\mathcal{W} = \mathcal{W}(\Omega)$ denote a collection of (closed) dyadic Whitney cubes of Ω , so that the cubes in $\mathcal{W}$ form a covering of Ω with non-overlapping interiors, which satisfy

$$4 \operatorname{diam}(I) \leq \operatorname{dist}(4I, \partial\Omega) \leq \operatorname{dist}(I, \partial\Omega) \leq 40 \operatorname{diam}(I), \quad \forall I \in \mathcal{W},$$

and also

$$(1/4) \operatorname{diam}(I_1) \leq \operatorname{diam}(I_2) \leq 4 \operatorname{diam}(I_1), \quad \text{whenever } I_1 \text{ and } I_2 \text{ touch.}$$

Let $X(I)$ be the center of I and $\ell(I)$ denote the sidelength of I .

Given $0 < \lambda < 1$ and $I \in \mathcal{W}$, we write $I^* = (1 + \lambda)I$ for the “fattening” of I . By taking λ small enough, we can arrange matters, so that for any $I, J \in \mathcal{W}$,

$$\begin{aligned} \operatorname{dist}(I^*, J^*) &\simeq \operatorname{dist}(I, J), \\ \operatorname{int}(I^*) \cap \operatorname{int}(J^*) &\neq \emptyset \iff \partial I \cap \partial J \neq \emptyset. \end{aligned}$$

(The fattening thus ensures overlap of I^* and J^* for any pair $I, J \in \mathcal{W}$ whose boundaries touch, so that the Harnack chain property then holds locally, with constants depending upon λ , in $I^* \cap J^*$.) By choosing λ sufficiently small, say $0 < \lambda < \lambda_0$, we may also suppose that there is a $\tau \in (1/2, 1)$ such that for distinct $I, J \in \mathcal{W}$, we have that $\tau J \cap I^* = \emptyset$. In what follows we will need to work with the dilations $I^{**} = (1 + 2\lambda)I$ or $I^{***} = (1 + 4\lambda)I$, and to ensure that the same properties hold we further assume that $0 < \lambda < \lambda_0/4$.

Given $\vartheta \in \mathbb{N}$, for every cube $Q \in \mathbb{D}$ we set

$$\mathcal{W}_Q^\vartheta := \{I \in \mathcal{W} : 2^{-\vartheta} \ell(Q) \leq \ell(I) \leq 2^\vartheta \ell(Q), \text{ and } \operatorname{dist}(I, Q) \leq 2^\vartheta \ell(Q)\}. \quad (2.8)$$

We will choose $\vartheta \geq \vartheta_0$, with ϑ_0 large enough depending on the constants of the corkscrew condition (cf. Definition 2.2) and in the dyadic cube construction (cf. Lemma 2.5), so that $X_Q \in I$ for some $I \in \mathcal{W}_Q^\vartheta$, and for each dyadic child Q^j of Q , the respective corkscrew points $X_{Q^j} \in I^j$ for some $I^j \in \mathcal{W}_{Q^j}^\vartheta$. Moreover, we may always find an $I \in \mathcal{W}_Q^\vartheta$ with the slightly more precise property that $\ell(Q)/2 \leq \ell(I) \leq \ell(Q)$ and

$$\mathcal{W}_{Q_1}^\vartheta \cap \mathcal{W}_{Q_2}^\vartheta \neq \emptyset, \quad \text{whenever } 1 \leq \frac{\ell(Q_2)}{\ell(Q_1)} \leq 2, \text{ and } \text{dist}(Q_1, Q_2) \leq 1000\ell(Q_2).$$

For each $I \in \mathcal{W}_Q^\vartheta$, we form a Harnack chain from the center $X(I)$ to the corkscrew point X_Q and call it $H(I)$. We now let $\mathcal{W}_Q^{\vartheta,*}$ denote the collection of all Whitney cubes which meet at least one ball in the Harnack chain $H(I)$ with $I \in \mathcal{W}_Q^\vartheta$, that is,

$$\mathcal{W}_Q^{\vartheta,*} := \{J \in \mathcal{W} : \text{there exists } I \in \mathcal{W}_Q^\vartheta \text{ such that } H(I) \cap J \neq \emptyset\}.$$

We also define

$$U_Q^\vartheta := \bigcup_{I \in \mathcal{W}_Q^{\vartheta,*}} (1 + \lambda)I =: \bigcup_{I \in \mathcal{W}_Q^{\vartheta,*}} I^*.$$

By construction, we then have that

$$\mathcal{W}_Q^\vartheta \subset \mathcal{W}_Q^{\vartheta,*} \subset \mathcal{W} \quad \text{and} \quad X_Q \in U_Q^\vartheta, \quad X_{Q^j} \in U_Q^\vartheta,$$

for each child Q^j of Q . It is also clear that there is a uniform constant k^* (depending only on the 1-sided CAD constants and ϑ) such that

$$\begin{aligned} 2^{-k^*} \ell(Q) &\leq \ell(I) \leq 2^{k^*} \ell(Q), \quad \forall I \in \mathcal{W}_Q^{\vartheta,*}, \\ X(I) &\rightarrow_{U_Q^\vartheta} X_Q, \quad \forall I \in \mathcal{W}_Q^{\vartheta,*}, \\ \text{dist}(I, Q) &\leq 2^{k^*} \ell(Q), \quad \forall I \in \mathcal{W}_Q^{\vartheta,*}. \end{aligned}$$

Here, $X(I) \rightarrow_{U_Q^\vartheta} X_Q$ means that the interior of U_Q^ϑ contains all balls in Harnack Chain (in Ω) connecting $X(I)$ to X_Q , and moreover, for any point Z contained in any ball in the Harnack Chain, we have $\text{dist}(Z, \partial\Omega) \simeq \text{dist}(Z, \Omega \setminus U_Q^\vartheta)$ with uniform control of implicit constants. The constant k^* and the implicit constants in the condition $X(I) \rightarrow_{U_Q^\vartheta} X_Q$, depend on at most allowable parameter, on λ , and on ϑ . Moreover, given $I \in \mathcal{W}$ we have that $I \in \mathcal{W}_{Q_I}^{\vartheta,*}$, where $Q_I \in \mathbb{D}(\partial\Omega)$ satisfies $\ell(Q_I) = \ell(I)$, and contains any fixed $\hat{y} \in \partial\Omega$ such that $\text{dist}(I, \partial\Omega) = \text{dist}(I, \hat{y})$. The reader is referred to [23] for full details. We note however that in that reference the parameter ϑ is fixed. Here we need to allow ϑ to depend on the aperture of the cones and hence it is convenient to include the superindex ϑ .

For a given $Q \in \mathbb{D}$, the “Carleson box” relative to Q is defined by

$$T_Q^\vartheta := \text{int} \left(\bigcup_{Q' \in \mathbb{D}_Q} U_{Q'}^\vartheta \right).$$

For a given family $\mathcal{F} = \{Q_j\}$ of pairwise disjoint cubes and a given $Q \in \mathbb{D}(\partial\Omega)$, we define the “local sawtooth region” relative to $\mathcal{F}$ by

$$\Omega_{\mathcal{F}, Q}^\vartheta := \text{int} \left(\bigcup_{Q' \in \mathbb{D}_{\mathcal{F}, Q}} U_{Q'}^\vartheta \right) = \text{int} \left(\bigcup_{I \in \mathcal{W}_{\mathcal{F}, Q}^\vartheta} I^* \right),$$

where $\mathcal{W}_{\mathcal{F}, Q}^\vartheta := \bigcup_{Q' \in \mathbb{D}_{\mathcal{F}, Q}} \mathcal{W}_{Q'}^{\vartheta, *}$. Analogously, we can slightly fatten the Whitney boxes and use I^{**} to define new fattened Whitney regions and sawtooth domains. More precisely, for every $Q \in \mathbb{D}(\partial\Omega)$,

$$T_Q^{\vartheta, *} := \text{int} \left(\bigcup_{Q' \in \mathbb{D}_Q} U_{Q'}^\vartheta \right), \quad \Omega_{\mathcal{F}, Q}^{\vartheta, *} := \text{int} \left(\bigcup_{Q' \in \mathbb{D}_{\mathcal{F}, Q}} U_{Q'}^{\vartheta, *} \right), \quad U_Q^{\vartheta, *} := \bigcup_{I \in \mathcal{W}_Q^{\vartheta, *}} I^{**}.$$

Similarly, we can define $T_Q^{\vartheta, **}$, $\Omega_{\mathcal{F}, Q}^{\vartheta, **}$ and $U_Q^{\vartheta, **}$ by using I^{***} in place of I^{**} . For later use, we recall that [23, Proposition 6.1]:

$$Q \setminus \left(\bigcup_{Q_j \in \mathcal{F}} Q_j \right) \subset \partial\Omega \cap \partial\Omega_{\mathcal{F}, Q}^\vartheta \subset \overline{Q} \setminus \left(\bigcup_{Q_j \in \mathcal{F}} \text{int}(Q_j) \right). \quad (2.9)$$

Following [23], one can easily see that there exist constants $0 < \kappa_1 < 1$ and $\kappa_0 \geq \max\{2C, 4/c\}$ (with C the constant in (2.6), and c such that $c\ell(Q) \leq r_Q$), depending only on the allowable parameters and on ϑ , so that

$$\kappa_1 B_Q \cap \Omega \subset T_Q^\vartheta \subset T_Q^{\vartheta, *} \subset T_Q^{\vartheta, **} \subset \overline{T_Q^{\vartheta, **}} \subset \kappa_0 B_Q \cap \overline{\Omega} =: \frac{1}{2} B_Q^* \cap \overline{\Omega}, \quad (2.10)$$

where B_Q is defined as in (2.7).

2.4 PDE Estimates

Now, we recall several facts concerning the elliptic measures and the Green functions. For our first results we will only assume that $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, is an open set, not necessarily connected, with $\partial\Omega$ being ADR. Later we will focus on the case where Ω is a 1-sided CAD.

Let $Lu = -\text{div}(A\nabla u)$ be a variable coefficient second order divergence form operator with $A(X) = (a_{i,j}(X))_{i,j=1}^{n+1}$ being a real (not necessarily symmetric) matrix

with $a_{i,j} \in L^\infty(\Omega)$ for $1 \leq i, j \leq n+1$, and A uniformly elliptic, that is, there exists $\Lambda \geq 1$ such that

$$\Lambda^{-1}|\xi|^2 \leq A(X)\xi \cdot \xi, \quad |A(X)\xi \cdot \eta| \leq \Lambda|\xi||\eta|, \quad \forall \xi, \eta \in \mathbb{R}^{n+1} \text{ and a.e. } X \in \Omega.$$

In what follows, we will only be working with this kind of operators and we will refer to them as “elliptic operators” for the sake of simplicity. We write $L^\top$ to denote the transpose of L , or, in other words, $L^\top u = -\operatorname{div}(A^\top \nabla u)$ with $A^\top$ being the transpose matrix of A .

We say that a function $u \in W_{\text{loc}}^{1,2}(\Omega)$ is a weak solution of $Lu = 0$ in Ω , or that $Lu = 0$ in the weak sense, if

$$\iint_{\Omega} A(X) \nabla u(X) \cdot \nabla \phi(X) = 0, \quad \forall \phi \in C_c^\infty(\Omega).$$

Here and elsewhere $C_c^\infty(\Omega)$ stands for the set of compactly supported smooth functions with all derivatives of all orders being continuous.

Associated with the operators L and $L^\top$, one can, respectively, construct the elliptic measures $\{\omega_L^X\}_{X \in \Omega}$ and $\{\omega_{L^\top}^X\}_{X \in \Omega}$, and the Green functions G_L and $G_{L^\top}$ (see [26] for full details). We next present some definitions and properties that will be used throughout this paper.

The following lemmas can be found in [26].

Lemma 2.11 *Suppose that $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, is an open set such that $\partial\Omega$ is ADR. Given an elliptic operator L , there exist $C > 1$ (depending only on dimension and on the ellipticity of L) and $c_\theta > 0$ (depending on the above parameters and on $\theta \in (0, 1)$) such that G_L , the Green function associated with L , satisfies*

$$G_L(X, Y) \leq C|X - Y|^{1-n}; \quad (2.12)$$

$$c_\theta|X - Y|^{1-n} \leq G_L(X, Y), \quad \text{if } |X - Y| \leq \theta\delta(X), \theta \in (0, 1); \quad (2.13)$$

$$G_L(\cdot, Y) \in C(\overline{\Omega} \setminus \{Y\}) \quad \text{and } G_L(\cdot, Y)|_{\partial\Omega} \equiv 0, \forall Y \in \Omega; \quad (2.14)$$

$$G_L(X, Y) \geq 0, \quad \forall X, Y \in \Omega, X \neq Y; \quad (2.15)$$

$$G_L(X, Y) = G_{L^\top}(Y, X), \quad \forall X, Y \in \Omega, X \neq Y; \quad (2.16)$$

Moreover, $G_L(\cdot, Y) \in W_{\text{loc}}^{1,2}(\Omega \setminus \{Y\})$ for every $Y \in \Omega$, and satisfies $LG_L(\cdot, Y) = \delta_Y$ in the weak sense in Ω , that is,

$$\iint_{\Omega} A(X) \nabla_X G_L(X, Y) \cdot \nabla \Phi(X) dX = \Phi(Y), \quad \forall \Phi \in C_c^\infty(\Omega). \quad (2.17)$$

Finally, the following Riesz formula holds

$$\iint_{\Omega} A^\top(Y) \nabla_Y G_{L^\top}(Y, X) \cdot \nabla \Phi(Y) dY = \Phi(X) - \int_{\partial\Omega} \Phi d\omega_L^X, \quad (2.18)$$

for a.e. $X \in \Omega$ and for every $\Phi \in C_c^\infty(\mathbb{R}^{n+1})$.

Lemma 2.19 Suppose that $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, is a 1-sided CAD. Let L be an elliptic operator. There exists a constant C (depending only on the dimension, the 1-sided CAD constants and the ellipticity of L) such that for every ball $B_0 := B(x_0, r_0)$ with $x_0 \in \partial\Omega$ and $0 < r_0 < \text{diam}(\partial\Omega)$, and $\Delta_0 = B_0 \cap \partial\Omega$ we have the following properties:

(a) There holds

$$\omega_L^Y(\Delta_0) \geq 1/C, \quad \forall Y \in \Omega \cap B(x_0, C^{-1}r_0). \quad (2.20)$$

(b) If $B = B(x, r)$ with $x \in \partial\Omega$ is such that $2B \subset B_0$, then for any $X \in \Omega \setminus B_0$,

$$C^{-1}\omega_L^X(\Delta) \leq r^{n-1}G_L(X, X_\Delta) \leq C\omega_L^X(\Delta). \quad (2.21)$$

(b) If $X \in \Omega \setminus 4B_0$, then we have

$$\omega_L^X(2\Delta_0) \leq C\omega_L^X(\Delta_0). \quad (2.22)$$

3 Proof of Theorem 1.3

The goal of this section is to prove Theorem 1.3. We start with the following observation which will be used throughout the paper:

Remark 3.1 For every $\alpha > 0$, $0 < r < r'$, and $\varpi \in \mathbb{R}$, if $F \subset \partial\Omega$ is a bounded set and $v \in L_{\text{loc}}^2(\Omega)$, then

$$\sup_{x \in F} \iint_{\Gamma_\alpha^{r'}(x) \setminus \Gamma_\alpha^r(x)} |v(Y)|^2 \delta(Y)^\varpi dY < \infty. \quad (3.2)$$

To see this we first note that since F is bounded we can find R large enough so that $F \subset B(0, R)$. Then, if $x \in F$ one readily sees that

$$\Gamma_\alpha^{r'}(x) \setminus \Gamma_\alpha^r(x) \subset \overline{B(0, r' + R)} \cap \left\{ Y \in \Omega : \frac{r}{1+\alpha} \leq \delta(Y) \leq r' \right\} =: K.$$

Note that $K \subset \Omega$ is a compact set. Then, since $v \in L_{\text{loc}}^2(\Omega)$, we conclude that

$$\sup_{x \in F} \iint_{\Gamma_\alpha^{r'}(x) \setminus \Gamma_\alpha^r(x)} |v(Y)|^2 \delta(Y)^\varpi dY \leq \max \left\{ r', \frac{1+\alpha}{r} \right\}^{|\varpi|} \iint_K |v(Y)|^2 dY < \infty. \quad (3.3)$$

We can now proceed to prove Theorem 1.3. We first note that it is immediate to see that (b) $\implies$ (a), (c) $\implies$ (d), and (d) $\implies$ (e). Moreover, (3.2) yields easily (e) $\implies$ (d). Thus, it suffices to prove the following implications:

$$(a) \implies (c), \quad (a) \implies (b), \quad \text{and} \quad (d) \implies (a).$$

3.1 Proof of (a) $\implies$ (c)

Assume that $\sigma \ll \omega_L$. Fix an arbitrary $Q_0 \in \mathbb{D}_{k_0}$ where $k_0 \in \mathbb{Z}$ is taken so that $2^{-k_0} = \ell(Q_0) < \text{diam}(\partial\Omega)/M_0$, where M_0 is large enough and will be chosen later. From the construction of $T_{Q_0}^\vartheta$ one can easily see that $T_{Q_0}^\vartheta \subset \frac{1}{2}B_{Q_0}^* := \kappa_0 B_{Q_0}$, see (2.10). Let $X_0 := X_{M_0\Delta_{Q_0}}$ be an interior corkscrew point relative to $M_0\Delta_{Q_0}$ so that $X_0 \notin 4B_{Q_0}^*$ provided that M_0 is taken large enough depending on the allowable parameters. Since $\partial\Omega$ is ADR, (2.20) and Harnack's inequality give that $\omega_L^{X_0}(Q_0) \geq C_0^{-1}$, where $C_0 > 1$ depends on 1-sided CAD constants and M_0 . We now normalize the elliptic measure and the Green function as follows

$$\omega := C_0\sigma(Q_0)\omega_L^{X_0} \quad \text{and} \quad \mathcal{G}(\cdot) := C_0\sigma(Q_0)G_L(X_0, \cdot). \quad (3.4)$$

The hypothesis $\sigma \ll \omega_L$ implies that $\sigma \ll \omega$. Note that $1 \leq \frac{\omega(Q_0)}{\sigma(Q_0)} \leq C_0$. Let $N > C_0$ and let $\mathcal{F}_N^+ := \{Q_j\} \subset \mathbb{D}_{Q_0} \setminus \{Q_0\}$, respectively, $\mathcal{F}_N^- := \{Q_j\} \subset \mathbb{D}_{Q_0} \setminus \{Q_0\}$, be the collection of descendants of Q_0 which are maximal (and therefore pairwise disjoint) with respect to the property that

$$\frac{\omega(Q_j)}{\sigma(Q_j)} < \frac{1}{N}, \quad \text{respectively} \quad \frac{\omega(Q_j)}{\sigma(Q_j)} > N. \quad (3.5)$$

Write $\mathcal{F}_N = \mathcal{F}_N^+ \cup \mathcal{F}_N^-$ and note that $\mathcal{F}_N^+ \cap \mathcal{F}_N^- = \emptyset$. By maximality, one has

$$\frac{1}{N} \leq \frac{\omega(Q)}{\sigma(Q)} \leq N, \quad \forall Q \in \mathbb{D}_{\mathcal{F}_N, Q_0}. \quad (3.6)$$

Write for every $N > C_0$,

$$E_N^\pm := \bigcup_{Q \in \mathcal{F}_N^\pm} Q, \quad E_N^0 = E_N^+ \cup E_N^-, \quad E_N := Q_0 \setminus E_N^0, \quad (3.7)$$

and

$$Q_0 = \left(\bigcap_{N > C_0} E_N^0 \right) \cup \left(\bigcup_{N > C_0} E_N \right) =: E_0 \cup \left(\bigcup_{N > C_0} E_N \right). \quad (3.8)$$

We claim that for every $N > C_0$

$$E_N^+ = \{x \in Q_0 : M_{Q_0, \omega}^d \sigma(x) > N\} \quad \text{and} \quad E_N^- = \{x \in Q_0 : M_{Q_0, \sigma}^d \omega(x) > N\}, \quad (3.9)$$

where given two non-negative Borel measures μ and ν we set

$$M_{Q_0, \mu}^d \nu(x) := \sup_{x \in Q \in \mathbb{D}_{Q_0}} \frac{\nu(Q)}{\mu(Q)}.$$

To see the first equality in (3.9), let $x \in E_N^+$. Then, there exists $Q_j \in \mathcal{F}_N^+ \subset \mathbb{D}_{Q_0}$ so that $Q_j \ni x$. Thus, by (3.5)

$$M_{Q_0, \omega}^d \sigma(x) \geq \frac{\sigma(Q_j)}{\omega(Q_j)} > N.$$

On the other hand, if $M_{Q_0, \omega}^d \sigma(x) > N$, there exists $Q \in \mathbb{D}_{Q_0}$ so that $\sigma(Q)/\omega(Q) > N$. By the maximality of $\mathcal{F}_N^+$ we therefore conclude that $Q \subset Q_j$ for some $Q_j \in \mathcal{F}_N^+$. Hence, $x \in E_N^+$ as desired. This completes the proof of the first equality in (3.9) and the second one follows using the same argument interchanging the roles of ω and σ .

Once we have shown (3.9), we clearly see that $\{E_N^+\}_N$, $\{E_N^-\}_N$, and $\{E_N^0\}_N$ are decreasing sequences of sets. This, together with the fact that $\omega(E_N^\pm) \leq \omega(Q_0) < \infty$ and $\sigma(E_N^\pm) \leq \sigma(Q_0) < \infty$, implies that

$$\omega\left(\bigcap_{N>C_0} E_N^\pm\right) = \lim_{N \rightarrow \infty} \omega(E_N^\pm), \quad \sigma\left(\bigcap_{N>C_0} E_N^\pm\right) = \lim_{N \rightarrow \infty} \sigma(E_N^\pm). \quad (3.10)$$

Our next goal is to show that $\sigma(E_0) = 0$. To see this we note that by (3.5)

$$\omega(E_N^+) = \sum_{Q \in \mathcal{F}_N^+} \omega(Q) < \frac{1}{N} \sum_{Q \in \mathcal{F}_N^+} \sigma(Q) = \frac{1}{N} \sigma(E_N^+) \leq \frac{1}{N} \sigma(Q_0)$$

and, by (3.10)

$$\omega\left(\bigcap_{N>C_0} E_N^+\right) = \lim_{N \rightarrow \infty} \omega(E_N^+) = 0.$$

Use this, the fact that $\sigma \ll \omega$, and (3.10) to derive

$$0 = \sigma\left(\bigcap_{N>C_0} E_N^+\right) = \lim_{N \rightarrow \infty} \sigma(E_N^+). \quad (3.11)$$

On the other hand, (3.5) yields

$$\sigma(E_N^-) = \sum_{Q \in \mathcal{F}_N^-} \sigma(Q) < \frac{1}{N} \sum_{Q \in \mathcal{F}_N^-} \omega(Q) = \frac{1}{N} \omega(E_N^-) \leq \frac{1}{N} \omega(Q_0),$$

and (3.10) implies

$$\sigma\left(\bigcap_{N>C_0} E_N^-\right) = \lim_{N \rightarrow \infty} \sigma(E_N^-) = 0. \quad (3.12)$$

All these, together with (3.10) and the fact that $\{E_N^0\}_N$ is a decreasing sequence of sets with $\sigma(E_N^0) \leq \sigma(Q_0) < \infty$, give

$$\sigma(E_0) = \lim_{N \rightarrow \infty} \sigma(E_N^0) \leq \lim_{N \rightarrow \infty} \sigma(E_N^+) + \lim_{N \rightarrow \infty} \sigma(E_N^-) = 0, \quad (3.13)$$

hence $\sigma(E_0) = 0$.

Next, by (2.9) and [23, Proposition 6.3], we have

$$E_N \subset F_N := \partial\Omega \cap \partial\Omega_{\mathcal{F}_N, Q_0}^\vartheta \quad \text{and} \quad \sigma(F_N \setminus E_N) = 0. \quad (3.14)$$

Note that [23, Lemma 3.61] yields that $\Omega_{\mathcal{F}_N, Q_0}^\vartheta$ is a bounded 1-sided CAD for any $\vartheta \geq \vartheta_0$. Now, we are going to bound the square function in $L^2(F_N, \sigma)$. Let $u \in W_{\text{loc}}^{1,2}(\Omega) \cap L^\infty(\Omega)$ be a weak solution of $Lu = 0$ in Ω . Let $\vartheta \geq \vartheta_0$ and note that by (2.10), we see that $2B_Q \subset B_{Q_0}^*$. Recalling (3.4) and the fact $X_0 \notin 4B_{Q_0}^*$, we use (2.21), (2.22), (3.6), Harnack's inequality, and the fact that $\partial\Omega$ is ADR to conclude that

$$\frac{\mathcal{G}(X)}{\delta(X)} \simeq \frac{\omega(Q)}{\sigma(Q)} \simeq_N 1, \quad (3.15)$$

for all $X \in I^*$ with $I \in \mathcal{W}_Q^{\vartheta,*}$ and $Q \in \mathbb{D}_{\mathcal{F}_N, Q_0}$. This and the definition of $\Omega_{\mathcal{F}_N, Q_0}^\vartheta$ yield

$$\iint_{\Omega_{\mathcal{F}_N, Q_0}^\vartheta} |\nabla u(Y)|^2 \delta(Y) dY \lesssim_N \iint_{\Omega_{\mathcal{F}_N, Q_0}^\vartheta} |\nabla u(Y)|^2 \mathcal{G}(Y) dY. \quad (3.16)$$

For every $M \geq 1$, we set $\mathcal{F}_{N,M}$ to be the family of maximal cubes of the collection $\mathcal{F}_N$ augmented by adding all the cubes $Q \in \mathbb{D}_{Q_0}$ such that $\ell(Q) \leq 2^{-M} \ell(Q_0)$. This means that $Q \in \mathbb{D}_{\mathcal{F}_{N,M}, Q_0}$ if and only if $Q \in \mathbb{D}_{\mathcal{F}_N, Q_0}$ and $\ell(Q) > 2^{-M} \ell(Q_0)$. Observe that $\mathbb{D}_{\mathcal{F}_{N,M}, Q_0} \subset \mathbb{D}_{\mathcal{F}_{N,M'}, Q_0}$ for all $M \leq M'$, and hence $\Omega_{\mathcal{F}_{N,M}, Q_0}^\vartheta \subset \Omega_{\mathcal{F}_{N,M'}, Q_0}^\vartheta \subset \Omega_{\mathcal{F}_N, Q_0}^\vartheta$. This, together with the monotone convergence theorem, gives

$$\iint_{\Omega_{\mathcal{F}_N, Q_0}^\vartheta} |\nabla u(Y)|^2 \mathcal{G}(Y) dY = \lim_{M \rightarrow \infty} \iint_{\Omega_{\mathcal{F}_{N,M}, Q_0}^\vartheta} |\nabla u(Y)|^2 \mathcal{G}(Y) dY. \quad (3.17)$$

Invoking [12, Proposition 3.58], one has

$$\iint_{\Omega_{\mathcal{F}_{N,M}, Q_0}^\vartheta} |\nabla u(Y)|^2 \mathcal{G}(Y) dY \lesssim_N \sigma(Q_0) \simeq 2^{-k_0 n}, \quad (3.18)$$

where the implicit constants are independent of M . Consequently, combining (3.16), (3.17) and (3.18), we deduce that

$$\iint_{\Omega_{\mathcal{F}_N, Q_0}^\vartheta} |\nabla u(Y)|^2 \delta(Y) dY \leq C_N. \quad (3.19)$$

To continue, we recall the dyadic square function defined in [27, Sect. 2.3]:

$$S_{Q_0}^\vartheta u(x) := \left(\iint_{\Gamma_{Q_0}^\vartheta(x)} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY \right)^{1/2}, \text{ where } \Gamma_{Q_0}^\vartheta(x) := \bigcup_{x \in Q \in \mathbb{D}_{Q_0}} U_Q^\vartheta.$$

Note that if $Q \in \mathbb{D}_{Q_0}$ is so that $Q \cap E_N \neq \emptyset$, then necessarily $Q \in \mathbb{D}_{\mathcal{F}_N, Q_0}$, otherwise, $Q \subset Q' \in \mathcal{F}_N$, hence $Q \subset Q_0 \setminus E_N$. In view of (3.19), we have

$$\begin{aligned} \int_{E_N} S_{Q_0}^\vartheta u(x)^2 d\sigma(x) &= \int_{E_N} \iint_{\bigcup_{x \in Q \in \mathbb{D}_{Q_0}} U_Q^\vartheta} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY d\sigma(x) \\ &\lesssim \sum_{Q \in \mathbb{D}_{Q_0}} \ell(Q)^{-n} \sigma(Q \cap E_N) \iint_{U_Q^\vartheta} |\nabla u(Y)|^2 \delta(Y) dY \\ &\lesssim \sum_{Q \in \mathbb{D}_{\mathcal{F}_N, Q_0}} \iint_{U_Q^\vartheta} |\nabla u(Y)|^2 \delta(Y) dY \\ &\lesssim \iint_{\Omega_{\mathcal{F}_N, Q_0}^\vartheta} |\nabla u(Y)|^2 \delta(Y) dY \leq C_N, \end{aligned} \quad (3.20)$$

where we have used that the family $\{U_Q^\vartheta\}_{Q \in \mathbb{D}}$ has bounded overlap. This along with the last condition in (3.14) yields

$$S_{Q_0}^\vartheta u \in L^2(F_N, \sigma), \quad \forall \vartheta \geq \vartheta_0. \quad (3.21)$$

We next claim that fixed $\alpha > 0$, we can find ϑ sufficiently large depending on α such that for any $r_0 \ll 2^{-k_0}$,

$$S_\alpha^{r_0} u(x) \leq S_{Q_0}^\vartheta u(x), \quad x \in Q_0. \quad (3.22)$$

It suffices to show $\Gamma_\alpha^{r_0}(x) \subset \Gamma_{Q_0}^\vartheta(x)$ for any $x \in Q_0$. Indeed, let $Y \in \Gamma_\alpha^{r_0}(x)$. Pick $I \in \mathcal{W}$ so that $Y \in I$, and hence, $\ell(I) \simeq \delta(Y) \leq |Y - x| < r_0 \ll 2^{-k_0} = \ell(Q_0)$. Pick $Q_I \in \mathbb{D}_{Q_0}$ such that $x \in Q_I$ and $\ell(Q_I) = \ell(I) \ll \ell(Q_0)$. Thus, one has

$$\text{dist}(I, Q_I) \leq |Y - x| < (1 + \alpha)\delta(Y) \leq C(1 + \alpha)\ell(I) = C(1 + \alpha)\ell(Q_I).$$

Recalling (2.8), if we take $\vartheta \geq \vartheta_0$ large enough so that

$$2^\vartheta \geq C(1 + \alpha), \quad (3.23)$$

then $Y \in I \in \mathcal{W}_{Q_I}^\vartheta \subset \mathcal{W}_{Q_I}^{\vartheta,*}$. The latter gives that $Y \in U_{Q_I}^\vartheta \subset \Gamma_{Q_0}^\vartheta(x)$ and consequently (3.22) holds. We should mention that the dependence of ϑ on α implies that all the sawtooth regions $\Omega_{\mathcal{F}_N, Q_0}^\vartheta$ above as well as all the implicit constants depend on α .

To complete the proof we note that, it follows from (3.21) and (3.22) that $S_\alpha^{r_0} u \in L^2(F_N, \sigma)$. This together with Remark 3.1 easily yields

$$S_\alpha^r u \in L^2(F_N, \sigma), \quad \text{for any } r > 0. \quad (3.24)$$

We note that the previous argument has been carried out for an arbitrary $Q_0 \in \mathbb{D}_{k_0}$. Hence, using (3.7), (3.8), and (3.14) with $Q_k \in \mathbb{D}_{k_0}$, we conclude, with the induced notation, that

$$\begin{aligned} \partial\Omega &= \bigcup_{Q_k \in \mathbb{D}_{k_0}} Q_k = \left(\bigcup_{Q_k \in \mathbb{D}_{k_0}} E_0^k \right) \cup \left(\bigcup_{Q_k \in \mathbb{D}_{k_0}} \bigcup_{N > C_0} E_N^k \right) \\ &= \left(\bigcup_{Q_k \in \mathbb{D}_{k_0}} E_0^k \right) \cup \left(\bigcup_{Q_k \in \mathbb{D}_{k_0}} \bigcup_{N > C_0} F_N^k \right) =: F_0 \cup \left(\bigcup_{k, N} F_N^k \right), \end{aligned} \quad (3.25)$$

where $\sigma(F_0) = 0$ and $F_N^k = \partial\Omega \cap \partial\Omega_{\mathcal{F}_N^k, Q_k}$ where each $\Omega_{\mathcal{F}_N^k, Q_k} \subset \Omega$ is a bounded 1-sided CAD. Combining (3.25) and (3.24) with F_N^k in place of F_N , the proof of (a) $\Rightarrow$ (c) is complete. $\square$

3.2 Proof of (a) $\Rightarrow$ (b)

We use an argument suggested by the anonymous referee. Fix $X_0 \in \Omega$ and write $\omega := \omega_L^{X_0}$. Our assumption $\sigma \ll \omega_L$ implies that there exists a non-negative ω -measurable function $h = d\sigma/d\omega \in L_{\text{loc}}^1(\omega)$ (we are implicitly assuming that h is a fixed element of the equivalence class of functions which agree with $d\sigma/d\omega$ ω -a.e., hence we may assume that h is defined everywhere in $\partial\Omega$). Set

$$F_0 := F_0^0 \cup F_0^\infty := \{x \in \partial\Omega : h(x) = 0\} \cup \{x \in \partial\Omega : h(x) = \infty\}$$

and

$$F_N := \{x \in \partial\Omega : N^{-1} \leq h(x) \leq N\}, \quad N \in \mathbb{N}.$$

Clearly, $\partial\Omega = \bigcup_{N \geq 0} F_N$. Since $h \in L_{\text{loc}}^1(\omega_L)$ we have that $\omega(F_0^\infty) = 0$, hence $\sigma(F_0^\infty) = 0$. Also

$$\sigma(F_0^0) = \int_{F_0^0} h \, d\omega = 0.$$

Hence $\sigma(F_0) = 0$. On the other hand, if $F \subset F_N$, $N \in \mathbb{N}$ we clearly have

$$\frac{1}{N} \omega(F) \leq \sigma(F) = \int_F h \, d\omega \leq N \omega(F).$$

This completes the proof. $\square$

Remark 3.26 Using the proof of (a) $\implies$ (c) one can provide an alternative approach to see that (a) $\implies$ (b). For that we borrow some ideas from [43] and address some small inaccuracies that do not affect their conclusion. As before we fix an arbitrary cube $Q_0 \in \mathbb{D}_{k_0}$ and an integer $N > C_0$. Recall that the family $\mathcal{F}_N$ of stopping cubes is constructed in (3.5) and $E_N^0 = E_N^+ \cup E_N^-$ defined in (3.7). As we have shown that $\{E_N^+\}_N$, $\{E_N^-\}_N$, and $\{E_N^0\}_N$ are decreasing sequence of sets it is easy to see that

$$\bigcap_{N > C_0} E_N^0 = \left(\bigcap_{N > C_0} E_N^+ \right) \cup \left(\bigcap_{N > C_0} E_N^- \right). \quad (3.27)$$

By (3.11), (3.12) we conclude that this set has null σ -measure. Recalling our assumption $\sigma \ll \omega$ we set $h = d\sigma/d\omega$ and

$$L_0 := \left\{ x \in Q_0 : \int_{Q_x} |h(y) - h(x)| \, d\omega(y) \rightarrow 0, \mathbb{D}_{Q_0} \ni Q_x \searrow \{x\} \right\}. \quad (3.28)$$

By the Lebesgue differentiation theorem for dyadic cubes it follows that $\omega(Q_0 \setminus L_0) = 0$, hence $\sigma(Q_0 \setminus L_0) = 0$. Then we can write

$$Q_0 = \left(\bigcup_{N > C_0} (L_0 \setminus E_N^0) \right) \cup \left(\bigcap_{N > C_0} E_N^0 \right) \cup (Q_0 \setminus L_0) =: \left(\bigcup_{N > C_0} (L_0 \setminus E_N^0) \right) \cup \widetilde{F}_0,$$

with $\sigma(\widetilde{F}_0) = 0$. We claim that for any $x \in L_0 \setminus E_N^0$ the following holds:

$$\frac{1}{N} \leq \frac{\sigma(Q)}{\omega(Q)} \leq N, \quad \forall Q \in \mathbb{D}_{Q_0}, Q \ni x. \quad (3.29)$$

Otherwise, by the maximality of $\mathcal{F}_N^+$ or $\mathcal{F}_N^-$, one has $Q \subset Q_j$ for some $Q_j \in \mathcal{F}_N^+ \cup \mathcal{F}_N^-$. Hence $x \in E_N^0$ by (3.7) which is a contradiction. Using (3.28) and (3.29) we then obtain that $N^{-1} \leq h(x) \leq N$ for all $x \in L_0 \setminus E_N^0$. Thus, for every $F \subset L_0 \setminus E_N^0$ we conclude that

$$\frac{1}{N} \omega(F) \leq \sigma(F) = \int_F h \, d\omega \leq N \omega(F).$$

To complete the proof we repeat this argument with any $Q_k \in \mathbb{D}_{k_0}$ to readily get (b) using that $\partial\Omega = \bigcup_{Q_k \in \mathbb{D}_{k_0}} Q_k$.

3.3 Proof of (d) $\implies$ (a)

Given $Q_0 \in \mathbb{D}$ and for any $\eta \in (0, 1)$, we define the modified dyadic square function

$$S_{Q_0}^{\vartheta_0, \eta} u(x) := \left(\iint_{\Gamma_{Q_0}^{\vartheta_0, \eta}(x)} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY \right)^{1/2},$$

where the modified non-tangential cone $\Gamma_{Q_0}^{\vartheta_0, \eta}(x)$ is given by

$$\Gamma_{Q_0}^{\vartheta_0, \eta}(x) := \bigcup_{x \in Q \in \mathbb{D}_{Q_0}} U_{Q, \eta^3}^{\vartheta_0}, \quad U_{Q, \eta^3}^{\vartheta_0} = \bigcup_{\substack{Q' \in \mathbb{D}_Q \\ \ell(Q') > \eta^3 \ell(Q)}} U_{Q'}^{\vartheta_0}.$$

Here we recall that ϑ_0 depends on the 1-sided CAD constants (see Sect. 2.3).

The following auxiliary result, whose proof is postponed to Appendix 1, extends [12, Lemma 3.10] (see also [31, 32]).

Lemma 3.30 *There exist $0 < \eta \ll 1$ (depending only on dimension, the 1-sided CAD constants and the ellipticity of L) such that for every $Q_0 \in \mathbb{D}$, and for every Borel set $\emptyset \neq F \subset Q_0$ satisfying $\omega_L^{X_{Q_0}}(F) = 0$, there exists a Borel set $S \subset Q_0$ such that the bounded weak solution $u(X) = \omega_L^X(S)$ satisfies*

$$S_{Q_0}^{\vartheta_0, \eta} u(x) = \infty, \quad \forall x \in F.$$

Assume that (d) holds. To prove that $\sigma \ll \omega_L$ on $\partial\Omega$, by Lemma 2.5 (a), it suffices to show that for any given $Q_0 \in \mathbb{D}$,

$$F \subset Q_0, \quad \omega_L(F) = 0 \implies \sigma(F) = 0. \quad (3.31)$$

Consider then $F \subset Q_0$ with $\omega_L(F) = 0$. By the mutually absolute continuity between elliptic measures, one has $\omega_L^{X_{Q_0}}(F) = 0$. Lemma 3.30 applied to F yields that there exists a Borel set $S \subset Q_0$ such that $u(X) = \omega_L^X(S)$, $X \in \Omega$, satisfies

$$S_{Q_0}^{\vartheta_0, \eta} u(x) = \infty, \quad \forall x \in F. \quad (3.32)$$

To continue, we claim that there exist $\alpha_0 > 0$ and $r > 0$ such that

$$\Gamma_{Q_0}^{\vartheta_0, \eta}(x) \subset \Gamma_{\alpha}^r(x), \quad \forall x \in Q_0 \text{ and } \forall \alpha \geq \alpha_0. \quad (3.33)$$

Indeed, let $Y \in \Gamma_{Q_0}^{\vartheta_0, \eta}(x)$. By definition, there exist $Q \in \mathbb{D}_{Q_0}$ and $Q' \in \mathbb{D}_Q$ with $\ell(Q') > \eta^3 \ell(Q)$ such that $Y \in U_{Q'}^{\vartheta_0}$ and $x \in Q$. Then $Y \in I^*$ for some $I \in \mathcal{W}_{Q'}^{\vartheta_0, *}$, and hence,

$$\delta(Y) \simeq \ell(I) \simeq \ell(Q') \leq \ell(Q) < \eta^{-3} \ell(Q'). \quad (3.34)$$

This further implies that

$$\begin{aligned} |Y - x| &\leq \text{diam}(I^*) + \text{dist}(I, x) \leq \text{diam}(I^*) + \text{dist}(I, Q') + \text{diam}(Q) \\ &\lesssim 2^{k^*} \ell(Q') + \ell(Q) \lesssim \ell(Q), \end{aligned} \quad (3.35)$$

where k^* depends on the 1-sided CAD constants (see Sect. 2.3). Combining (3.34) with (3.35), we get

$$|Y - x| \leq C_1 \ell(Q_0) =: r/2 \quad \text{and} \quad |Y - x| \leq (1 + C_{1,\eta}) \delta(Y) =: (1 + \alpha_0) \delta(Y),$$

where C_1 depends only on the allowable parameters, and $C_{1,\eta}$ depends only on the allowable parameters and also on η . Eventually, this justifies (3.33).

Combining (3.32), (3.33), and (d), we readily see that $\sigma(F) = 0$ and (3.31) follows. This completes the proof of (d) $\implies$ (a) and hence that of Theorem 1.3. $\square$

4 Proof of Theorems 1.6 and 1.8

To prove Theorems 1.6 and 1.8, we will make use of Theorem 1.3 and show that the truncated square function is finite σ -a.e. for every bounded weak solution. Indeed, we are going to show the following more general result, which is a qualitative version of [12, Theorem 4.13].

Theorem 4.1 *Let $\Omega \subset \mathbb{R}^{n+1}$, $n \geq 2$, be a 1-sided CAD. There exists $\tilde{\alpha}_0 > 0$ (depending only on the 1-sided CAD constants) such that if $L_0 u = -\text{div}(A_0 \nabla u)$ and $L_1 u = -\text{div}(A_1 \nabla u)$ are real (not necessarily symmetric) elliptic operators such that $A_0 - A_1 = A + D$, where $A, D \in L^\infty(\Omega)$ are real matrices satisfying the following conditions:*

(i) *there exist $\alpha_1 \geq \tilde{\alpha}_0$ and $r_1 > 0$ such that*

$$\iint_{\Gamma_{\alpha_1}^{r_1}(x)} a(X)^2 \delta(X)^{-n-1} dX < \infty, \quad \sigma\text{-a.e. } x \in \partial\Omega, \quad (4.2)$$

where

$$a(X) := \sup_{Y \in B(X, \delta(X)/2)} |A(Y)|, \quad X \in \Omega;$$

(ii) *$D \in \text{Lip}_{\text{loc}}(\Omega)$ is antisymmetric and there exist $\alpha_2 \geq \tilde{\alpha}_0$ and $r_2 > 0$ such that*

$$\iint_{\Gamma_{\alpha_2}^{r_2}(x)} |\text{div}_C D(X)|^2 \delta(X)^{1-n} dX < \infty, \quad \sigma\text{-a.e. } x \in \partial\Omega; \quad (4.3)$$

then $\sigma \ll \omega_{L_0}$ if and only if $\sigma \ll \omega_{L_1}$.

Proof By symmetry, it suffices to assume that $\sigma \ll \omega_{L_0}$ and prove $\sigma \ll \omega_{L_1}$. Let $u \in W_{\text{loc}}^{1,2}(\Omega) \cap L^\infty(\Omega)$ be a weak solution of $L_1 u = 0$ in Ω and $\|u\|_{L^\infty(\Omega)} = 1$. Applying Theorem 1.3 (d) $\Rightarrow$ (a) to u , we are reduced to showing that for some $r > 0$,

$$S_{\alpha_0}^r u(x) < \infty, \quad \text{for } \sigma\text{-a.e. } x \in \partial\Omega,$$

where α_0 is given in Theorem 1.3. Proceeding as in Sect. 3.1 and invoking (3.22), it suffices to see that for every fixed $Q_0 \in \mathbb{D}_{k_0}$ and for some fixed large ϑ (which depends on α_0 and hence solely on the 1-sided CAD constants) one has

$$Q_0 = \bigcup_{N \geq 0} \widehat{E}_N, \quad \sigma(\widehat{E}_0) = 0 \quad \text{and} \quad S_{Q_0}^\vartheta u \in L^2(\widehat{E}_N, \sigma), \quad \forall N \geq 1. \quad (4.4)$$

Fix then $Q_0 \in \mathbb{D}_{k_0}$, where k_0 is given in the beginning of Sect. 3.1. We use the normalization in (3.4) with $L = L_0$:

$$\omega := C_0 \sigma(Q_0) \omega_{L_0}^{X_0} \quad \text{and} \quad \mathcal{G}(\cdot) := C_0 \sigma(Q_0) G_{L_0}(X_0, \cdot). \quad (4.5)$$

Much as in (3.5), we introduce the families for each $N > C_0$ we let $\mathcal{F}_N^+ := \{Q_j\} \subset \mathbb{D}_{Q_0} \setminus \{Q_0\}$, respectively, $\mathcal{F}_N^- := \{Q_j\} \subset \mathbb{D}_{Q_0} \setminus \{Q_0\}$, be the collection of descendants of Q_0 which are maximal (and therefore pairwise disjoint) with respect to the property that

$$\frac{\omega(Q_j)}{\sigma(Q_j)} < \frac{1}{N}, \quad \text{respectively} \quad \frac{\omega(Q_j)}{\sigma(Q_j)} > N. \quad (4.6)$$

Write $\mathcal{F}_N = \mathcal{F}_N^+ \cup \mathcal{F}_N^-$ and note that $\mathcal{F}_N^+ \cap \mathcal{F}_N^- = \emptyset$. By maximality, one has

$$\frac{1}{N} \leq \frac{\omega(Q)}{\sigma(Q)} \leq N, \quad \forall Q \in \mathbb{D}_{\mathcal{F}_N, Q_0}. \quad (4.7)$$

Write for every $N > C_0$,

$$E_N^\pm := \bigcup_{Q \in \mathcal{F}_N^\pm} Q, \quad E_N^0 = E_N^+ \cup E_N^-, \quad E_N := Q_0 \setminus E_N^0. \quad (4.8)$$

Set

$$S_{Q_0} \gamma^\vartheta(x) := \left(\sum_{x \in Q \in \mathbb{D}_{Q_0}} \gamma_Q^\vartheta \right)^{1/2},$$

where for every $Q \in \mathbb{D}_{Q_0}$ we write

$$\gamma_Q^\vartheta := \iint_{U_Q^{\vartheta,*}} a(X)^2 \delta(X)^{-n-1} dX + \iint_{U_Q^{\vartheta,*}} |\operatorname{div}_C D(X)|^2 \delta(X)^{1-n} dX.$$

We claim that there exist $\tilde{\alpha}_0 > 0$ and $\tilde{r} > 0$ such that

$$\Gamma_{Q_0}^{\vartheta,*}(x) := \bigcup_{x \in Q \in \mathbb{D}_{Q_0}} U_Q^{\vartheta,*} \subset \Gamma_{\tilde{\alpha}_0}^{\tilde{r}}(x), \quad x \in \partial\Omega. \quad (4.9)$$

Indeed, let $Y \in \Gamma_{Q_0}^{\vartheta,*}(x)$. Then, there exists $Q \in \mathbb{D}_{Q_0}$ with $Q \ni x$ and $I \in \mathcal{W}_Q^{\vartheta,*}$ such that $Y \in I^{**}$. Using these, one has

$$|Y - x| \leq \text{diam}(I^{**}) + \text{dist}(I, Q) + \text{diam}(Q) \lesssim_{\vartheta} \ell(Q) \simeq_{\vartheta} \ell(I) \simeq \delta(Y),$$

which implies

$$|Y - x| < C_1 \delta(Y) =: (1 + \tilde{\alpha}_0) \delta(Y) \quad \text{and} \quad |Y - x| < C_2 \ell(Q_0) = C_2 2^{-k_0} =: \tilde{r},$$

where both C_1 and C_2 depend only on the allowable parameters —note that they depend on ϑ , hence on the 1-sided CAD constants. Thus, (4.9) holds for the choice of $\tilde{\alpha}_0$ and $\tilde{r}$, and as a result

$$\begin{aligned} S_{Q_0} \gamma^{\vartheta}(x)^2 &\lesssim \iint_{\Gamma_{Q_0}^{\vartheta,*}(x)} a(X)^2 \delta(X)^{-n-1} dX + \iint_{\Gamma_{Q_0}^{\vartheta,*}(x)} |\text{div}_C D(X)|^2 \delta(X)^{1-n} dX \\ &\leq \iint_{\Gamma_{\alpha_1}^{\max\{\tilde{r}, r_1\}}(x)} a(X)^2 \delta(X)^{-n-1} dX + \iint_{\Gamma_{\alpha_2}^{\max\{\tilde{r}, r_2\}}(x)} |\text{div}_C D(X)|^2 \delta(X)^{1-n} dX \\ &< \infty, \quad \text{for } \sigma\text{-a.e. } x \in Q_0, \end{aligned} \quad (4.10)$$

where we have used that the fact that the family $\{U_Q^{\vartheta,*}\}_{Q \in \mathbb{D}}$ has bounded overlap, that $\alpha_1, \alpha_2 \geq \tilde{\alpha}_0$ and the last estimate follows from (4.2), (4.3) together with Remark 3.1.

Given $N > C_0$ (C_0 is the constant that appeared in Sect. 3.1), let $\tilde{\mathcal{F}}_N \subset \mathbb{D}_{Q_0}$ be the collection of maximal cubes (with respect to the inclusion) $Q_j \in \mathbb{D}_{Q_0}$ such that

$$\sum_{Q_j \subset Q \in \mathbb{D}_{Q_0}} \gamma_Q^{\vartheta} > N^2. \quad (4.11)$$

Observe that

$$S_{Q_0} \gamma^{\vartheta}(x) \leq N, \quad \forall x \in Q_0 \setminus \bigcup_{Q_j \in \tilde{\mathcal{F}}_N} Q_j. \quad (4.12)$$

Otherwise, there exists a cube $Q_x \ni x$ such that $\sum_{Q_x \subset Q \in \mathbb{D}_{Q_0}} \gamma_Q^{\vartheta} > N^2$, hence $x \in Q_x \subset Q_j$ for some $Q_j \in \tilde{\mathcal{F}}_N$, which is a contradiction.

We next set

$$\tilde{E}_0 := \bigcap_{N > C_0} \tilde{E}_N^0 := \bigcap_{N > C_0} \left(\bigcup_{Q_j \in \tilde{\mathcal{F}}_N} Q_j \right). \quad (4.13)$$

Let $x \in \tilde{E}_{N+1}^0$. Then there exists $Q_x \in \tilde{\mathcal{F}}_{N+1}$ such that $x \in Q_x$. By (4.11), one has

$$\sum_{Q_x \subset Q \in \mathbb{D}_{Q_0}} \gamma_Q^\vartheta > (N+1)^2 > N^2.$$

Therefore, the maximality of the cubes in $\tilde{\mathcal{F}}_N$ gives that $Q_x \subset Q'_x$ for some $Q'_x \in \tilde{\mathcal{F}}_N$ with $x \in Q'_x \subset \tilde{E}_N^0$. This shows that $\{\tilde{E}_N^0\}_N$ is a decreasing sequence of sets, and since $\tilde{E}_N^0 \subset Q_0$ for every N we conclude that

$$\omega(\tilde{E}_0) = \lim_{N \rightarrow \infty} \omega(\tilde{E}_N^0), \quad \sigma(\tilde{E}_0) = \lim_{N \rightarrow \infty} \sigma(\tilde{E}_N^0).$$

Note that for every $N > C_0$, if $x \in \tilde{E}_0$ there exists $Q_x^N \in \tilde{\mathcal{F}}_N$ such that $Q_x^N \ni x$. By the definition of $\mathcal{F}_N$, we have

$$S_{Q_0} \gamma^\vartheta(x)^2 = \sum_{x \in Q \in \mathbb{D}_{Q_0}} \gamma_Q^\vartheta \geq \sum_{Q_x^N \subset Q \in \mathbb{D}_{Q_0}} \gamma_Q^\vartheta > N^2,$$

and, therefore,

$$\begin{aligned} \sigma(\tilde{E}_0) &= \lim_{N \rightarrow \infty} \sigma(\tilde{E}_N^0) \leq \lim_{N \rightarrow \infty} \sigma(\{x \in Q_0 : S_{Q_0} \gamma^\vartheta(x) > N\}) \\ &= \sigma(\{x \in Q_0 : S_{Q_0} \gamma^\vartheta(x) = \infty\}) = 0, \end{aligned} \quad (4.14)$$

by (4.10).

To proceed, let $\hat{\mathcal{F}}_N$ be the collection of maximal, hence pairwise disjoint, cubes in $\mathcal{F}_N \cup \tilde{\mathcal{F}}_N$. Note that $\mathbb{D}_{\hat{\mathcal{F}}_N, Q_0} \subset \mathbb{D}_{\mathcal{F}_N, Q_0} \cap \mathbb{D}_{\tilde{\mathcal{F}}_N, Q_0}$. This along with (4.7) yields

$$\frac{1}{N} \leq \frac{\omega(Q)}{\sigma(Q)} \leq N, \quad \forall Q \in \mathbb{D}_{\hat{\mathcal{F}}_N, Q_0}. \quad (4.15)$$

We next set

$$\hat{E}_0 := \bigcap_{N > C_0} \hat{E}_N^0 := \bigcap_{N > C_0} \left(\bigcup_{Q_j \in \hat{\mathcal{F}}_N} Q_j \right). \quad (4.16)$$

Note that $\hat{\mathcal{F}}_N \subset \mathcal{F}_N \cup \tilde{\mathcal{F}}_N$ and also that if $Q \in \mathcal{F}_N \cup \tilde{\mathcal{F}}_N$ then there exists $Q' \in \mathcal{F}_N \cup \tilde{\mathcal{F}}_N$ so that $Q \subset Q'$. This shows that $\hat{E}_N^0 = E_N^0 \cup \tilde{E}_N^0$, where E_N^0 and $\tilde{E}_N^0$ are defined in (4.8) and (4.13) respectively. As we showed that $\{E_N^0\}_N$ and $\{\tilde{E}_N^0\}_N$ are decreasing sequence of sets, then so is $\{\hat{E}_N^0\}_N$. This together with the fact that $\hat{E}_N^0 \subset Q_0$ lead to

$$\sigma(\hat{E}_0) = \lim_{N \rightarrow \infty} \sigma(\hat{E}_N^0) \leq \lim_{N \rightarrow \infty} \sigma(E_N^0) + \lim_{N \rightarrow \infty} \sigma(\tilde{E}_N^0) = 0,$$

as shown in (3.13) and (4.14), hence $\sigma(\hat{E}_0) = 0$.

Next we write

$$Q_0 = \widehat{E}_0 \cup \left(\bigcup_{N > C_0} \widehat{E}_N \right) := \widehat{E}_0 \cup \left(\bigcup_{N > C_0} (Q_0 \setminus \widehat{E}_N^0) \right). \quad (4.17)$$

Therefore, to get (4.4), we are left with proving

$$S_{Q_0}^\vartheta u \in L^2(\widehat{E}_N, \sigma), \quad \forall N > C_0. \quad (4.18)$$

With this goal in mind, we apply (4.15), (4.17) and proceed as in the proof of (3.20) and (3.16), to conclude that

$$\int_{\widehat{E}_N} S_{Q_0}^\vartheta u(x)^2 d\sigma(x) \lesssim \iint_{\Omega_{\widehat{\mathcal{F}}_N, Q_0}^\vartheta} |\nabla u|^2 \delta dY \lesssim_N \iint_{\Omega_{\widehat{\mathcal{F}}_N, Q_0}^\vartheta} |\nabla u|^2 \mathcal{G} dY. \quad (4.19)$$

As in Sect. 3.1, for every $M \geq 1$, we consider the pairwise disjoint collection $\widehat{\mathcal{F}}_{N,M}$ that is the family of maximal cubes of the collection $\widehat{\mathcal{F}}_N$ augmented by adding all the cubes $Q \in \mathbb{D}_{Q_0}$ such that $\ell(Q) \leq 2^{-M}\ell(Q_0)$. In particular, $Q \in \mathbb{D}_{\widehat{\mathcal{F}}_{N,M}, Q_0}$ if and only if $Q \in \mathbb{D}_{\widehat{\mathcal{F}}_N, Q_0}$ and $\ell(Q) > 2^{-M}\ell(Q_0)$. Moreover, $\mathbb{D}_{\widehat{\mathcal{F}}_{N,M}, Q_0} \subset \mathbb{D}_{\widehat{\mathcal{F}}_{N,M'}, Q_0}$ for all $M \leq M'$, and hence $\Omega_{\widehat{\mathcal{F}}_{N,M}, Q_0}^\vartheta \subset \Omega_{\widehat{\mathcal{F}}_{N,M'}, Q_0}^\vartheta \subset \Omega_{\widehat{\mathcal{F}}_N, Q_0}^\vartheta$. Then the monotone convergence theorem implies

$$\iint_{\Omega_{\widehat{\mathcal{F}}_N, Q_0}^\vartheta} |\nabla u|^2 \mathcal{G} dY = \lim_{M \rightarrow \infty} \iint_{\Omega_{\widehat{\mathcal{F}}_{N,M}, Q_0}^\vartheta} |\nabla u|^2 \mathcal{G} dY =: \lim_{M \rightarrow \infty} \mathcal{J}_M. \quad (4.20)$$

To continue with the proof, we are going to follow [12, Proof of Proposition 4.18]. Let $\Psi \in C_c^\infty(\mathbb{R}^{n+1})$ be the smooth cut-off function associated with the sawtooth domain $\Omega_{\widehat{\mathcal{F}}_{N,M}, Q_0}^\vartheta$ (see [12, Lemma 3.61] or [25, Lemma 4.44]) and note that since $\Psi \gtrsim 1$ in $\Omega_{\widehat{\mathcal{F}}_N, M}^\vartheta$ we have

$$\mathcal{J}_M \lesssim \widetilde{\mathcal{J}}_M := \iint_{\Omega} |\nabla u|^2 \mathcal{G} \Psi^2 dY. \quad (4.21)$$

Note that $\widetilde{\mathcal{J}}_M < \infty$ because $\text{supp } \Psi \subset \overline{\Omega_{\widehat{\mathcal{F}}_{N,M}, Q_0}^{\vartheta,*}} \subset \Omega$ and $u \in W_{\text{loc}}^{1,2}(\Omega)$. A careful examination of [12, Proof of Proposition 4.18] gives

$$\begin{aligned} \widetilde{\mathcal{J}}_M &\lesssim_N \sigma(Q_0) + \widetilde{\mathcal{J}}_M^{\frac{1}{2}} \left(\iint_{\Omega_{\widehat{\mathcal{F}}_{N,M}, Q_0}^{\vartheta,*}} \frac{a(X)^2}{\delta(X)} dX \right)^{\frac{1}{2}} \\ &\quad + \widetilde{\mathcal{J}}_M^{\frac{1}{2}} \sigma(Q_0)^{\frac{1}{2}} + \sigma(Q_0)^{\frac{1}{2}} \left(\iint_{\Omega_{\widehat{\mathcal{F}}_{N,M}, Q_0}^{\vartheta,*}} |\text{div}_C D(X)|^2 \delta(X) dX \right)^{\frac{1}{2}}. \end{aligned}$$

In turn, applying Young's inequality and hiding, we readily get

$$\tilde{\mathcal{I}}_M \lesssim_N \sigma(Q_0) + \iint_{\Omega_{\tilde{\mathcal{F}}_N, Q_0}^{\vartheta, *}} \frac{a(X)^2}{\delta(X)} dX + \iint_{\Omega_{\tilde{\mathcal{F}}_N, Q_0}^{\vartheta, *}} |\operatorname{div}_C D(X)|^2 \delta(X) dX, \quad (4.22)$$

where the implicit constant is independent of M . Collecting (4.19), (4.20), (4.21), and (4.22), we obtain

$$\begin{aligned} \int_{\widehat{E}_N} S_{Q_0}^{\vartheta} u(x)^2 d\sigma(x) &\lesssim \sigma(Q_0) + \iint_{\Omega_{\tilde{\mathcal{F}}_N, Q_0}^{\vartheta, *}} \frac{a(X)^2}{\delta(X)} dX + \iint_{\Omega_{\tilde{\mathcal{F}}_N, Q_0}^{\vartheta, *}} |\operatorname{div}_C D(X)|^2 \delta(X) dX \\ &\leq \sigma(Q_0) + \sum_{Q \in \mathbb{D}_{\tilde{\mathcal{F}}_N, Q_0}} \left(\iint_{U_Q^{\vartheta, *}} \frac{a(X)^2}{\delta(X)} dX + \iint_{U_Q^{\vartheta, *}} |\operatorname{div}_C D(X)|^2 \delta(X) dX \right) \\ &\lesssim \sigma(Q_0) + \sum_{Q \in \mathbb{D}_{\tilde{\mathcal{F}}_N, Q_0}} \gamma_Q^{\vartheta} \sigma(Q), \end{aligned} \quad (4.23)$$

where we used that $\sigma(Q) \simeq \ell(Q)^n \simeq \delta(X)^n$ for every $X \in U_Q^{\vartheta, *}$. On the other hand,

$$\begin{aligned} \sum_{Q \in \mathbb{D}_{\tilde{\mathcal{F}}_N, Q_0}} \gamma_Q^{\vartheta} \sigma(Q) &= \int_{Q_0} \sum_{x \in Q \in \mathbb{D}_{\tilde{\mathcal{F}}_N, Q_0}} \gamma_Q^{\vartheta} d\sigma(x) \\ &\leq \int_{\widehat{E}_N} S_{Q_0} \gamma^{\vartheta}(x)^2 d\sigma(x) + \sum_{Q_j \in \tilde{\mathcal{F}}_N} \sum_{Q \in \mathbb{D}_{\tilde{\mathcal{F}}_N, Q_0}} \gamma_Q^{\vartheta} \sigma(Q \cap Q_j). \end{aligned} \quad (4.24)$$

As observed above $\widetilde{E}_N^0 \subset \widehat{E}_N^0$, hence, (4.12) leads to

$$\int_{\widehat{E}_N} S_{Q_0} \gamma^{\vartheta}(x)^2 d\sigma(x) \leq N^2 \sigma(Q_0). \quad (4.25)$$

In order to control the second term in (4.24), we fix $Q_j \in \widehat{\mathcal{F}}_N$. Note that if $Q \in \mathbb{D}_{\widehat{\mathcal{F}}_N, Q_0}$ is so that $Q \cap Q_j \neq \emptyset$ then necessarily $Q_j \subsetneq Q$. Write $\widehat{Q}_j$ for the dyadic father of Q_j , that is, $\widehat{Q}_j$ is the unique dyadic cube containing Q_j with $\ell(\widehat{Q}_j) = 2\ell(Q_j)$. We claim that

$$\sum_{\widehat{Q}_j \subset Q \in \mathbb{D}_{Q_0}} \gamma_Q^{\vartheta} = \sum_{Q_j \subsetneq Q \in \mathbb{D}_{Q_0}} \gamma_Q^{\vartheta} \leq N^2. \quad (4.26)$$

Otherwise, recalling the construction of $\tilde{\mathcal{F}}_N$ in (4.11), it follows that $\widehat{Q}_j \subset Q'$ for some $Q' \in \tilde{\mathcal{F}}_N$. From the definition of $\widehat{\mathcal{F}}_N$, we then have that $Q' \subset Q''$ for some

$Q'' \in \widehat{\mathcal{F}}_N$. Consequently, $Q_j \subsetneq Q''$ with $Q_j, Q'' \in \widehat{\mathcal{F}}_N$ contradicting the maximality of the family $\widehat{\mathcal{F}}_N$. Then it follows from (4.26) that

$$\begin{aligned} \sum_{Q_j \in \widehat{\mathcal{F}}_N} \sum_{Q \in \mathbb{D}_{\widehat{\mathcal{F}}_N, Q_0}} \gamma_Q^\vartheta \sigma(Q \cap Q_j) &= \sum_{Q_j \in \widehat{\mathcal{F}}_N} \sigma(Q_j) \sum_{Q_j \subsetneq Q \in \mathbb{D}_{Q_0}} \gamma_Q^\vartheta \\ &\leq N^2 \sum_{Q_j \in \widehat{\mathcal{F}}_N} \sigma(Q_j) \leq N^2 \sigma\left(\bigcup_{Q_j \in \widehat{\mathcal{F}}_N} Q_j\right) \leq N^2 \sigma(Q_0). \end{aligned} \quad (4.27)$$

Collecting (4.23), (4.24), (4.25), and (4.27), we deduce that

$$\int_{\widehat{E}_N} S_{Q_0} u(x)^2 d\sigma(x) \leq C_N \sigma(Q_0) \simeq C_N 2^{-k_0 n}.$$

This shows (4.18) and completes the proof of Theorem 4.1. $\square$

Now let us see how we deduce Theorems 1.6 and 1.8 from Theorem 4.1.

Proof of Theorem 1.6 Let L_0 and L_1 be the elliptic operators given in Theorem 1.6. If we take $A = A_0 - A_1$ and $D = 0$ in Theorem 4.1, then (4.2) coincides with the assumption (1.7) and (4.3) holds automatically. Therefore, Theorem 1.6 immediately follows from Theorem 4.1. $\square$

Proof of Theorem 1.8 Let A be the matrix as stated in Theorem 1.8. If we take $A_0 = A$, $A_1 = A^\top$, $\widetilde{A} = 0$ and $D = A - A^\top$ in Theorem 4.1, then one has $A_0 - A_1 = \widetilde{A} + D$ with $D \in \text{Lip}_{\text{loc}}(\Omega)$ antisymmetric, (4.2) holds trivially and (4.3) agrees with (1.9). Thus, Theorem 4.1 implies that $\sigma \ll \omega_L$ if and only if $\sigma \ll \omega_{L^\top}$.

Similarly, the conclusion that $\sigma \ll \omega_L$ if and only if $\sigma \ll \omega_{L^{\text{sym}}}$ follows if we set $A_0 = A$, $A_1 = (A + A^\top)/2$, $\widetilde{A} = 0$ and $D = (A - A^\top)/2$. $\square$

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix A. Extending the Construction of Kenig, Kirchheim, Pipher, Toro: Proof of Lemma 3.30

In this appendix we prove Lemma 3.30. We will follow the construction in [12, Sect. 3] which in turn extends that of [31] (see also [32]). In those scenarios the set F is

sufficiently small, that is, it satisfies $\omega_L^{X_{Q_0}}(F) \leq \beta \omega_L^{X_{Q_0}}(Q_0)$ and it is shown that there is a set S_β so that $u_\beta(X) = \omega_L^X(S_\beta)$, $X \in \Omega$, satisfies $S_{Q_0}^{\beta_0, \eta} u(x)^2 \gtrsim_\eta \log(\beta^{-1})$ for every $x \in F$. Here we obtain the limiting case $\beta = 0$.

We start with some definition and some auxiliary result:

Definition A.1 Let $E \subset \mathbb{R}^{n+1}$ be an n -dimensional ADR set. Fix $Q_0 \in \mathbb{D}(E)$ (cf. Lemma (2.5)) and let μ be a regular Borel measure on Q_0 . Given $\varepsilon_0 \in (0, 1)$ and a Borel set $\emptyset \neq F \subset Q_0$, a good ε_0 -cover of F with respect to μ , of length $k \in \mathbb{N}$, is a collection $\{\mathcal{O}_\ell\}_{\ell=1}^k$ of Borel subsets of Q_0 , together with pairwise disjoint families $\mathcal{F}_\ell = \{Q^\ell\} \subset \mathbb{D}_{Q_0}$, $1 \leq \ell \leq k$, such that the following hold:

- (a) $F \subset \mathcal{O}_k \subset \mathcal{O}_{k-1} \subset \cdots \subset \mathcal{O}_2 \subset \mathcal{O}_1 \subset Q_0$.
- (b) $\mathcal{O}_\ell = \bigcup_{Q^\ell \in \mathcal{F}_\ell} Q^\ell$, for every $1 \leq \ell \leq k$.
- (c) $\mu(\mathcal{O}_\ell \cap Q^{\ell-1}) \leq \varepsilon_0 \mu(Q^{\ell-1})$, for each $Q^{\ell-1} \in \mathcal{F}_{\ell-1}$ and $2 \leq \ell \leq k$.

Analogously, a good ε_0 -cover of F with respect to μ , of length ∞ , is a collection $\{\mathcal{O}_\ell\}_{\ell=1}^\infty$ of Borel subsets of Q_0 , together with pairwise disjoint families $\mathcal{F}_\ell = \{Q^\ell\} \subset \mathbb{D}_{Q_0}$, $\ell \geq 1$, such that the following hold:

- (a) $F \subset \cdots \subset \mathcal{O}_k \subset \mathcal{O}_{k-1} \subset \cdots \subset \mathcal{O}_2 \subset \mathcal{O}_1 \subset Q_0$.
- (b) $\mathcal{O}_\ell = \bigcup_{Q^\ell \in \mathcal{F}_\ell} Q^\ell$, for every $\ell \geq 1$,
- (c) $\mu(\mathcal{O}_\ell \cap Q^{\ell-1}) \leq \varepsilon_0 \mu(Q^{\ell-1})$, for each $Q^{\ell-1} \in \mathcal{F}_{\ell-1}$ and $\ell \geq 2$.

Remark A.2 In the previous definition we implicitly assume that $F \cap Q^\ell \neq \emptyset$ for every $Q^\ell \in \mathcal{F}_\ell$ and for all $1 \leq \ell \leq k$ if the length is k , or all $\ell \geq 1$ if the length is infinity. Otherwise, we can remove all the cubes Q^ℓ for which $F \cap Q^\ell = \emptyset$, and all the required conditions clearly hold.

Observe also that if $\{\mathcal{O}_\ell\}_{\ell=1}^k$ is a good ε_0 -cover of F then, by Definition A.1, we have for every $2 \leq \ell \leq k$

$$\mu(\mathcal{O}_\ell^1) = \mu(\mathcal{O}_\ell^1 \cap \mathcal{O}_{\ell-1}^1) = \sum_{Q \in \mathcal{F}_{\ell-1}^1} \mu(\mathcal{O}_\ell^1 \cap Q) \leq \varepsilon_0 \sum_{Q \in \mathcal{F}_{\ell-1}^1} \mu(Q) = \varepsilon_0 \mu(\mathcal{O}_{\ell-1}^1).$$

Iterating this for every $2 \leq \ell \leq k$ we conclude that

$$\mu(\mathcal{O}_\ell^1) \leq \varepsilon_0 \mu(\mathcal{O}_{\ell-1}^1) \leq \varepsilon_0^2 \mu(\mathcal{O}_{\ell-2}^1) \leq \cdots \leq \varepsilon_0^{\ell-1} \mu(\mathcal{O}_1^1) \leq \varepsilon_0^{\ell-1} \mu(Q_0). \quad (\text{A.3})$$

Lemma A.4 [12, Lemma 3.5] Let $E \subset \mathbb{R}^{n+1}$ be an n -dimensional ADR set. Let μ be a regular Borel measure on Q_0 and assume that it is dyadically doubling on Q_0 , that is, there exists $C_\mu \geq 1$ such that $\mu(Q^*) \leq C_\mu \mu(Q)$ for every $Q \in \mathbb{D}_{Q_0} \setminus \{Q_0\}$, with $Q^* \supset Q$ and $\ell(Q^*) = 2\ell(Q)$ (i.e., Q^* is the “dyadic parent” of Q). For every $0 < \varepsilon_0 \leq e^{-1}$, if $\emptyset \neq F \subset Q_0$ with $\mu(F) \leq \alpha \mu(Q_0)$ and $0 < \alpha \leq \varepsilon_0^2 / (2C_\mu^2)$ then F has a good ε_0 -cover with respect to μ of length $k_0 = k_0(\alpha, \varepsilon_0, C_\mu) \in \mathbb{N}$, $k_0 \geq 2$, which satisfies $k_0 \approx \frac{\log \alpha^{-1}}{\log \varepsilon_0^{-1}}$. In particular, if $\mu(F) = 0$, then F has a good ε_0 -cover of arbitrary length.

We would like to mention that in the case $\mu(F) = 0$ this result gives an ε_0 -cover of arbitrary length. Our goal is to show that in such an scenario one can iterate the construction and construct an ε_0 -cover of infinite length:

Lemma A.5 *Let $E \subset \mathbb{R}^{n+1}$ be an n -dimensional ADR set and fix $Q_0 \in \mathbb{D}(E)$. Let μ be a regular Borel measure on Q_0 and assume that it is dyadically doubling on Q_0 . For every $0 < \varepsilon_0 \leq e^{-1}$, if $\emptyset \neq F \subset Q_0$ with $\mu(F) = 0$, then F has a good ε_0 -cover of length ∞ .*

Proof We are going to iterate Lemma A.4. Given $0 < \varepsilon_0 \leq e^{-1}$ and $0 < \alpha < \varepsilon_0^2/(2C_\mu^2)$, let $k_0 = k_0(\alpha, \varepsilon_0, C_\mu) \in \mathbb{N}$, $k_0 \geq 2$, be the value from Lemma A.4 so that $k_0 \approx \frac{\log \alpha^{-1}}{\log \varepsilon_0^{-1}}$. Let $F \subset Q_0$ with $\mu(F) = 0$. Using that $\mu(F) < \alpha\mu(Q_0)$, Lemma A.4 gives $\{\mathcal{O}_\ell^1\}_{\ell=1}^{k_0}$, a good ε_0 -cover of length k_0 of F with $\mathcal{F}_\ell^1 \subset \mathbb{D}_{Q_0}$ the associated families of pairwise disjoint cubes. This is the first generation in the construction.

To obtain the second generation take an arbitrary $Q \in \mathcal{F}_{k_0}^1$ and note that $\mu(F \cap Q) = 0 < \alpha\mu(Q)$. We apply again Lemma A.4 in Q to $F \cap Q$ (which is not empty by Remark A.2) and obtain a good ε_0 -cover $\{\tilde{\mathcal{O}}_\ell^2(Q)\}_{\ell=1}^{k_0}$ of $F \cap Q$ with associated families of pairwise disjoint cubes $\tilde{\mathcal{F}}_\ell^2(Q) \subset \mathbb{D}_Q$, $1 \leq \ell \leq k_0$, and so that

$$F \cap Q \subset \tilde{\mathcal{O}}_{k_0}^2(Q) \subset \tilde{\mathcal{O}}_{k_0-1}^2(Q) \subset \cdots \subset \tilde{\mathcal{O}}_2^2(Q) \subset \tilde{\mathcal{O}}_1^2(Q) \subset Q \in \mathcal{F}_{k_0}^1$$

Write $\tilde{\mathcal{O}}_\ell^2 := \bigcup_{Q \in \mathcal{F}_{k_0}^1} \tilde{\mathcal{O}}_\ell^2(Q)$ and $\tilde{\mathcal{F}}_\ell^2 = \bigcup_{Q \in \mathcal{F}_{k_0}^1} \tilde{\mathcal{F}}_\ell^2(Q)$ for $1 \leq \ell \leq k_0$. Since for each $Q \in \mathcal{F}_{k_0}^1$ and for each $1 \leq \ell \leq k_0$ the family $\tilde{\mathcal{F}}_\ell^2(Q) \subset \mathbb{D}_Q$ is pairwise disjoint, and the family $\mathcal{F}_{k_0}^1$ is also pairwise disjoint we easily conclude that $\tilde{\mathcal{F}}_\ell^2$ is a pairwise disjoint family. Besides,

$$\begin{aligned} F &= F \cap \mathcal{O}_{k_0}^1 = \bigcup_{Q \in \mathcal{F}_{k_0}^1} F \cap Q \subset \tilde{\mathcal{O}}_{k_0}^2 \subset \tilde{\mathcal{O}}_{k_0-1}^2 \subset \cdots \subset \tilde{\mathcal{O}}_2^2 \subset \tilde{\mathcal{O}}_1^2 \\ &\subset \bigcup_{Q \in \mathcal{F}_{k_0}^1} Q = \mathcal{O}_{k_0}^1 \subset \mathcal{O}_{k_0-1}^1 \subset \cdots \subset \mathcal{O}_1^1 \subset Q_0. \end{aligned} \quad (\text{A.6})$$

Set $\mathcal{O}_\ell^2 := \mathcal{O}_\ell^1$ for $1 \leq \ell \leq k_0$ and $\mathcal{O}_\ell^2 := \tilde{\mathcal{O}}_{\ell-k_0+1}^2$ for $k_0 + 1 \leq \ell \leq 2(k_0 - 1) + 1$. Write $\mathcal{F}_\ell^2 \subset \mathbb{D}_{Q_0}$ for the associated families of pairwise disjoint dyadic cubes for $1 \leq \ell \leq 2(k_0 - 1) + 1$. Our goal is to show that $\{\mathcal{O}_\ell^2\}_{\ell=1}^{2(k_0-1)+1}$ is a good ε_0 -cover of F whose length is $2(k_0 - 1) + 1$. By (A.6) and the previous construction, (a) and (b) in Definition A.1 clearly hold. We then need to verify (c). With this goal in mind we note that since $\{\mathcal{O}_\ell^1\}_{\ell=1}^{k_0}$ is a good ε_0 -cover, we obtain

$$\mu(\mathcal{O}_\ell^2 \cap Q) = \mu(\mathcal{O}_\ell^1 \cap Q) \leq \varepsilon_0 \mu(Q), \quad \forall Q \in \mathcal{F}_{\ell-1}^2 = \mathcal{F}_{\ell-1}^1, \quad 2 \leq \ell \leq k_0.$$

Also, if $Q \in \mathcal{F}_{k_0}^2 = \mathcal{F}_{k_0}^1$ then (A.3) in Remark A.2 applied to $\tilde{\mathcal{O}}_2^2(Q)$ gives

$$\mu\left(\mathcal{O}_{k_0+1}^2 \cap Q\right) = \mu\left(\tilde{\mathcal{O}}_2^2 \cap Q\right) = \mu\left(\tilde{\mathcal{O}}_2^2(Q)\right) \leq \varepsilon_0 \mu(Q).$$

On the other hand, let $k_0 + 2 \leq \ell \leq 2(k_0 - 1) + 1$ and let $Q \in \mathcal{F}_{\ell-1}^2 = \tilde{\mathcal{F}}_{\ell-k_0}^2$. By construction, there exists $Q' \in \mathcal{F}_{k_0}^1$ so that $Q \in \tilde{\mathcal{F}}_{\ell-k_0}^2(Q') \subset \mathbb{D}_{Q'}$. Then,

$$\mu\left(\mathcal{O}_\ell^2 \cap Q\right) = \mu\left(\tilde{\mathcal{O}}_{\ell-k_0+1}^2 \cap Q\right) = \mu\left(\tilde{\mathcal{O}}_{\ell-k_0+1}^2(Q') \cap Q\right) \leq \varepsilon_0 \mu(Q),$$

where we have used that $\{\tilde{\mathcal{O}}_\ell^2(Q')\}_{\ell=1}^{k_0}$ is a good ε_0 -cover. All these show (c) and as result $\{\mathcal{O}_\ell^2\}_{\ell=1}^{2(k_0-1)+1}$ is a good ε_0 -cover of F whose length is $2(k_0 - 1) + 1$.

The third generation is obtained in the very same way, we take $Q \in \mathcal{F}_{2(k_0-1)+1}^2$ and note that $\mu(F \cap Q) = 0 < \alpha \mu(Q)$. We apply again Lemma A.4 in Q to $F \cap Q$ and obtain $\{\tilde{\mathcal{O}}_\ell^3(Q)\}_{\ell=1}^{k_0}$, a good ε_0 -cover of $F \cap Q$ (which is not empty by Remark A.2) with $\tilde{\mathcal{F}}_\ell^3(Q) \subset \mathbb{D}_Q$ the associated families of pairwise disjoint cubes. We set $\tilde{\mathcal{O}}_\ell^3 := \bigcup_{Q \in \mathcal{F}_{2(k_0-1)+1}^2} \tilde{\mathcal{O}}_\ell^3(Q)$ and $\tilde{\mathcal{F}}_\ell^3 = \bigcup_{Q \in \mathcal{F}_{2(k_0-1)+1}^2} \tilde{\mathcal{F}}_\ell^3(Q)$ for $1 \leq \ell \leq k_0$. Define $\mathcal{O}_\ell^3 := \mathcal{O}_\ell^2$ for $1 \leq \ell \leq 2(k_0 - 1) + 1 = 2k_0 - 1$ and $\mathcal{O}_\ell^3 := \tilde{\mathcal{O}}_{\ell-2(k_0-1)}^3$ for $2k_0 \leq \ell \leq 3(k_0 - 1) + 1$. Write $\mathcal{F}_\ell^3 \subset \mathbb{D}_{Q_0}$ for the associated families of pairwise disjoint dyadic cubes for $1 \leq \ell \leq 3(k_0 - 1) + 1$. The same argument allows us to show that $\{\mathcal{O}_\ell^3\}_{\ell=1}^{3(k_0-1)+1}$ is a good ε_0 -cover of F whose length is $3(k_0 - 1) + 1$.

If we iterate this construction in N steps we will have constructed $\{\mathcal{O}_\ell^N\}_{\ell=1}^{N(k_0-1)+1}$, a good ε_0 -cover of F whose length is $N(k_0 - 1) + 1$. We observe that such iteration procedure works because $\mu(F) = 0$, hence $\mu(F \cap Q) = 0$ for every $Q \in \mathbb{D}_{Q_0}$ and also because $F \cap Q \neq \emptyset$ for every Q in each of the families that define the good ε_0 -cover (see Remark A.2). Since $k_0 \geq 2$ and we can continue with this iteration infinitely many times we eventually obtain an infinite good ε_0 -cover. $\square$

To continue we need to introduce some notation and some auxiliary result from [12]. Given $\eta = 2^{-k_*} < 1$ small enough to be chosen momentarily and given $Q \in \mathbb{D}(\partial\Omega)$ we define $Q^{(\eta)} \in \mathbb{D}_Q$ to be the unique dyadic cube such that $x_Q \in Q^{(\eta)}$ and $\ell(Q^{(\eta)}) = \eta \ell(Q)$.

Lemma A.7 [12, Lemma 3.24] *There exist $0 < \eta = 2^{-k_*} \ll 1$ and $\varepsilon_0 \ll 1$ small enough, and $c_0 \in (0, 1/2)$ (depending only on dimension, the 1-sided CAD constants and the ellipticity of L) with the following significance. Suppose that $\emptyset \neq F \subset Q_0 \in \mathbb{D}(\partial\Omega)$ is a Borel set and that $\{\mathcal{O}_\ell\}_{\ell=1}^k$ is a good ε_0 -cover of F with respect to $\omega_L^{X_{Q_0}}$ of length $k \in \mathbb{N}$, with associated pairwise disjoint families $\{\mathcal{F}_\ell\}_{1 \leq \ell \leq k} \subset \mathbb{D}_{Q_0}$. Define, $\mathcal{O}_\ell^{(\eta)} = \bigcup_{Q \in \mathcal{F}_\ell} Q^{(\eta)}$, for each $1 \leq \ell \leq k$, and consider the Borel set $S_k := \bigcup_{\ell=2}^k (\mathcal{O}_{\ell-1}^{(\eta)} \setminus \mathcal{O}_\ell)$. For each $y \in F$ and $1 \leq \ell \leq k$, let $Q^\ell(y) \in \mathcal{F}_\ell$ be the unique dyadic cube containing y , and let $P^\ell(y) \in \mathbb{D}_{Q^\ell(y)}$ be the unique dyadic cube*

containing y with $\ell(P^\ell(y)) = \eta\ell(Q^\ell(y))$. Then $u_k(X) := \omega_L^X(S_k)$, $X \in \Omega$, satisfies

$$\left| u_k \left(X_{(Q^\ell(y))^{(\eta)}} \right) - u_k \left(X_{(P^\ell(y))^{(\eta)}} \right) \right| \geq c_0, \quad 1 \leq \ell \leq k-1. \quad (\text{A.8})$$

We are now ready to prove Lemma 3.30:

Proof of Lemma 3.30 Fix $Q_0 \in \mathbb{D}$ and a Borel set $\emptyset \neq F \subset Q_0$ with $\omega_L^{X_{Q_0}}(F) = 0$. Let $\eta = 2^{-k_*}$ and ε_0 be small enough, and c_0 from Lemma A.7. From Lemma 2.19 and Harnack's inequality we have that $\omega_L^{X_{Q_0}}$ is Borel regular dyadically doubling measure on Q_0 . Applying Lemma A.5 with $\mu = \omega_L^{X_{Q_0}}$, one can find $\{\mathcal{O}_\ell\}_{\ell=1}^\infty$, a good ε_0 -cover of F of length ∞ . In particular, for every $N \in \mathbb{N}$, $\{\mathcal{O}_\ell\}_{\ell=1}^N$ is a good ε_0 -cover of F of length N . As such we can invoke Lemma A.7 to obtain $S_N := \bigcup_{\ell=2}^N (\mathcal{O}_{\ell-1}^{(\eta)} \setminus \mathcal{O}_\ell)$ so that with the notation introduced in that result the L -solution $u_N(X) := \omega_L^X(S_N)$, $X \in \Omega$, satisfies

$$\left| u_N \left(X_{(Q^\ell(y))^{(\eta)}} \right) - u_N \left(X_{(P^\ell(y))^{(\eta)}} \right) \right| \geq c_0, \quad \forall y \in F, \quad 1 \leq \ell \leq N-1. \quad (4.9)$$

Define next $S := \bigcup_{\ell=2}^\infty (\mathcal{O}_{\ell-1}^{(\eta)} \setminus \mathcal{O}_\ell)$ and $u(X) := \omega_L^X(S)$, $X \in \Omega$. By the monotone convergence theorem $u_N(X) \rightarrow u(X)$ as $N \rightarrow \infty$ for every $X \in \Omega$. Thus, (4.9) readily gives

$$\left| u \left(X_{(Q^\ell(y))^{(\eta)}} \right) - u \left(X_{(P^\ell(y))^{(\eta)}} \right) \right| \geq c_0, \quad \forall y \in F, \quad \ell \geq 1. \quad (4.10)$$

This and the argument in [12, p. 7919] imply

$$\left(\iint_{U_{Q^\ell(y), \eta^3}^{\vartheta_0}} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY \right)^{\frac{1}{2}} \gtrsim_\eta c_0, \quad \forall \ell \geq 1. \quad (4.11)$$

Thus, for every $N \geq 1$ and every $y \in F$

$$\begin{aligned} Nc_0^2 &\lesssim_\eta \sum_{\ell=1}^N \iint_{U_{Q^\ell(y), \eta^3}^{\vartheta_0}} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY \lesssim_\eta \iint_{\bigcup_{\ell=1}^N U_{Q^\ell(y), \eta^3}^{\vartheta_0}} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY \\ &\leq \iint_{\bigcup_{y \in Q \in \mathbb{D}_{Q_0}} U_{Q, \eta^3}^{\vartheta_0}} |\nabla u(Y)|^2 \delta(Y)^{1-n} dY = S_{Q_0}^{\vartheta_0, \eta} u(y)^2, \end{aligned}$$

where we have used that the family $\{U_{Q, \eta^3}^{\vartheta_0}\}_{Q \in \mathbb{D}_{Q_0}}$ has bounded overlap albeit with a constant that depends on η . Letting $N \rightarrow \infty$ we conclude that $S_{Q_0}^{\vartheta_0, \eta} u(y) = \infty$ for every $y \in F$ and the proof is complete. $\square$

References

1. Akman, M., Azzam, J., Mourougolou, M.: Absolute continuity of harmonic measure for domains with lower regular boundaries. *Adv. Math.* **345**, 1206–1252 (2019)
2. Akman, M., Badger, M., Hofmann, S., Martell, J.M.: Rectifiability and elliptic measures on 1-sided NTA domains with Ahlfors-David regular boundaries. *Trans. Am. Math. Soc.* **369**, 5711–5745 (2017)
3. Akman, M., Bortz, S., Hofmann, S., Martell, J.M.: Rectifiability, interior approximation and harmonic measure. *Ark. Mat.* **57**, 1–22 (2019)
4. Akman, M., Hofmann, S., Martell, J.M., Toro, T.: Square function and non-tangential maximal function estimates for elliptic operators in 1-sided NTA domains satisfying the capacity density condition, [arXiv:2103.10046](https://arxiv.org/abs/2103.10046)
5. Akman, M., Hofmann, S., Martell, J.M., Toro, T.: Perturbation of elliptic operators in 1-sided NTA domains satisfying the capacity density condition, [arXiv:1901.08261v3](https://arxiv.org/abs/1901.08261v3)
6. Azzam, J., Hofmann, S., Martell, J.M., Mayboroda, S., Mourougolou, M., Tolsa, X., Volberg, A.: Rectifiability of harmonic measure. *Geom. Funct. Anal.* **26**, 703–728 (2016)
7. Azzam, J., Hofmann, S., Martell, J.M., Nyström, K., Toro, T.: A new characterization of chord-arc domains. *J. Eur. Math. Soc.* **19**, 967–981 (2017)
8. Azzam, J., Mourougolou, M., Tolsa, X.: Singular sets for harmonic measure on locally flat domains with locally finite surface measure. *Int. Math. Res. Not. IMRN* **2017**(12), 3751–3773 (2017)
9. Badger, M.: Null sets of harmonic measure on NTA domains: Lipschitz approximation revisited. *Math. Z.* **270**, 241–262 (2012)
10. Bishop, C.J., Jones, P.W.: Harmonic measure and arclength. *Ann. Math.* **2**(132), 511–547 (1990)
11. Caverio, J., Hofmann, S., Martell, J.M.: Perturbations of elliptic operators in 1-sided chord-arc domains. Part I: Small and large perturbation for symmetric operators. *Trans. Am. Math. Soc.* **371**, 2797–2835 (2019)
12. Caverio, J., Hofmann, S., Martell, J.M., Toro, T.: Perturbations of elliptic operators in 1-sided chord-arc domains. Part II: Non-symmetric operators and Carleson measure estimates. *Trans. Am. Math. Soc.* **373**(11), 7901–7935 (2020)
13. Christ, M.: A Tb theorem with remarks on analytic capacity and the Cauchy integral. *Colloq. Math.* **60**(61), 601–628 (1990)
14. Dahlberg, B.: Estimates of harmonic measure. *Arch. Ration. Mech. Anal.* **65**, 275–288 (1977)
15. Dahlberg, B.: Weighted norm inequalities for the Lusin area integral and the nontangential maximal functions for functions harmonic in a Lipschitz domain. *Studia Math.* **67**, 297–314 (1980)
16. Dahlberg, B.: On the absolute continuity of elliptic measures. *Am. J. Math.* **108**, 1119–1138 (1986)
17. David, G., Jerison, D.: Lipschitz approximation to hypersurfaces, harmonic measure, and singular integrals. *Indiana Univ. Math. J.* **39**, 831–845 (1990)
18. David, G., Semmes, S.: Singular integrals and rectifiable sets in $\mathbb{R}^n$, Au-delà des graphes lipschitziens. *Asterisque* **193** (1991)
19. David, G., Semmes, S.: Analysis of and on Uniformly Rectifiable Sets, Mathematical Monographs and Surveys, vol. 38. AMS (1993)
20. Fabes, E.B., Jerison, D.S., Kenig, C.E.: Necessary and sufficient conditions for absolute continuity of elliptic-harmonic measure. *Ann. Math.* **2**(119), 121–141 (1984)
21. Fefferman, R.: A criterion for the absolute continuity of the harmonic measure associated with an elliptic operator. *J. Am. Math. Soc.* **2**(1), 127–135 (1989)
22. Fefferman, R., Kenig, C.E., Pipher, J.: The theory of weights and the Dirichlet problem for elliptic equations. *Ann. Math.* **2**(134), 65–124 (1991)
23. Hofmann, S., Martell, J.M.: Uniform rectifiability and harmonic measure I: Uniform rectifiability implies Poisson kernels in L^p . *Ann. Sci. Éc. Norm. Supér.* **47**, 577–654 (2014)
24. Hofmann, S., Martell, J.M., Mayboroda, S., Toro, T., Zhao, Z.: Uniform rectifiability and elliptic operators satisfying a Carleson measure condition. *Geom. Funct. Anal.* **31**(2), 325–401 (2021)
25. Hofmann, S., Martell, J.M., Toro, T.: A_∞ implies NTA for a class of variable coefficient elliptic operators. *J. Differ. Equ.* **263**, 6147–6188 (2017)
26. Hofmann, S., Martell, J.M., Toro, T.: Elliptic Operators on Non-smooth Domains (Book in preparation)
27. Hofmann, S., Martell, J.M., Uriarte-Tuero, I.: Uniform rectifiability and harmonic measure, II: Poisson kernels in L^p imply uniform rectifiability. *Duke Math. J.* **163**, 1601–1654 (2014)
28. Hofmann, S., Mitrea, D., Mitrea, M., Morris, A.J.: L^p -square function estimates on spaces of homogeneous type and on uniformly rectifiable sets. *Mem. Am. Math. Soc.* **245**, 1159 (2017)

29. Jerison, D.S., Kenig, C.E.: The Dirichlet problem in nonsmooth domains. *Ann. Math.* **2**(113), 367–382 (1981)
30. Jerison, D.S., Kenig, C.E.: Boundary behavior of harmonic functions in nontangentially accessible domains. *Adv. Math.* **46**, 80–147 (1982)
31. Kenig, C.E., Kirchheim, B., Pipher, J., Toro, T.: Square functions and the A_∞ property of elliptic measures. *J. Geom. Anal.* **26**, 2383–2410 (2016)
32. Kenig, C.E., Koch, H., Pipher, J., Toro, T.: A new approach to absolute continuity of elliptic measure, with applications to non-symmetric equations. *Adv. Math.* **153**(2), 231–298 (2000)
33. Kenig, C.E., Pipher, J.: The Dirichlet problem for elliptic equations with drift terms. *Publ. Mat.* **45**, 199–217 (2001)
34. Kenig, C.E., Toro, T.: Harmonic measure on locally flat domains. *Duke Math. J.* **87**, 509–551 (1997)
35. Kaufman, R., Wu, J.: Distortion of the boundary under conformal mapping. *Michigan Math. J.* **29**, 267–280 (1982)
36. Lavrentiev, M.: Boundary problems in the theory of univalent functions (Russian). *Mat. Sb.* **43**, 815–844 (1936)
37. McMillan, J.E.: Boundary behavior of a conformal mapping. *Acta Math.* **123**, 43–67 (1973)
38. Milakis, E., Pipher, J., Toro, T.: Harmonic analysis on chord arc domains. *J. Geom. Anal.* **23**, 2091–2157 (2013)
39. Milakis, E., Pipher, J., Toro, T.: Perturbations of elliptic operators in chord arc domains. In: *Harmonic Analysis and Partial Differential Equations, Contemp. Math.*, vol. 612, pp. 143–161. Amer. Math. Soc., Providence, RI (2014)
40. Øksendal, B.: Sets of harmonic measure zero. In: *Aspects of Contemporary Complex Analysis, Proc. NATO Adv. Study Inst., Univ. Durham, Durham, 1979*, Academic Press, London-New York, pp. 469–473 (1980)
41. Riesz, F., Riesz, M.: Über die randwerte einer analitischen funktion, *Compte Rendues du Quatrième Congrès des Mathématiciens Scandinaves*, Stockholm 1916. Almqvists and Wilksels, Upsala (1920)
42. Semmes, S.: Analysis vs. geometry on a class of rectifiable hypersurfaces in $\mathbb{R}^n$. *Indiana Univ. Math. J.* **39**, 1005–1035 (1990)
43. Toro, T., Zhao, Z.: Boundary rectifiability and elliptic operators with $W^{1,1}$ coefficients. *Adv. Calc. Var.* **14**(1), 37–62 (2021)
44. Wu, J.: On singularity of harmonic measure in space. *Pac. J. Math.* **121**, 485–496 (1986)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Mingming Cao¹ · José María Martell¹ · Andrea Olivo²

✉ José María Martell
chema.martell@icmat.es

Mingming Cao
mingming.cao@icmat.es

Andrea Olivo
aolivo@dm.uba.ar

¹ Instituto de Ciencias Matemáticas CSIC-UAM-UC3M-UCM, Consejo Superior de Investigaciones Científicas, C/ Nicolás Cabrera, 13-15, 28049 Madrid, Spain

² Departamento de Matemática, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires and IMAS-CONICET, Pabellón I, C1428EGA Ciudad de Buenos Aires, Argentina