On the Catlin Multitype of Sums of Squares Domains

Nicholas Aidoo¹

Received: 23 June 2021 / Accepted: 12 February 2022 / Published online: 26 February 2022
© The Author(s) 2022

Abstract

For a sum of squares domain of finite D'Angelo 1-type at the origin, we show that the polynomial model obtained from the computation of the Catlin multitype at the origin of such a domain is likewise a sum of squares domain. We also prove, under the same finite type assumption that the multitype is an invariant of the ideal of holomorphic functions defining the domain. Both results are proven using Martin Kolář's algorithm for the computation of the multitype introduced in Kolář (Int Math Res Not (IMRN) 18:3530–3548, 2010). Given a sum of squares domain, we rewrite the Kolář algorithm in terms of ideals of holomorphic functions and also introduce an approach that explicitly constructs the homogeneous polynomial transformations used in the algorithm.

Keywords Sum of squares domains · Catlin multitype · Finite type domains in $\mathbb{C}^n$ · The Kolář algorithm

Mathematics Subject Classification Primary 32F18 · Secondary 32T27; 32T25

1 Introduction

Domains defined by sums of squares of holomorphic functions constitute a very important class in the field of several complex variables as they connect in a very natural way complex analysis with algebraic geometry. This class of domains was introduced by Kohn in his Acta paper [12] under the term *special domains*. Kohn's novel introduction of the technique of subelliptic multiplier ideals to address the local regularity problem paved the way for significant discoveries in several complex variables. In [5,6], D'Angelo studied the local geometry of real hypersurfaces by assigning to every point on the hypersurface an associated family of ideals of holomorphic functions and explor-

✉ Nicholas Aidoo
nicholas.aidoo@ntnu.no

¹ Department of Mathematical Sciences, Norwegian University of Science and Technology, Trondheim, Norway

ing various invariants in commutative algebra and algebraic geometry. He established a close connection between the geometry of sums of squares domains and complex algebraic geometry. Further work by Siu in [16] and [17] on sums of squares domains introduced new approaches for generating multipliers for general systems of partial differential equations. Owing to his initial work on sums of squares domains, Siu gave an extension of the special domain approach to real analytic and smooth cases. Kim and Zaitsev in [9] proposed a new class of geometric invariants called *the jet-vanishing orders* and used them to establish a new selection algorithm in the Kohn's construction of subelliptic multipliers of sums of squares domains in dimension 3. Also, in a recent paper by the same authors [10], they provide a solution to the effectiveness problem in Kohn's algorithm for generating multipliers for domains including those defined by sums of squares of holomorphic functions in all dimensions. Other important results pertaining to sums of squares domains can be found in [3,4,7,8,14].

A key motivation for this paper is to introduce some important tools and techniques necessary for the study of the multitype level set stratification of sums of squares domains. We will focus here on the multitype computations for such domains. Our main tool is an algorithm devised by Kolář in [13] for the computation of the Catlin multitype when it has finite entries. In order to ensure this condition is satisfied, we will assume finite D'Angelo 1-type throughout since the latter bounds from above the last entry of the multitype; see [2].

Owing to the connection between the properties of the associated family of ideals of holomorphic functions and the geometry of sums of squares domains, a natural question is whether or not the multitype could be computed from the corresponding ideals of holomorphic functions. We provide an answer to this question by showing that the multitype of a sum of squares domain can be computed from the related ideal of holomorphic functions via the restatement of the Kolář algorithm at the level of ideals. Besides the fact that working with ideals aligns better with complex algebraic geometry, this restatement also reduces significantly the amount of work involved in computing the Catlin multitype for sums of squares domains.

A *sum of squares domain* $\Omega \subset \mathbb{C}^{n+1}$ is one of which boundary-defining function $r(z)$ is given by

$$r(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_{n+1})|^2, \quad (1.1)$$

where $f_j(z_1, \dots, z_{n+1})$ for all j , $1 \leq j \leq N$ are holomorphic functions vanishing at the origin in $\mathbb{C}^{n+1}$. We shall denote by $M \subset \mathbb{C}^{n+1}$ the hypersurface defined by $\{z \in \mathbb{C}^{n+1} \mid r(z) = 0\}$.

The *model hypersurface* associated to M at the origin is defined as follows:

$$M_H = \{z \in \mathbb{C}^{n+1} : H(z, \bar{z}) = 0\}, \quad (1.2)$$

the zero locus of the homogeneous polynomial $H(z, \bar{z})$ consisting of all monomials from the Taylor expansion of the defining function that have weight 1 with respect to the multitype weight. We refer to $H(z, \bar{z})$ as the *model polynomial*.

Our first main result concerns the form of the model polynomial $H(z, \bar{z})$:

Theorem 1.1 *Let $M \subset \mathbb{C}^{n+1}$ be a hypersurface of which defining function is given as follows:*

$$r(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_{n+1})|^2,$$

where $f_1, \dots, f_N$ are holomorphic functions defined on a neighborhood of the origin and assume that the D'Angelo 1-type of M at the origin is finite. Then the model associated to M is also a sum of squares domain.

In other words, the model polynomial

$$H(z, \bar{z}) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |h_j(z_1, \dots, z_n)|^2, \quad (1.3)$$

where h_j is a polynomial consisting of all terms from the Taylor expansion of f_j of weight $1/2$ with respect to the multitype at the origin, $j = 1, \dots, N$. Note that the corresponding model hypersurface $M_H = \{z \in \mathbb{C}^{n+1} : H(z, \bar{z}) = 0\}$ is a decoupled sum of squares domain since variable z_{n+1} has weight 1, so no h_j can depend on it. By Catlin's results in [2], M_H has the same multitype at the origin as the original domain. In fact, with respect to multitype computations, every model polynomial is decoupled.

Our second main result answers in the affirmative a question posed by D. Zaitsev during a discussion with the author:

Theorem 1.2 *Let $0 \in M \subset \mathbb{C}^{n+1}$ be a hypersurface of which defining function is given as follows:*

$$r(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_{n+1})|^2,$$

where $f_1, \dots, f_N$ are holomorphic functions defined on a neighborhood of the origin and assume that the D'Angelo 1-type of M is finite at the origin.

Then the Catlin multitype is an invariant of the ideal of holomorphic functions defining the domain.

Computing the multitype via the Kolář algorithm requires one to use weighted homogeneous polynomial transformations at each step in order to generate an intermediate weight and a partial model polynomial that depends on a minimal number of variables. How these homogeneous polynomial transformations are obtained was not explicitly described in [13], however. In this paper, we characterize these weighted homogeneous polynomial transformations and explicitly construct them by relating them to elementary row and column operations on the Levi matrix associated to the domain. More specifically, we devise an algorithm for the construction of these

homogeneous polynomial transformations, the row reduction algorithm. We call a polynomial change of variables at step j of the Kolář algorithm *allowable with respect to variable* z_k if it eliminates the occurrence of variable z_k from the leading polynomial P_j and reduces the number of variables in P_j by at least one. For the correspondence between polynomial changes of variables and elementary row and column operations on the Levi matrix associated to the domain, we introduce a variant of the notion of dependence compared to the standard one in linear algebra as follows: Let A_{P_j} be the Levi matrix of the leading polynomial P_j at step j of the Kolář algorithm. For a given k , denote by R_k and C_k the k th row and the k th column of the matrix A_{P_j} respectively. Let $\mathcal{R}$ be the set of all rows of the matrix A_{P_j} . We call a row R_k of $\mathcal{R}$ *dependent* if it satisfies the condition:

$$R_k = \sum_{\substack{l=1 \\ l \neq k}}^n \alpha_l R_l,$$

where $\alpha_l \in \mathbb{C}[z]$, $\alpha_l \neq 0$ for at least one l . In other words, the row R_k is dependent if it can be written as a linear combination of the other rows with polynomial coefficients. The set $\mathcal{R}$ is said to be *dependent* if at least one of the rows is dependent.

Since the matrix A_{P_j} is Hermitian, a similar definition holds for the k th column C_k of A_{P_j} namely

$$C_k = \sum_{\substack{l=1 \\ l \neq k}}^n \bar{\alpha}_l C_l.$$

Hence, we prove the following:

Proposition 1.3 *Assume that the D'Angelo 1-type of the hypersurface M in $\mathbb{C}^{n+1}$ at 0 is finite. At step j of the Kolář algorithm for the computation of the multitype at 0, let Λ_j be an intermediate weight, P_j the leading polynomial, and let Q_j be the leftover polynomial, which consists of monomials of weighted degree greater than 1 with respect to Λ_j . Let k be given, $1 \leq k \leq n$. There exists an allowable polynomial transformation on P_j with respect to the variable z_k if and only if the k th row of A_{P_j} is dependent.*

Using this characterization of the polynomial transformations and the restatement of the Kolář algorithm in terms of ideals of holomorphic functions, the row reduction algorithm connects nicely the notion of simplifying the Jacobian module associated to a sum of squares domain with elementary row operations on the complex Jacobian matrix of the same domain. By employing this algorithm at every step of the Kolář algorithm for the computation of the multitype, we are able to construct the weighted homogeneous polynomial transformations needed in the Kolář algorithm.

This paper is structured in the following manner: Sect. 2 describes some notation and provides definitions that are pertinent to our discussions in subsequent sections. Section 3 defines the Catlin multitype and provides a thorough description of the Kolář

algorithm as introduced in [13]. Section 4 presents a key lemma for the characterization of the multitype entries of the sum of squares domain. Specifically, we establish the fact that each multitype entry can be realized by the modulus square of some monomial of vanishing order at least one in the multivariate ring of polynomials over $\mathbb{C}$. Section 5 contains the proofs of Theorems 1.1 and 1.2. Section 6 presents a modified version of the Kolář algorithm in terms of ideals of holomorphic polynomials. In the same section, a corollary to Theorem 1.1 is stated and proven. Finally, Sect. 7 contains the proof of Proposition 1.3 and also provides a detailed description of the row reduction algorithm for the construction of the weighted homogeneous polynomials required in the Kolář algorithm.

This article constitutes part of the author's Ph.D thesis at Trinity College Dublin under the supervision of Andreea Nicoara.

2 Definitions and Notation

We present some definitions that we use in the article following the setup of Kolář in [13]. Let M be a hypersurface in $\mathbb{C}^{n+1}$ and $p \in M$ be a Levi-degenerate point. We will assume that p is a point of finite D'Angelo 1-type. Let (z, w) be local holomorphic coordinates centered at the point p , where $w = u + iv$ is the complex non-tangential variable and the complex tangential variables are in the n -tuple $z = (z_1, \dots, z_n)$ with $z_k = x_k + iy_k$. Throughout this paper, we will compute and define weights by considering only the complex tangential variables $z_1, \dots, z_n$ as in [13].

Definition 2.1 A weight $\Lambda = (\mu_1, \dots, \mu_n)$ is an n -tuple of rational numbers with $0 \leq \mu_j \leq \frac{1}{2}$ satisfying

- i. $\mu_j \geq \mu_{j+1}$ for $1 \leq j \leq n-1$;
- ii. For each t , either $\mu_t = 0$ or there exists a sequence of nonnegative integers $a_1, \dots, a_t$ satisfying $a_t > 0$ such that

$$\sum_{j=1}^t a_j \mu_j = 1.$$

Let Λ be a weight. If $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multiindex, then we define the weighted length of α by

$$|\alpha|_\Lambda = \sum_{j=1}^n \alpha_j \mu_j.$$

Also if $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\hat{\alpha} = (\hat{\alpha}_1, \dots, \hat{\alpha}_n)$ are multiindices, then the weighted length of the pair $(\alpha, \hat{\alpha})$ is defined by

$$|(\alpha, \hat{\alpha})|_\Lambda = \sum_{j=1}^n (\alpha_j + \hat{\alpha}_j) \mu_j.$$

Definition 2.2 A monomial $A_{\alpha\hat{\alpha}l}z^\alpha\bar{z}^{\hat{\alpha}}u^l$ is said to be of weighted degree κ if

$$\kappa := l + |(\alpha, \hat{\alpha})|_\Lambda.$$

Similarly, we define the weighted order of the differential operator $D^\alpha \bar{D}^{\hat{\alpha}} D^l$ to equal to $\kappa := l + |(\alpha, \hat{\alpha})|_\Lambda$, where

$$D^\alpha = \frac{\partial^{|\alpha|}}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}, \quad \bar{D}^{\hat{\alpha}} = \frac{\partial^{|\hat{\alpha}|}}{\partial \bar{z}_1^{\hat{\alpha}_1} \dots \partial \bar{z}_n^{\hat{\alpha}_n}}, \quad \text{and} \quad D^l = \frac{\partial^l}{\partial u^l}.$$

A polynomial $P(z, \bar{z}, u)$ is said to be Λ -homogeneous of weighted degree κ if it is a sum of monomials of weighted degree κ .

We shall set the variable w as well as the variables u and v to have a weight of one.

Definition 2.3 A weight $\Lambda = (\mu_1, \dots, \mu_n)$ is said to be *distinguished* if there exist local holomorphic coordinates (z, w) mapping p to the origin such that the boundary-defining equation for M in the new coordinates is of the form:

$$v = P(z, \bar{z}) + o_\Lambda(1), \quad (2.1)$$

where $P(z, \bar{z})$ is a Λ -homogeneous polynomial of weighted degree 1 without pluri-harmonic terms and $o_\Lambda(1)$ denotes a smooth function of which derivatives of weighted order less than or equal to 1 vanish at zero.

We order the weights lexicographically. This means that for the pair of weights $\Lambda_1 = (\mu_1, \dots, \mu_n)$ and $\Lambda_2 = (\mu'_1, \dots, \mu'_n)$, $\Lambda_1 > \Lambda_2$ if for some t , $\mu_j = \mu'_j$ for $j < t$ and $\mu_t > \mu'_t$.

Definition 2.4 Let $\Lambda = (\lambda_1, \dots, \lambda_n)$ be a weight and

$$\tilde{w} = w + g(z_1, \dots, z_n, w) \text{ and } \tilde{z}_j = z_j + f_j(z_1, \dots, z_n, w),$$

for $1 \leq j \leq n$, be a holomorphic change of variables. We say that this transformation is

- i. Λ -homogeneous if f_j is a Λ -homogeneous polynomial of weighted degree λ_j and g is a Λ -homogeneous polynomial of weighted degree 1,
- ii. Λ -superhomogeneous if f_j has a Taylor expansion consisting of monomials that have weighted degree $\geq \lambda_j$ and g consists of terms of weighted degree ≥ 1 ,
- iii. Λ -subhomogeneous if the Taylor expansion of f_j consists of terms of weighted degree $\leq \lambda_j$ and g consists of weighted degree ≤ 1 .

J. J. Kohn introduced the notion of type of a point on a pseudoconvex hypersurface in $\mathbb{C}^2$ in [11]. In [1], Thomas Bloom and Ian Graham generalized Kohn's notion to $\mathbb{C}^n$ and gave a geometric characterization of type of points on real hypersurfaces in $\mathbb{C}^n$. There is the definition given by Bloom and Graham: Let $\mathcal{N}$ be a real C^∞ hypersurface

defined in an open subset $U \subset \mathbb{C}^n$ with defining function r . Let $\mathcal{L}_k$ for $k \geq 0$ an integer, be the module, over $C^\infty(U)$, of vector fields generated by the tangential holomorphic vector fields to $\mathcal{N}$, their conjugates, and commutators of order less than or equal to k of such vector fields.

Definition 2.5 A point $p \in \mathcal{N}$ is of type m if $\langle \partial r(p), F(p) \rangle = 0$ for all $F \in \mathcal{L}_{m-1}$ while $\langle \partial r(p), F(p) \rangle \neq 0$ for some $F \in \mathcal{L}_m$.

Here, we denote the contractions between a cotangent vector and a tangent vector by $\langle \cdot, \cdot \rangle$. We shall refer to the type at a point p as defined above as the *Bloom-Graham type*.

3 Catlin Multitype and the Kolář Algorithm

In this section, we describe the Catlin multitype and the Kolář algorithm for the computation of the multitype at the origin in [13]. The notion of weights, distinguished weights, and the multitype was introduced by Catlin in [2]. We will follow the notation and definitions as given by Kolář in [13] and further describe some of the tools that M. Kolář introduced:

Definition 3.1 Let M be a hypersurface in $\mathbb{C}^{n+1}$, and let $p \in M$. Let $\Lambda^* = (\mu_1, \dots, \mu_n)$ be the greatest lower bound with respect to the lexicographic ordering of all the distinguished weights at p . The *multitype* $\mathcal{M}$ at p is defined to be the n -tuple $(m_1, \dots, m_n)$, where $m_j = \infty$ if $\mu_j = 0$ and $m_j = \frac{1}{\mu_j}$ if $\mu_j \neq 0$. We call the multitype $\mathcal{M}$ at p finite, if the last entry $m_n < \infty$.

The next theorem from [2] clarifies the relationship between the multitype and the D'Angelo type for a pseudoconvex domain:

Theorem 3.1 (Catlin). *Let $\Omega \subset \mathbb{C}^{n+1}$ be a pseudoconvex domain smooth boundary. Let $p_0 \in b\Omega$ be a boundary point. If $\mathcal{M}(p_0) = (m_1, \dots, m_n)$ is the multitype at p_0 , then for each $q = 1, \dots, n$, $m_{n+1-q} \leq \Delta_q(b\Omega, p_0)$, where $\Delta_q(b\Omega, p_0)$ is the D'Angelo q -type at p_0 .*

For the purposes of this paper, we shall assume finite D'Angelo 1-type at any point $p \in M$, since this assumption by Theorem 3.1 ensures that all the entries of the multitype are finite, which is the exact setting in which the Kolář algorithm works.

For a weight Λ , we say the local coordinates on M at p are Λ -adapted if M is described locally to have the form in (2.1), where P is Λ -homogeneous. We shall refer to Λ^* -adapted coordinates as the multitype coordinates given such that P is Λ^* -homogeneous.

Let γ_j , $j = 1, \dots, c$, be the length of the j th constant piece of the multitype weight given such that c is the number of distinct entries in the multitype. Let $\sum_{i=1}^j \gamma_i = k_j$, then we have

$$\mu_1 = \dots = \mu_{k_1} > \mu_{k_1+1} = \dots = \mu_{k_2} > \dots = \mu_{k_{c-1}} > \dots = \mu_{k_{c-1}+1} = \dots = \mu_n,$$

where $n = k_c$. We define a monotone sequence of weights $\Lambda_1, \dots, \Lambda_c$ which are ordered lexicographically as follows. Λ_1 is a constant n -tuple $(\mu_1, \dots, \mu_1)$ and $\Lambda_c = \Lambda^*$ is the multitype weight. We then define the weight $\Lambda_j = (\lambda_1^j, \dots, \lambda_n^j)$ for $1 < j < c$, by $\lambda_i^j = \mu_i$ for $i \leq k_{j-1}$ and $\lambda_i^j = \mu_{k_{j-1}+1}$ for $i > k_{j-1}$. Note that this construction yields a finite sequence of weights even if Λ^* has some zero entries.

Definition 3.2 Fix Λ^* -adapted local coordinates. The *leading polynomial* P is defined as follows:

$$P(z, \bar{z}) = \sum_{|(\alpha, \hat{\alpha})|_{\Lambda^*} = 1} C_{\alpha, \hat{\alpha}} z^\alpha \bar{z}^{\hat{\alpha}}. \quad (3.1)$$

The polynomial defined in (3.1) is exactly the polynomial that only retains the terms of weight 1. Put differently, it is a Λ^* -homogeneous polynomial of weighted degree 1 with no pluriharmonic terms, where Λ^* is the multitype weight. Following Kolář in [13], we will also denote by *leading polynomial* the polynomial consisting of all terms of weight 1 with respect to each intermediate weight Λ_j in the Kolář algorithm.

Theorem 3.2 (Kolář) *A biholomorphic transformation takes Λ^* -adapted coordinates into Λ^* -adapted coordinates if and only if this transformation is Λ^* -superhomogeneous.*

We will now describe the Kolář algorithm for the computation of the multitype and apply this theorem under the assumption that all entries of the multitype are finite.

The Kolář Algorithm We start by considering local holomorphic coordinates in which the leading polynomial in the variables z and $\bar{z}$ contains no pluriharmonic terms. The degree of the lowest order monomial in this polynomial is then equal to the Bloom-Graham type of M at p as defined in [1]. This gives the first multitype component m_1 ; see [2]. By our assumption, $1 < m_1 < +\infty$. Let $m_1 = \frac{1}{\mu_1}$ and set $\Lambda_1 = (\mu_1, \dots, \mu_1)$. We then consider all Λ_1 -homogeneous transformations and choose one that will make the leading polynomial P_1 to be independent of the largest number of variables. We denote this number by d_1 . So for such coordinates, we get the defining function of M to be of the form:

$$v = P_1(z_1, \dots, z_{n-d_1}, \bar{z}_1, \dots, \bar{z}_{n-d_1}) + Q_1(z, \bar{z}) + o(u),$$

where P_1 is Λ_1 -homogeneous of weighted degree 1 and Q_1 is $o_{\Lambda_1}(1)$. We use the result that for any weight Λ which is lexicographically smaller than Λ_1 , Λ -adapted coordinates are also Λ_1 -adapted coordinates and the fact that Λ_1 -homogeneous transformation is linear, to conclude that $\mu_1 = \dots = \mu_{n-d_1}$ and $\mu_{n-d_1+1} < \mu_1$. We define the following important tools:

$$\Theta_1 = \left\{ (\alpha, \hat{\alpha}) \mid C_{\alpha, \hat{\alpha}}^1 \neq 0 \text{ and } \sum_{i=1}^{n-d_1} (\alpha_i + \hat{\alpha}_i) \mu_i < 1 \right\},$$

and also

$$Q_1(z, \bar{z}) = \sum_{|(\alpha, \hat{\alpha})|_{\Lambda_1} > 1} C_{\alpha, \hat{\alpha}}^1 z^\alpha \bar{z}^{\hat{\alpha}}.$$

For every $(\gamma, \hat{\gamma}) \in \Theta_1$,

$$W_1(\gamma, \hat{\gamma}) = \frac{1 - \sum_{i=1}^{n-d_1} (\gamma_i + \hat{\gamma}_i) \mu_i}{\sum_{i=n-d_1+1}^n (\gamma_i + \hat{\gamma}_i) \mu_i}. \quad (3.2)$$

We define the next weight Λ_2 by letting

$$\lambda_j^2 = \max_{(\alpha, \hat{\alpha}) \in \Theta_1} W_1(\alpha, \hat{\alpha})$$

for $j > n - d_1$ and $\lambda_j^2 = \mu_1$ for $j \leq n - d_1$. We then complete the second by letting P_2 be the new leading polynomial corresponding to the weight Λ_2 . P_2 depends on more than $n - d_1$ variables.

We proceed by induction. At the j th step, for $j > 2$, using coordinates from the previous step, we consider all Λ_{j-1} -homogeneous transformations and choose one that makes the leading polynomial P_{j-1} to be independent of the largest number of variables. We fix such coordinates, and let d_{j-1} be the largest number of variables, which do not show up in P_{j-1} after this change of variables. By Theorem 3.2, the transformations taking Λ_{j-1} -adapted coordinates into Λ_{j-1} -adapted coordinates are always Λ_{j-1} -superhomogeneous. The number of multitype entries that are added at each step of the computation depends on the difference $(d_{j-2} - d_{j-1})$. Hence, we consider two cases in this step:

Case 1 Assume that $d_{j-2} > d_{j-1}$. Also recall that for any weight Λ that is smaller than Λ_{j-1} with respect to the lexicographic ordering, Λ -adapted coordinates are also Λ_{j-1} -adapted. This implies that we get $(d_{j-2} - d_{j-1})$ multitype entries

$$\mu_{n-d_{j-2}+1} = \cdots = \mu_{n-d_{j-1}} = \lambda_{n-d_{j-2}+1}^{j-1}$$

and let $\lambda_i^j = \mu_i$ for $i \leq n - d_{j-1} - 1$. To obtain λ_i^j for $j > n - d_{j-1} - 1$, we consider

$$v = P_{j-1}(z_1, \dots, z_{n-d_{j-1}}, \bar{z}_1, \dots, \bar{z}_{n-d_{j-1}}) + Q_{j-1}(z, \bar{z}) + o(u),$$

where Q_{j-1} is $o_{\Lambda_{j-1}}(1)$ and P_{j-1} is Λ_{j-1} -homogeneous of weighted degree 1. We define Q_{j-1} , Θ_{j-1} , and W_{j-1} in a similar way as in step two. Thus,

$$\Theta_{j-1} = \left\{ (\alpha, \hat{\alpha}) \mid C_{\alpha, \hat{\alpha}}^{j-1} \neq 0 \text{ and } \sum_{i=1}^{n-d_{j-1}} (\alpha_i + \hat{\alpha}_i) \mu_i < 1 \right\},$$

and also

$$Q_{j-1}(z, \bar{z}) = \sum_{|(\alpha, \hat{\alpha})|_{\Lambda_{j-1}} > 1} C_{\alpha, \hat{\alpha}}^{j-1} z^{\alpha} \bar{z}^{\hat{\alpha}}.$$

For every $(\gamma, \hat{\gamma}) \in \Theta_{j-1}$,

$$W_{j-1}(\gamma, \hat{\gamma}) = \frac{1 - \sum_{i=1}^{n-d_{j-1}} (\gamma_i + \hat{\gamma}_i) \mu_i}{\sum_{i=n-d_{j-1}+1}^n (\gamma_i + \hat{\gamma}_i)}. \quad (3.3)$$

So for the remaining multitype entries of Λ_j , we let

$$\lambda_i^j = \max_{(\alpha, \hat{\alpha}) \in \Theta_{j-1}} W_{j-1}(\alpha, \hat{\alpha}),$$

for $i > n - d_{j-1}$.

Case 2 Assume that $d_{j-1} = d_{j-2}$. There are zero multitype entries computed in this case and so we only determine λ_i^j for $j > n - d_{j-1}$ using (3.3). This completes the j th step of the computation.

The process terminates after a finite number of steps to give all the entries of the multitype weight Λ^* . It is clear that case 1 advances the process. We just need to show that the number of times case 2 occurs where no multitype entries are determined can only happen finitely many times. We claim case 2 can take place at most $\lceil \frac{1}{\mu_n} \rceil^{n-d_{j-1}+1}$ times, where $\lceil \frac{1}{\mu_n} \rceil$ is the ceiling for the rational number $\frac{1}{\mu_n}$. Indeed, it comes down to the number of different values that (3.3) can have. The upper bound for the numerator is given by $\lceil \frac{1}{\mu_n} \rceil^{n-d_{j-1}}$ as the μ_i entries are decreasing, whereas the upper bound for the denominator is given by $\lceil \frac{1}{\mu_n} \rceil$.

Example 1 Let $M \subset \mathbb{C}^4$ be a real hypersurface near the origin given by $r = 0$ with

$$r = 2\operatorname{Re}(z_4) + |z_1 - z_2 + z_3^2|^2 + |z_1^2 - z_2^2|^2.$$

Using the Kolář algorithm, we proceed as follows: The Bloom-Graham type is 2, which implies that $\mu_1 = \frac{1}{2}$ and $\Lambda_1 = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$. Thus, $P_1 = |z_1 - z_2|^2$. We consider all Λ_1 -homogeneous transformation and choose

$$\tilde{z}_1 = z_1 - z_2, \quad \text{and} \quad \tilde{z}_j = z_j,$$

for $j = 2, 3, 4$, to obtain a leading polynomial P_1 independent of the variables z_2 and z_3 . We write r in the new variables and ignore $\sim$ when no confusion arises. Then $P_1 = |z_1|^2$ with $d_1 = 2$. The leftover polynomial is

$$Q_1 = |z_3^2|^2 + 2\operatorname{Re}(z_1 \bar{z}_3^2) + |z_1^2|^2 + 4\operatorname{Re}(z_1^2 \bar{z}_1 \bar{z}_2) + 4|z_1 z_2|^2.$$

$\Lambda_2 = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$, where the maximum number $\max(W_1) = \frac{1}{4}$. We consider all Λ_2 -homogeneous transformations and choose

$$\tilde{z}_1 = z_1 + z_3^2 \text{ and } \tilde{z}_j = z_j$$

for $j = 2, 3, 4$, which makes the leading polynomial independent of the variables z_2 and z_3 . In the new variables, $P_2 = |z_1|^2$ with $d_2 = 2$ and

$$\begin{aligned} Q_2 = & |z_1^2|^2 + 4|z_1 z_3^2|^2 + 4\operatorname{Re}(z_1^2 \bar{z}_1 \bar{z}_3^2) + |z_3^4|^2 + 2\operatorname{Re}(z_1^2 \bar{z}_3^4) \\ & + 4\operatorname{Re}(z_1 z_3^2 \bar{z}_3^4) + 4|z_1 z_2|^2 + 4\operatorname{Re}(z_1^2 \bar{z}_1 \bar{z}_2) \\ & + 8\operatorname{Re}(z_1 z_3^2 \bar{z}_1 \bar{z}_2) + 4\operatorname{Re}(z_3^4 \bar{z}_1 \bar{z}_2) + 4|z_2 z_3^2|^2 - 4\operatorname{Re}(z_1^2 \bar{z}_2 \bar{z}_3^2) - 8\operatorname{Re}(z_1 z_3^2 \bar{z}_2 \bar{z}_3^2) \\ & - 4\operatorname{Re}(z_3^4 \bar{z}_2 \bar{z}_3^2) - 8\operatorname{Re}(z_1 z_2 \bar{z}_2 \bar{z}_3^2). \end{aligned}$$

$\Lambda_3 = (\frac{1}{2}, \frac{1}{6}, \frac{1}{6})$ with $\max(W_2) = \frac{1}{6}$. Here, no Λ_3 -homogeneous transformation can make P_3 to be independent of any variables and so $d_3 = 0$. Hence,

$$P_3 = |z_1|^2 + 4|z_2 z_3^2|^2.$$

4 Squares of Monomials and the Multitype Entries

We will show in this section that the set of monomials that give the maximum W -value at each step of the Kolář algorithm always consists of both squares of moduli of monomials and cross terms. Since the entries in the multitype depend on the maximum W -values, it will suffice to establish that for each entry of the multitype, there is always a square that gives the corresponding multitype entry.

As hinted in the introduction, we rely on the Kolář algorithm for the computation of the multitype in [13] to develop tools to study the stratification by multitype level sets of the boundary of a domain given by a sum of squares of holomorphic functions. We seek a tool that will enable us to effectively interpret our results both geometrically and algebraically. As such, we resort to reducing the problem to the study of the ideal of the holomorphic functions in the sum. For the transition to be effective, we need to establish that a natural modification of the Kolář algorithm still holds at the level of ideals.

The ensuing lemma gives the foundational result we need in order to transition from the sums of squares case to the case of ideals of holomorphic functions in $\mathbb{C}^n$. We use the result from this lemma to prove Theorems 1.1 and 1.2.

Lemma 4.1 *Let f and g be monomials with nonzero coefficients from the Taylor expansion of h , where h is a generator from a sum of squares domain in $\mathbb{C}^{n+1}$. Let P_t for $t \geq 1$ be the leading polynomial at step t of the Kolář algorithm, and let W_t be the number computed at the $(t + 1)$ th step.*

- A. *If $W_t(|f|^2) = W_t(|g|^2)$, then $W_t(f\bar{g}) = W_t(|f|^2) = W_t(|g|^2)$.*
- B. *If $W_t(|f|^2) < W_t(|g|^2)$, then $W_t(|f|^2) < W_t(f\bar{g}) < W_t(|g|^2)$.*

C. If $W_t(|f|^2)$ cannot be computed, then

- i. $W_t(f\bar{g}) \leq W_t(|g|^2)$ for any monomial g for which both $W_t(|g|^2)$ and $W_t(f\bar{g})$ can be computed.
- ii. $W_t(f\bar{g})$ cannot be computed for any monomial g for which $W_t(|g|^2)$ cannot be computed.

D. If f is such that $|f|^2$ is in the leading polynomial P_t , then

- i. For any monomial g for which $W_t(|g|^2)$ can be computed, $W_t(f\bar{g}) = W_t(|g|^2)$.
- ii. For any monomial g for which $W_t(|g|^2)$ cannot be computed, $W_t(f\bar{g})$ cannot be computed as well.

Here $W_t(f) := W_t(\alpha, \hat{\alpha})$ where $(\alpha, \hat{\alpha})$ is the pair of multiindices corresponding to the monomial f .

Remark 4.1.1 We shall say the quantity $W_t(f)$ “cannot be computed” if the pair of multiindices $(\alpha, \hat{\alpha})$ corresponding to the monomial f is not an element of Θ_t . In other words, $W_t(f)$ cannot be computed if the numerator of the fraction giving $W_t(f)$ is not positive; see (4.1) below.

Remark 4.1.2 We note here that $W_t(f\bar{g})$ and $W_t(\bar{f}g)$ corresponding to the cross terms $f\bar{g}$ and $\bar{f}g$, respectively, are equal.

Proof Let $z_1, \dots, z_c$ be the variables in the leading polynomial P_t , and let the weight $\Lambda_t = (\mu_1, \dots, \mu_c, \mu_{c+1}, \dots, \mu_n)$. We begin by recalling that

$$W_t(\alpha, \hat{\alpha}) = \frac{1 - \sum_{i=1}^c (\alpha_i + \hat{\alpha}_i) \mu_i}{\sum_{i=c+1}^n (\alpha_i + \hat{\alpha}_i)}, \quad (4.1)$$

where $(\alpha, \hat{\alpha}) = (\alpha_1, \dots, \alpha_n, \hat{\alpha}_1, \dots, \hat{\alpha}_n)$ is the multiindex of the monomial of which W_t is being computed.

Let Γ_1 be the set of all nonzero monomials that consist of only variables not in P_t , let Γ_2 be the set of all nonzero monomials which consist of variables both in P_t as well as variables not in P_t , and let Γ_3 be the set of all nonzero monomials which consist of only variables in P_t . If $W_t(|f|^2)$ can be computed, then $f \in \Gamma_1$ or $f \in \Gamma_2$ only. We will now prove f cannot belong to Γ_3 . We assume the opposite, namely that $f \in \Gamma_3$ and that $W_t(|f|^2)$ can be computed. The monomial $|f|^2$ has weight ≥ 1 with respect to Λ_t , and since $f \in \Gamma_3$, it follows that the numerator of $W_t(|f|^2)$ is ≤ 0 . Therefore, $W_t(|f|^2)$ cannot be computed, which is a contradiction. Without loss of generality, we now specify to $f \in \Gamma_1$ or $f \in \Gamma_2$ for all parts of the lemma pertaining to the case when $W_t(|f|^2)$ can be computed (and similarly for g).

Since $f \in \Gamma_1$ or $f \in \Gamma_2$, we can write $f = f_1 f_2$ where f_1 and f_2 are monomials satisfying $f_1 \in \Gamma_3$ and $f_2 \in \Gamma_1$. Let $f_1 = z_1^{\alpha_1} \dots z_c^{\alpha_c}$, where $z_1, \dots, z_c$ are the variables in the leading polynomial P_t , and $f_2 = C_f z_{c+1}^{\beta_{c+1}} \dots z_n^{\beta_n}$ for $C_f \in \mathbb{C}$. If $W_t(|f|^2)$ can be computed, by our definition of f , the multiindices α and β corresponding to monomials f_1 and f_2 , respectively, must satisfy $|\alpha| \geq 0$ and $|\beta| > 0$. If $|\alpha| = 0$, then $f \in \Gamma_1$, and if $|\alpha| > 0$, then $f \in \Gamma_2$. Here $|\alpha| = \alpha_1 + \dots + \alpha_c$ as the rest of the entries are

zero, and $|\beta| = \beta_{c+1} + \cdots + \beta_n$ for the same reason. Now, $|f|^2 = f_1 \bar{f}_1 f_2 \bar{f}_2$ with $|\alpha| = |\hat{\alpha}|$ and $|\beta| = |\hat{\beta}|$. Hence,

$$W_t(|f|^2) = \frac{1 - \sum_{i=1}^c (\alpha_i + \hat{\alpha}_i) \mu_i}{|\beta| + |\hat{\beta}|} = \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i}{|\beta|}. \quad (4.2)$$

Similarly, let $g = g_1 g_2$, where $g_1 \in \Gamma_3$ and $g_2 \in \Gamma_1$. Let γ and τ be the multiindices corresponding to monomials g_1 and g_2 , respectively. Here $g_1 = z_1^{\gamma_1} \cdots z_c^{\gamma_c}$ and $g_2 = C_g z_{c+1}^{\tau_{c+1}} \cdots z_n^{\tau_n}$ for $C_g \in \mathbb{C}$. By a similar computation as carried out above for f ,

$$W_t(|g|^2) = \frac{\frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\tau|}. \quad (4.3)$$

A. Suppose that $W_t(|f|^2) = W_t(|g|^2)$ and consider

$$\begin{aligned} W_t(f\bar{g}) &= \frac{1 - \sum_{i=1}^c \alpha_i \mu_i - \sum_{i=1}^c \hat{\gamma}_i \mu_i}{|\beta| + |\tau|} \\ &= \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i + \frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\beta| + |\tau|} \\ &= \frac{|\beta| W_t(|f|^2) + |\tau| W_t(|g|^2)}{|\beta| + |\tau|} \text{ from (4.2) and (4.3) by cross multiplication.} \\ &= W_t(|f|^2) = W_t(|g|^2) \text{ by our hypothesis.} \end{aligned} \quad (4.4)$$

B. Suppose that $W_t(|f|^2) < W_t(|g|^2)$. Then from (A.) we get that

$$\begin{aligned} W_t(f\bar{g}) &= \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i + \frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\beta| + |\tau|} \\ &= \frac{|\beta| W_t(|f|^2) + |\tau| W_t(|g|^2)}{|\beta| + |\tau|} \text{ from (4.2) and (4.3)} \\ &< \frac{|\beta| W_t(|g|^2) + |\tau| W_t(|g|^2)}{|\beta| + |\tau|} \text{ since } W_t(|f|^2) < W_t(|g|^2) \\ &= W_t(|g|^2). \end{aligned} \quad (4.5)$$

Again

$$W_t(f\bar{g}) = \frac{|\beta| W_t(|f|^2) + |\tau| W_t(|g|^2)}{|\beta| + |\tau|} > W_t(|f|^2) \quad (4.6)$$

since $W_t(|f|^2) < W_t(|g|^2)$. Thus, from (4.5) and (4.6), we obtain

$$W_t(|f|^2) < W_t(f\bar{g}) < W_t(|g|^2).$$

C. i. Suppose that $W_t(|f|^2)$ cannot be computed. Then clearly $f \notin \Gamma_1$. We know that $f = f_1 f_2$ and that $|\alpha| > 0$ and $|\beta| \geq 0$. If $W_t(|f|^2)$ cannot be computed, then we get that $\sum_{i=1}^c (\alpha_i + \hat{\alpha}_i) \mu_i \geq 1$. Thus, $\sum_{i=1}^c \alpha_i \mu_i \geq 1/2$ since $\alpha_i = \hat{\alpha}_i$ for all i , $1 \leq i \leq c$.

Let $g = g_1 g_2$, where $g_1 \in \Gamma_3$ and $g_2 \in \Gamma_1$. $W_t(|g|^2)$ and $W_t(f \bar{g})$ can be computed, and so g cannot belong to Γ_3 , which implies the multiindex corresponding to g_2 must satisfy $|\tau| > 0$. Therefore,

$$\begin{aligned} W_t(f \bar{g}) &= \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i + \frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\beta| + |\tau|} \\ &= \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i + |\tau| W_t(|g|^2)}{|\beta| + |\tau|} \quad \text{from (4.3)} \\ &\leq \frac{|\tau|}{|\beta| + |\tau|} W_t(|g|^2) \quad \text{since } \frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i \leq 0 \\ &\leq W_t(|g|^2) \quad \text{since } \frac{|\tau|}{|\beta| + |\tau|} \leq 1. \end{aligned} \quad (4.7)$$

ii. From (C.i.) we know that $f = f_1 f_2 \notin \Gamma_1$ and also that $\sum_{i=1}^c \alpha_i \mu_i \geq 1/2$ since $\alpha_i = \hat{\alpha}_i$ for all i , $1 \leq i \leq c$. Similarly, for $g = g_1 g_2$ it means that $g \notin \Gamma_1$ and that $\sum_{i=1}^c \gamma_i \mu_i \geq 1/2$. Thus, $\sum_{i=1}^c (\alpha_i + \gamma_i) \mu_i \geq 1$, and so the pair of multiindices corresponding to the cross term $f \bar{g}$ is not in the set Θ_t . Hence, the number $W_t(f \bar{g})$ cannot be computed.

D. i. Suppose that f is such that $|f|^2$ is in the leading polynomial P_t . Then $f \in \Gamma_3$, which implies that $f = f_1 f_2$ with $|\beta| = 0$, that is, $f = C f_1$ for C a nonzero constant. Let $g = g_1 g_2$ be any monomial such that $W_t(|g|^2)$ can be computed. Clearly, $g \notin \Gamma_3$, and so $|\tau| > 0$ whereas $|\gamma| \geq 0$. We know from (4.3) that

$$W_t(|g|^2) = \frac{\frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\tau|}.$$

Given that $|f|^2$ is a term in P_t , $\sum_{i=1}^c (\alpha_i + \hat{\alpha}_i) \mu_i = 1$ since P_t is a Λ_t -homogeneous polynomial of weighted degree 1. Thus, $\sum_{i=1}^c \alpha_i \mu_i = 1/2$ since $\alpha_i = \hat{\alpha}_i$ for all i , $1 \leq i \leq c$. Now

$$\begin{aligned} W_t(f \bar{g}) &= \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i + \frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\beta| + |\tau|} \\ &= \frac{\frac{1}{2} - \sum_{i=1}^c \alpha_i \mu_i + \frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\tau|} \quad \text{since } |\beta| = 0 \\ &= \frac{\frac{1}{2} - \sum_{i=1}^c \gamma_i \mu_i}{|\tau|} \quad \text{since } \sum_{i=1}^c \alpha_i \mu_i = \frac{1}{2} \\ &= W_t(|g|^2). \end{aligned} \quad (4.8)$$

- ii. Let $g = g_1 g_2$ be any monomial such that $W_t(|g|^2)$ cannot be computed. From (D.i.) we know that $\sum_{i=1}^c \alpha_i \mu_i = 1/2$ and from (C.ii.) we know that $\sum_{i=1}^c \gamma_i \mu_i \geq 1/2$. Thus,

$$\sum_{i=1}^c (\alpha_i + \gamma_i) \mu_i = \sum_{i=1}^c (\alpha_i + \gamma_i) \mu_i = \frac{1}{2} + \sum_{i=1}^c \gamma_i \mu_i \geq 1.$$

Hence the pair of multiindices corresponding to the cross term $f \bar{g}$ does not belong to the set Θ_t , and so the number $W_t(f \bar{g})$ cannot be computed.

□

Remark 4.1.3 The terms that are added to the leading polynomial P_t at step t of the Kolář algorithm are the terms needed to compute the multitype. Clearly, the maximum W_t at every step is always assumed by a square of a monomial. All the cross terms that are added to the leading polynomial could be eliminated without changing the multitype. Most importantly, the multitype of M at the origin will not change if:

- All the cross terms are eliminated from the expansion of $|h|^2$, where h is a generator of a sum of squares domain and
- All the squares that do not appear in the final leading polynomial are added to our data.

5 Proofs of the Main Results

We give the proofs of the two main theorems in this section. As hinted in the introduction, we shall use the result from Lemma 4.1 to prove Theorems 1.1 and 1.2. By our convention in this paper, we assume without loss of generality that the domains we work with are decoupled as mentioned in the introduction. We start by proving a more general result than Theorem 1.1.

Theorem 5.1 *Let $0 \in M \subset \mathbb{C}^{n+1}$ be a hypersurface of which defining function is given as*

$$r(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2,$$

where $f_1, \dots, f_N$ are holomorphic functions near the origin and assume that the D'Angelo 1-type of M at the origin is finite. Then the leading polynomial $P_t(z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n)$ obtained at step t of the Kolář algorithm, for $t \geq 1$, is a sum of squares of holomorphic polynomials. In particular, the final leading polynomial $P(z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n)$ corresponding to the multitype weight is likewise a sum of squares.

Proof We order the generators f_k by vanishing order. We truncate each generator f_k up to order $\beta = \lceil \Delta_1(M, 0) \rceil$, the ceiling of the D'Angelo 1-type of M at the origin,

and denote it by f_k^β . Since a sum of squares domain is pseudoconvex, by Theorem 3.1, no terms of higher order than β ought to come into the computation of the multitype. We order the terms in the truncated generator f_k^β by vanishing order and also use the reverse lexicographic ordering to reorder the monomials with the same combined degree. Now let $f_{k,i} = C_{k,i} z_1^{\alpha_{k,i}^1} \dots z_n^{\alpha_{k,i}^n}$ be the i th monomial in the generator f_k^β after ordering by vanishing order. Let the number of distinct combined degrees in the Taylor expansion f_k^β be κ_k , and let $\eta_{k,j}$ be the number of nonzero monomials with the same combined degree $v_{k,j}$ in f_k^β . Thus,

$$|f_k^\beta|^2 = |f_{k,1} + \dots + f_{k,\eta_k}|^2, \text{ where } \eta_k = \sum_{j=1}^{\kappa_k} \eta_{k,j}.$$

Here η_k is the total number of monomials with nonzero coefficients in the power series expansion of the generator f_k up to order β . In the expansion of $|f_k^\beta|^2$, we have two types of terms: squares $|f_{k,i}|^2$ and cross terms $2\operatorname{Re}(f_{k,j} \bar{f}_{k,i})$. For simplicity sake, for each monomial $f_{k,i}$ in the generator f_k^β , we write the terms from the expansion of $|f_k^\beta|^2$ into an expression of the form:

$$|f_{k,i}|^2 + \sum_{j=1}^{i-1} 2\operatorname{Re}(f_{k,j} \bar{f}_{k,i}), \quad (5.1)$$

for $i = 1, 2, \dots, \eta_k$.

Define $P_{t,k}$ for $t \geq 1$ to be the sum in the leading polynomial P_t at step t consisting of terms from the expansion of $|f_k^\beta|^2$. We could have $P_{t,k} = 0$ for some k , $1 \leq k \leq N$, and for some $t \geq 1$ if there are no terms from the expansion of $|f_k^\beta|^2$ in the leading polynomial P_t after the t th step. Thus, $P_t = \sum_{k=1}^N P_{t,k}$. In order to show that the final leading polynomial P_t is a sum of squares, it will suffice to show that each $P_{t,k}$ obtained at each step of the Kolář algorithm is a sum of squares. Note that trivially 0 is a sum of squares.

The Bloom–Graham type is $2v_1$, where $v_1 := \min_{1 \leq k \leq N} \{v_{k,1}\} = v_{1,1}$ since the monomials from the expansion of each $|f_k^\beta|^2$ as well as the generators f_k are ordered by vanishing order. Thus, the leading polynomial P_1 consists of all terms of combined degree $2v_1$. Clearly, $P_{1,1} \neq 0$, but $P_{1,k} = 0$ for every k such that $v_{k,1} > v_1$. Let k be such that $P_{1,k} \neq 0$, and denote by $m_k \geq 1$ the number of monomials from the Taylor expansion of f_k^β that have combined degree v_1 . By writing the terms from the expansion of $|f_k^\beta|^2$ into the form given in (5.1), it is easy to see that the first m_k squares $|f_{k,i}|^2$ for $i = 1, \dots, m_k$ as well as the $\binom{m_k}{2}$ cross terms $2\operatorname{Re}(f_{k,j} \bar{f}_{k,i})$ for $j < i$ all have combined degree $2v_1$. Thus,

$$P_{1,k} = |f_{k,1} + \dots + f_{k,m_k}|^2, \quad (5.2)$$

which is obviously a sum of squares. If $d_1 = 0$, then we are done and P_1 becomes the leading polynomial with multitype weight Λ_1 . On the other hand, if $d_1 = n - c$ for $c < n$, then we proceed to the next step.

In the second step, we assume without loss of generality that the W_1 value of at least one square in Q_1 can be computed. Then the maximum W_1 value exists, which further implies that some terms from the expansion of $|f_k^\beta|^2$ for some k will be added to the leading polynomial P_1 in order to obtain P_2 . $P_{2,k}$ might be 0 for some k if no terms from the expansion of $|f_k^\beta|^2$ end up in P_2 after this step. Obviously, the interesting case is when $P_{2,k} \neq 0$. Consider k such that $P_{2,k} \neq 0$. Suppose that u_k squares from the expansion of $|f_k^\beta|^2$ give the maximum W_1 value, and let $|f_{k,c_l}|^2$ for $l = 1, \dots, u_k$ be these squares. Here c_l for $l = 1, \dots, u_k$ is some positive integer between 1 and η_k , and $c_l < c_{l+1}$ for $l = 1, \dots, u_k - 1$. The argument now splits into two cases:

Case 1 $P_{1,k} = 0$. By Lemma 4.1, part A, we get exactly $\binom{u_k}{2}$ cross terms $2\operatorname{Re}(f_{k,c_e} \bar{f}_{k,c_l})$, which give the maximal value for W_1 and $c_e < c_l$ for $1 \leq e, l \leq u_k$. Combining the u_k squares with the $\binom{u_k}{2}$ cross terms gives

$$P_{2,k} = |f_{k,c_1} + \dots + f_{k,c_{u_k}}|^2. \quad (5.3)$$

Case 2 $P_{1,k} \neq 0$. Then from Lemma 4.1, part D(i), we know that for each square $|f_{k,c_l}|^2$ there are exactly m_k cross terms $2\operatorname{Re}(f_{k,j} \bar{f}_{k,c_l})$ for all $j < c_l$ and $j = 1, \dots, m_k$ as well as $l = 1, \dots, u_k$ that give the maximal value for W_1 . We also know from the first case that there are $\binom{u_k}{2}$ cross terms $2\operatorname{Re}(f_{k,c_e} \bar{f}_{k,c_l})$ that give the maximal value for W_1 , and so we obtain the result that

$$P_{2,k} = |f_{k,1} + \dots + f_{k,m_k} + f_{k,c_1} + \dots + f_{k,c_{u_k}}|^2. \quad (5.4)$$

In the $(t+1)$ th step, we begin by first assuming that the sum $P_{t,k}$ for some k is a sum of squares because if $P_{t,k} = 0$ the argument is identical to the one given in Case 1. Thus, let $P_{t,k}$ be given as $P_{t,k} = |f_{k,b_1} + \dots + f_{k,b_{v_k}}|^2$, where v_k is the total number of squares from the expansion of $|f_k^\beta|^2$ in P_t after step t with $b_j < b_{j+1}$ for $j = 1, \dots, v_k - 1$. Let's assume that the W_t value of at least one square in Q_t can be computed. This implies that the maximal value for W_t exists. Assume that s_k squares from the expansion of $|f_k^\beta|^2$ give the maximum W_t value, and let $|f_{k,a_l}|^2$ for $l = 1, \dots, s_k$ be these squares.

From Lemma 4.1, part D(i), we know that for each square $|f_{k,a_l}|^2$ there are exactly v_k cross terms $2\operatorname{Re}(f_{k,j} \bar{f}_{k,a_l})$ for all $j < a_l$ where $j = b_1, \dots, b_{v_k}$ and $l = 1, \dots, s_k$, which give the maximal value for W_t . By Lemma 4.1, part A, there are $\binom{s_k}{2}$ cross terms $2\operatorname{Re}(f_{k,a_e} \bar{f}_{k,a_l})$ which give the maximal value for W_t , and so we obtain the result that

$$P_{t+1,k} = |f_{k,b_1} + \dots + f_{k,b_{v_k}} + f_{k,a_1} + \dots + f_{k,a_{s_k}}|^2. \quad (5.5)$$

Hence, $P_{t+1,k}$ is a sum of squares. Since the leading polynomial P_t at each step is a sum of the $P_{t,k}$'s, the leading polynomial at each step and subsequently the final leading polynomial is a sum of squares.

Since every change of variables that is allowed in the Kolář algorithm sends each square to a square and keeps the weight of terms in the leading polynomial P_t the same, it is easy to see that the leading polynomial is still a sum of squares after any change of variables.

Our assumption of finite D'Angelo 1-type implies that the last entry of the multitype is bounded, and so all entries of the multitype weight will be finite. This means that the Kolář algorithm will definitely terminate after a finite number of steps, and so the above procedure can only occur finitely many times. We conclude, therefore that the final leading polynomial corresponding to the last weight in the procedure, the multitype weight, is always a sum of squares, too. $\square$

Proof of Theorem 1.1 We know that the final leading polynomial in the Kolář algorithm for computing the multitype at the origin is the model polynomial. Thus, from Theorem 5.1, we conclude that the model of a sum of squares is a sum of squares. $\square$

Before we proof Theorem 1.2 we shall consider the following lemma:

Lemma 5.2 *Let $M \subset \mathbb{C}^{n+1}$ with $0 \in M$ be a hypersurface of which defining function is given by*

$$r(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2,$$

where $f_1, \dots, f_N$ are holomorphic functions near the origin. Let $M' \subset \mathbb{C}^{n+1}$ be another hypersurface of which defining function is given as

$$u(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^{l-1} |f_j|^2 + |h_l f_l - \sum_{c \neq l} h_c f_c|^2 + \sum_{j=l+1}^N |f_j|^2,$$

for some fixed l , where h_j is a holomorphic function near the origin for every $j = 1, \dots, N$. Assume that the D'Angelo 1-type of M is finite at the origin. Then the multitype obtained by applying Kolář algorithm to both $r(z)$ and $u(z)$ is the same provided that $\langle f_1, \dots, f_N \rangle$ and

$$\left\langle f_1, \dots, f_{l-1}, h_l f_l - \sum_{c \neq l} h_c f_c, f_{l+1}, \dots, f_N \right\rangle$$

represent the same ideal in the ring $\mathcal{O}$.

Remark 5.2.1 Modifying only one generator at a time makes the bookkeeping in the computation of the multitype easier to follow.

Remark 5.2.2 Assume that there exist some l such that $1 \leq l \leq N$ and holomorphic functions near the origin $h_1, \dots, h_{l-1}, h_{l+1}, \dots, h_N$ such that $f_l = \sum_{c \neq l} h_c f_c$. It is clear that $\langle f_1, \dots, f_N \rangle$ and $\langle f_1, \dots, f_{l-1}, f_{l+1}, \dots, f_N \rangle$ represent the same ideal in $\mathcal{O}$. Therefore, applying Lemma 5.2 with $h_l \equiv 1$ shows that adding in the square $|f_l|^2$ or taking it away makes absolutely no difference as far as the multitype computation goes. This observation will be crucial in the proof of Theorem 1.2.

Proof We will show that the multitype obtained by applying the Kolář algorithm to both $r(z)$ and $u(z)$ is the same. Let $\beta = \lceil \Delta_1(M, 0) \rceil$ be the ceiling of the D'Angelo 1-type of M at the origin. We truncate each generator f_k as well as each holomorphic function h_k at the order β and denote them by f_k^β and h_k^β , respectively. Denote by $f_{k,i}$ the i th monomial from the Taylor expansion of f_k^β after ordering by vanishing order and reverse lexicographic order for the monomials with the same vanishing order.

By Lemma 4.1, we know that each entry of the multitype is realized by a square. If no square from the expansion of $|f_l^\beta|^2$ contributes to the entries of the multitype, then the multitype entries for both defining functions $r(z)$ and $u(z)$ are the same, and there is nothing to prove. We, thus, assume that there exists at least one square from the expansion of $|f_l^\beta|^2$ that contributes to the entries of the multitype and that no nonzero multiple of that square exists in any of the expansions of $|f_1^\beta|^2, \dots, |f_{l-1}^\beta|^2, |f_{l+1}^\beta|^2, \dots, |f_N^\beta|^2$.

Next, we claim that if $h_l(0) = 0$, then no square from the expansion of $|h_l^\beta f_l^\beta|^2$ can contribute to the entries of the multitype. Indeed, let $h_{l,i}$ be the i th monomial from the Taylor expansion of h_l^β after ordering by vanishing order and reverse lexicographic order for the monomials with the same vanishing order. For every monomial $f_{l,j}$ in f_l^β , the monomial $h_{l,i} f_{l,j}$ in $h_l^\beta f_l^\beta$ has greater combined degree than that of $f_{l,j}$ for every $i \geq 1$. As a result, $|h_{l,i} f_{l,j}|^2$ cannot give the Bloom-Graham type since the combined degree of $|f_{l,j}|^2$ is strictly less than the combined degree of $|h_{l,i} f_{l,j}|^2$. Furthermore, $W_t(|h_{l,i} f_{l,j}|^2)$ cannot be computed if $W_t(|f_{l,j}|^2)$ gives the maximum W_t -value. By the same argument, if $f_l = \sum_{c \neq l} g_c f_c$, for g_c with $c = 1, \dots, l-1, l+1, \dots, N$ holomorphic functions near the origin, then no square from the expansion of $|f_l^\beta|^2$ can contribute to the entries of the multitype unless it is the square of the nonzero constant term of some g_c multiplied by a monomial of f_c that in itself gives that same multitype entry. Since we assumed the contrary, f_l cannot be written in terms of the other generators. By our hypothesis, however, $\langle f_1, \dots, f_N \rangle$ and $\langle f_1, \dots, f_{l-1}, h_l f_l - \sum_{c \neq l} f_c h_c, f_{l+1}, \dots, f_N \rangle$ represent the same ideal in the ring $\mathcal{O}$. Putting these two facts together along with our assumption that there exists at least one square from the expansion of $|f_l^\beta|^2$ that contributes to the entries of the multitype and that no nonzero multiple of that square exists in any of the expansions of $|f_1^\beta|^2, \dots, |f_{l-1}^\beta|^2, |f_{l+1}^\beta|^2, \dots, |f_N^\beta|^2$, we conclude that $h_l(0) \neq 0$. Without loss of generality, assume $h_l \equiv 1$. We shall show that modifying the function f_l^β in the sum of squares by the sum $\sum_{c \neq l} f_c^\beta h_c^\beta$ does not alter the multitype. We further assume that the sum $\sum_{c \neq l} f_c^\beta h_c^\beta \neq 0$. By Lemma 4.1, it suffices to focus on how the omission

of some squares of monomials from the term $|f_l^\beta - \sum_{c \neq l} h_c^\beta f_c^\beta|^2$ in the defining function $u(z)$ affects our results. We now break our argument into two cases:

Case 1 Assume that there exists a monomial m in f_l^β of which square $|m|^2$ from the expansion of $|f_l^\beta|^2$ gives the Bloom-Graham type. We consider two subcases here:

- i. Assume that m is in the expression $f_l^\beta - \sum_{c \neq l} h_c^\beta f_c^\beta$. Then no monomial in the sum $\sum_{c \neq l} h_c^\beta f_c^\beta$ cancels out m . Clearly, the weights obtained at the first step of the Kolář algorithm are the same for both $r(z)$ and $u(z)$ since $|m|^2$ belongs to both defining functions.
- ii. Assume that m is not in the expression $f_l^\beta - \sum_{c \neq l} h_c^\beta f_c^\beta$. Hence, m gets canceled out in the expression $f_l^\beta - \sum_{c \neq l} h_c^\beta f_c^\beta$ and so does not appear in $u(z)$. Let ψ be the monomial in the sum $\sum_{c \neq l} h_c^\beta f_c^\beta$ that cancels out m , and write $\psi = h_{c,i} f_{c,j}$ for some $c \neq l$, where $h_{c,i}$ is some monomial in h_c^β and $f_{c,j}$ is some monomial in f_c^β . By our assumption, ψ equals m , and its square $|\psi|^2$ gives the Bloom-Graham type as well. The monomial $h_{c,i}$ cannot have vanishing order 1 or higher; otherwise, $f_{c,j}$ must have combined degree less than that of ψ , which contradicts the fact that $|\psi|^2$ gives the Bloom-Graham type. Thus, $h_{c,i} = h_{c,1} \in \mathbb{C}$ and $m = h_{c,1} f_{c,j}$. Hence, the square $|f_{c,j}|^2$ gives the Bloom-Graham type as well. Even though there is the cancellation in $u(z)$, the weight obtained at the first step having applied the Kolář algorithm to $r(z)$ and $u(z)$ is the same. More specifically, the squares $|f_{c,j}|^2$ and $|m|^2$ appear in the expansions of $|f_c^\beta|^2$ and $|f_l^\beta|^2$, respectively.

Case 2 Assume that there exists a monomial m in f_l^β of which square $|m|^2$ from the expansion of $|f_l^\beta|^2$ gives the maximum W_t -value at the $(t+1)$ th step for $t \geq 1$.

- i. Assume that m is in the expression $f_l^\beta - \sum_{c \neq l} h_c^\beta f_c^\beta$. Then no monomial in the sum $\sum_{c \neq l} h_c^\beta f_c^\beta$ cancels out m , and so the weights obtained at the $(t+1)$ th step are the same for both $r(z)$ and $u(z)$ since $|m|^2$ belongs to both defining functions.
- ii. Assume that m is not in the expression $f_l^\beta - \sum_{c \neq l} h_c^\beta f_c^\beta$. Then m gets canceled out by some monomial ψ in the sum $\sum_{c \neq l} h_c^\beta f_c^\beta$. Let $\psi = h_{c,s} f_{c,j}$ for some s and j , where $h_{c,s}$ is some monomial in h_c^β and $f_{c,j}$ is some monomial in f_c^β . This implies that $m = \psi$ and $|\psi|^2$ gives the maximal W_t -value at step $t+1$ as well. Now let's assume that $h_{c,s} \notin \mathbb{C}$. Then the combined degree of $f_{c,j}$ is less than that of ψ .

If $W_t(|f_{c,j}|^2)$ cannot be computed, then $W_t(|\psi|^2)$ cannot be computed, which gives a contradiction. Therefore, we assume that $W_t(|f_{c,j}|^2)$ can be computed.

At this point let us recall the definition of Γ_1 , Γ_2 , and Γ_3 as given in the proof of Lemma 4.1. Let Γ_1 be the set of all nonzero monomials that consist of only variables not in P_t , let Γ_2 be the set of all nonzero monomials which consist of variables both in P_t as well as variables not in P_t , and let Γ_3 be the set of all nonzero monomials which consist of only variables in P_t . Also, recall that for any monomial f if $W_t(|f|^2)$ can

be computed, then $f \in \Gamma_1$ or $f \in \Gamma_2$ only. Again, we shall write any monomial f in the form $f = \gamma_1 \gamma_2$ where γ_1 and γ_2 are monomials satisfying $\gamma_1 \in \Gamma_3$ and $\gamma_2 \in \Gamma_1$. Recall that

$$W_t(|f|^2) = \frac{1 - \sum_{i=1}^{\kappa} (\alpha_i + \hat{\alpha}_i) \mu_i}{\sum_{i=\kappa+1}^n (\alpha_i + \hat{\alpha}_i)},$$

where $(\alpha_1, \dots, \alpha_n, \hat{\alpha}_1, \dots, \hat{\alpha}_n)$ is the multiindex of the monomial $|f|^2$ of which W_t is being computed, κ is the number of variables in the leading polynomial P_t , and $W_t(|f|^2)$ is the W_t -value of the term $|f|^2$.

We shall now consider the number $W_t(|\psi|^2)$ given that $W_t(|f_{c,j}|^2)$ can be computed. From Lemma 4.1, $f_{c,j} \in \Gamma_1$ or Γ_2 . Since the monomial $h_{c,s}$ can belong to Γ_1 , Γ_2 , or Γ_3 , we shall consider three subcases below and assume that $f_{c,j} \in \Gamma_1$ or Γ_2 in each case:

- Assume that $h_{c,s} \in \Gamma_1$. Clearly, $W_t(|f_{c,j}|^2)$ and $W_t(|\psi|^2)$ both have the same numerator and the denominator of $W_t(|\psi|^2)$ is greater than that of $W_t(|f_{c,j}|^2)$ because $h_{c,s} \in \Gamma_1$. Thus, $W_t(|f_{c,j}|^2)$ is greater than $W_t(|\psi|^2)$, which is a contradiction to our hypothesis that $W_t(|\psi|^2)$ is maximal at step $t + 1$.
- Assume that $h_{c,s} \in \Gamma_2$. Here, the numerator of $W_t(|\psi|^2)$ is smaller than the numerator of $W_t(|f_{c,j}|^2)$ since $h_{c,s}$ contains a monomial in Γ_3 . Also, the denominator of $W_t(|\psi|^2)$ is greater than the denominator of $W_t(|f_{c,j}|^2)$ because $h_{c,s}$ contains a monomial in Γ_1 . Thus, $W_t(|f_{c,j}|^2)$ is greater than $W_t(|\psi|^2)$, which is again a contradiction.
- Assume that $h_{c,s} \in \Gamma_3$. Then $W_t(|f_{c,j}|^2)$ is always greater than $W_t(|\psi|^2)$ for $f_{c,j} \in \Gamma_1$ or Γ_2 since the denominators of both numbers are equal and the numerator of $W_t(|\psi|^2)$ is less than that of $W_t(|f_{c,j}|^2)$. This gives a contradiction since $W_t(|\psi|^2)$ is maximal at step $t + 1$.

From cases (a), (b), and (c) we can see that if $h_{c,s} \notin \mathbb{C}$, then $W_t(|\psi|^2)$ cannot be the maximum at step $t + 1$, and so we have a contradiction to our hypothesis in all three cases. Hence, $h_{c,s} \in \mathbb{C}$ and so $m = h_{c,s} f_{c,j}$. Clearly, $W_t(|f_{c,j}|^2)$ gives the maximal value at step $t + 1$, too. This implies that if we apply the Kolář algorithm to both $r(z)$ and $u(z)$, then the multitype entry at the $(t + 1)$ th step will be the same for both defining functions. The squares $|f_{c,j}|^2$ and $|m|^2$ appear in the expansions of $|f_c^\beta|^2$ and $|f_l^\beta|^2$, respectively. We see that regardless of the cancellation in $u(z)$, the weight obtained at the $(t + 1)$ th step remains unchanged.

Clearly, the case when $h_l(0) \neq 0$ combines the analysis for the cases when $h_l \equiv 1$ and $h_l(0) = 0$. $\square$

Proof of Theorem 1.2 Let $M \subset \mathbb{C}^{n+1}$ with $0 \in M$ be a hypersurface of which defining function is given by

$$r(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2, \quad (5.6)$$

where $f_1, \dots, f_N$ are holomorphic functions near the origin. We begin by letting $M' \subset \mathbb{C}^{n+1}$ be another hypersurface of which defining function is given as

$$u(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^S |g_j(z_1, \dots, z_n)|^2, \quad (5.7)$$

where $g_1, \dots, g_S$ are also holomorphic functions near the origin, and show that the multitype obtained by applying the Kolář algorithm to both $r(z)$ and $u(z)$ is the same provided that $\langle f_1, \dots, f_N \rangle$ and $\langle g_1, \dots, g_S \rangle$ represent the same ideal in the ring $\mathcal{O}$.

Let the ideals associated to the hypersurfaces M and M' be given by $\langle f \rangle = \langle f_1, \dots, f_N \rangle$ and $\langle g \rangle = \langle g_1, \dots, g_S \rangle$, respectively, and suppose that $\langle f \rangle = \langle g \rangle$. By Remark 5.2.2 following the statement of Lemma 5.2, we know that adding in the square of any element of the ideal $\langle f_1, \dots, f_N \rangle$ does not modify the multitype because that element can be written in terms of the generators $f_1, \dots, f_N$ with coefficients in $\mathcal{O}$. Since $\langle f_1, \dots, f_N \rangle = \langle g_1, \dots, g_S \rangle$, each g_j is an element of $\langle f_1, \dots, f_N \rangle$ and can be written in terms of $f_1, \dots, f_N$ with coefficients in $\mathcal{O}$. Therefore, (5.6) has the same multitype at the origin as

$$r_1(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2 + |g_1(z_1, \dots, z_n)|^2,$$

and inductively, the same multitype at the origin as

$$r_S(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2 + \sum_{k=1}^S |g_k(z_1, \dots, z_n)|^2.$$

Now, we apply the argument in reverse. Since $\langle g_1, \dots, g_S \rangle = \langle f_1, \dots, f_N \rangle$, each f_j is an element of $\langle g_1, \dots, g_S \rangle$ and can be written in terms of $g_1, \dots, g_S$ with coefficients in $\mathcal{O}$. Therefore, by Remark 5.2.2, (5.7) has the same multitype at the origin as

$$u_1(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{k=1}^S |g_k(z_1, \dots, z_n)|^2 + |f_1(z_1, \dots, z_n)|^2,$$

and inductively, as

$$r_S(z) = 2\operatorname{Re}(z_{n+1}) + \sum_{k=1}^S |g_k(z_1, \dots, z_n)|^2 + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2.$$

We conclude that $r(z)$ and $u(z)$ have the same multitype at the origin, namely that the multitype is an invariant of the ideal of generators. $\square$

6 An Ideal Restatement of the Kolář Algorithm

In this section, we give the ideal restatement of the Kolář algorithm for the multitype computation for sum of squares domains in $\mathbb{C}^{n+1}$. Before that, we shall consider the following:

Let $M \subset \mathbb{C}^{n+1}$ be a hypersurface given by $r = 0$ with

$$r = 2\operatorname{Re}(z_{n+1}) + \sum_{k=1}^N |f_k(z_1, \dots, z_n)|^2,$$

where $f_1, \dots, f_N$ are holomorphic functions defined on a neighborhood of the origin. Assume that the D'Angelo 1-type is finite at the origin and that $\beta = \lceil \Delta_1(M, 0) \rceil$, the ceiling of the D'Angelo 1-type of M at the origin. We truncate each holomorphic function f_k at the order β and let $f_{k,i}$ be the i th monomial of the generator f_k^β after ordering by vanishing order. The ideal corresponding to the defining function r is given as $\mathcal{I} = (z_{n+1}, f_1^\beta, \dots, f_N^\beta)$. We know that the term z_{n+1} has weight 1. Following the original algorithm of Kolář, we shall ignore the term z_{n+1} and work with the corresponding ideal $\mathcal{I} = (f_1^\beta, \dots, f_N^\beta)$.

From Theorem 1.1 we know that all leading polynomials produced are sums of squares. Therefore, any leading polynomial P_j can be written in the form:

$$P_j = \sum_{k=1}^N \left| \sum_{i=1}^{v_k} f_{k,a_i} \right|^2, \quad (6.1)$$

where the f_{k,a_i} 's are the monomials from the generator f_k^β of weighted degree $\frac{1}{2}$ with respect to Λ_j . We will associate to every leading polynomial P_j the ideal $\mathcal{I}_{P_j}$ given by

$$\mathcal{I}_{P_j} = \left(\sum_{i=1}^{v_1} f_{1,a_i}, \dots, \sum_{i=1}^{v_N} f_{N,a_i} \right). \quad (6.2)$$

It is convenient to introduce notation for each square in P_j . Let $P_{j,k} = \left| \sum_{i=1}^{v_k} f_{k,a_i} \right|^2$. Then its associated ideal $\mathcal{I}_{P_{j,k}}$ can be expressed as

$$\mathcal{I}_{P_{j,k}} = \left(\sum_{i=1}^{v_k} f_{k,a_i} \right) = (f_{k,a_1} + \dots + f_{k,a_{v_k}}) \quad (6.3)$$

Clearly,

$$\mathcal{I}_{P_j} = \sum_{k=1}^N \mathcal{I}_{P_{j,k}}. \quad (6.4)$$

Recall also that each monomial in every leading polynomial is of weighted degree one with respect to the corresponding weight. As a result, the weighted degree of any monomial f_{k,a_i} is exactly one half with respect to the corresponding weight.

Thus, given $\mathcal{I}_{P_{j,k}} = (f_{k,a_1} + \cdots + f_{k,a_{v_k}})$, the f_{k,a_i} 's are exactly the monomials from the generator f_k^β of weighted degree $\frac{1}{2}$ with respect to Λ_j .

Set the ideal $\mathcal{I} = \mathcal{I}_0$. For $j \geq 1$, the ideals $\mathcal{I}_{P_j}$, $\mathcal{I}_j$, and $\mathcal{I}_{P_{j,k}}$ can be described as follows:

1. $\mathcal{I}_{P_j}$ is the ideal of which generators are precisely the terms from the generators of the ideal $\mathcal{I}_{j-1}$ having weighted order exactly $\frac{1}{2}$ with respect to the weight Λ_j . We refer to the ideal $\mathcal{I}_{P_j}$ as the *leading polynomial ideal*.
2. The ideal $\mathcal{I}_j$ is the ideal obtained after applying the chosen Λ_j -homogeneous transformation, which makes the generators of $\mathcal{I}_{P_j}$ to be independent of the largest number of variables, to $\mathcal{I}_j$. Simply put, $\mathcal{I}_j$ is the ideal obtained after changing variables in the ideal $\mathcal{I}_{j-1}$.
3. $\mathcal{I}_{P_{j,k}}$ is the principal ideal of which generator is the sum of monomials in the generator f_k^β with weighted degree exactly $\frac{1}{2}$ with respect to the weight Λ_j . The ideal $\mathcal{I}_{P_{j,k}}$ is the zero ideal if no monomial in the generator f_k^β has weighted degree $\frac{1}{2}$ with respect to the weight Λ_j .

The Kolář Algorithm (Ideal Version) Set the ideal $\mathcal{I} = \mathcal{I}_0$, and compute the vanishing order at the origin of $\mathcal{I}_0$, which is the same as the degree v_1 of the lowest order monomial in $\mathcal{I}_0$. We define the Bloom-Graham type as twice the vanishing order of $\mathcal{I}_0$. This gives the first entry of the multitype m_1 and so let $m_1 = 1/\mu_1$, where $\mu_1 = 2v_1$. Set the first weight to be $\Lambda_1 = (\mu_1, \dots, \mu_1)$.

In the second step, consider all Λ_1 -homogeneous transformations, and choose one that will make the set of all generators of the leading polynomial ideal $\mathcal{I}_{P_1}$ to be independent of the largest number of variables. Denote this number by d_1 . In the local coordinates after such a Λ_1 -homogeneous transformation, we obtain that $\mathcal{I}_{P_1}$ is the ideal of which generators consist of those monomials in the variables $z_1, \dots, z_{n-d_1}$, which are of weighted degree $1/2$ with respect to Λ_1 . Apply the chosen Λ_1 -homogeneous transformation to the ideal $\mathcal{I}_0$ to obtain the ideal $\mathcal{I}_1$. The rest of the terms from the generators in $\mathcal{I}_1$, which are not in $\mathcal{I}_{P_1}$, have weighted degrees strictly greater than $\frac{1}{2}$ with respect to Λ_1 .

We shall now give a slightly modified version of Kolář's Θ_1 and W_1 . If $\alpha^{k,j} = (\alpha_1^{k,j}, \dots, \alpha_n^{k,j})$ is the multiindex of a monomial $f_{k,j}$ from any generator of $\mathcal{I}_1$, which is not in $\mathcal{I}_{P_1}$, then $f_{k,j}$ is of the form:

$$f_{k,j} = C_{k,j}^1 z^{\alpha^{k,j}} \quad \text{and} \quad |\alpha^{k,j}|_{\Lambda_1} > \frac{1}{2}.$$

Let

$$\Theta_1 = \left\{ \alpha^{k,j} \mid C_{k,j}^1 \neq 0 \text{ and } \sum_{i=1}^{n-d_1} \alpha_i^{k,j} \mu_i < \frac{1}{2} \right\}.$$

For every $\alpha^{k,j} \in \Theta_1$,

$$W_1(\alpha^{k,j}) = \frac{\frac{1}{2} - \sum_{i=1}^{n-d_1} \alpha_i^{k,j} \mu_i}{\sum_{i=n-d_1+1}^n \alpha_i^{k,j}}. \quad (6.5)$$

The next weight Λ_2 is defined by letting

$$\lambda_i^2 = \max_{\alpha^{k,j} \in \Theta_1} W_1(\alpha^{k,j})$$

for $i > n - d_1$, and $\lambda_i^2 = \mu_i$ for $i \leq n - d_1$. To complete the second step, we let $\mathcal{I}_{P_2}$ be the second leading polynomial ideal corresponding to the weight Λ_2 . The generators of $\mathcal{I}_{P_2}$ depend on more than $n - d_1$ variables.

We proceed by induction. At the step t , for $t > 2$, we consider all Λ_{t-1} -homogeneous transformations and choose one that makes the generators of the leading polynomial ideal $\mathcal{I}_{P_{t-1}}$ to be independent of the largest number of variables. Denote this number by d_{t-1} . Apply this Λ_{t-1} -homogeneous transformation to the previous ideal $\mathcal{I}_{t-2}$ in the $(t-1)$ th step to obtain the ideal $\mathcal{I}_{t-1}$. We know from the Kolář algorithm that the number of multitype entries that are added at each step of the computation depends on the difference $(d_{t-2} - d_{t-1})$. We consider two cases:

Case 1: Assume that $d_{t-2} > d_{t-1}$. Again recall that for any weight Λ that is smaller than Λ_{t-1} with respect to the lexicographic ordering, Λ -adapted coordinates are also Λ_{t-1} -adapted. This implies that we get $(d_{t-2} - d_{t-1})$ multitype entries

$$\mu_{n-d_{t-2}+1} = \cdots = \mu_{n-d_{t-1}} = \lambda_{n-d_{t-2}+1}^{t-1}$$

and let $\lambda_i^t = \mu_i$ for $i \leq n - d_{t-2}$. Here, $\mathcal{I}_{P_{t-1}}$ is the ideal of which generators are sums of monomials in the variables $z_1, \dots, z_{n-d_{t-1}}$ that are Λ_{t-1} -homogeneous of weighted degree $\frac{1}{2}$. To obtain λ_i^t for $i > n - d_{t-1}$, we consider the rest of the monomials from the generators in $\mathcal{I}_{t-1}$ that are not in $\mathcal{I}_{P_{t-1}}$. Using these monomials that have weighted degree strictly greater than $\frac{1}{2}$ with respect to Λ_{t-1} , we define Θ_{t-1} and compute W_{t-1} in a similar way as in step two. Such monomials are of the form $f_{k,j} = C_{k,j}^{t-1} z^{\alpha^{k,j}}$ for multiindex $\alpha^{k,j} = (\alpha_1^{k,j}, \dots, \alpha_n^{k,j})$ satisfying $|\alpha^{k,j}|_{\Lambda_{t-1}} > \frac{1}{2}$. Thus,

$$\Theta_{t-1} = \left\{ \alpha^{k,j} \mid C_{k,j}^{t-1} \neq 0 \text{ and } \sum_{i=1}^{n-d_{t-1}} \alpha_i^{k,j} \mu_i < \frac{1}{2} \right\}.$$

For every $\alpha^{k,j} \in \Theta_{t-1}$,

$$W_{t-1}(\alpha^{k,j}) = \frac{\frac{1}{2} - \sum_{i=1}^{n-d_{t-1}} \alpha_i^{k,j} \mu_i}{\sum_{i=n-d_{t-1}+1}^n \alpha_i^{k,j}}. \quad (6.6)$$

So for the remaining multitype entries of Λ_j , we let

$$\lambda_i^t = \max_{\alpha^{k,j} \in \Theta_{t-1}} W_{t-1}(\alpha^{k,j}),$$

for $i > n - d_{t-1}$.

Case 2 Assume that $d_{t-1} = d_{t-2}$. There are zero multitype entries computed in this case, and so we only determine λ_i^t for $t > n - d_{t-1}$ using (6.6). This completes the step t of the algorithm.

We can, thus, establish a one-to-one correspondence between the leading polynomial P_t for $t \geq 1$ and the intermediate ideal $\mathcal{I}_{P_t}$ introduced above. Since working with ideals of holomorphic functions is often easier than with real-valued polynomials, the restatement of the Kolář algorithm simplifies multitype computations for a sum of squares domain. We work with considerably fewer terms in the case of the ideals as compared to sums of squares. In particular, for each modulus square of a generator consisting of m monomials, Kolář's original algorithm involves working with m squares plus $\binom{m}{2}$ cross terms, whereas this restatement in terms of ideals involves computations for only m monomials.

We give a corollary to Theorem 1.1:

Corollary 6.0.1 *Let $M \subset \mathbb{C}^{n+1}$ be a hypersurface given by $r = 0$ with*

$$r = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2,$$

where $f_1, \dots, f_N$ are holomorphic functions defined on a neighborhood of the origin and assume that the D'Angelo 1-type of M at the origin is finite. Then for $\ell \geq 1$, each monomial from every generator of the leading polynomial ideal $\mathcal{I}_{P_\ell}$ obtained at the ℓ th step of the Kolář algorithm has weighted degree $\frac{1}{2}$ with respect to the weight Λ_ℓ .

Proof Let f_k^β be the Taylor expansion of the holomorphic function f_k to the order β , where β is the ceiling of the D'Angelo 1-type. We order the generators by vanishing order and let $\mathcal{I} = (f_1^\beta, \dots, f_N^\beta)$. Now assume that the vanishing order of the ideal $\mathcal{I}$ is $\nu > 0$. Then the Bloom-Graham type is precisely 2ν and the weight $\mu_1 = \frac{1}{2\nu}$ with $\Lambda_1 = (\frac{1}{2\nu}, \dots, \frac{1}{2\nu})$. Thus, $\mathcal{I}_{P_1}$ is not the zero ideal.

For every k such that $\mathcal{I}_{P_{1,k}}$ is not the zero ideal,

$$\mathcal{I}_{P_{1,k}} = (f_{k,1}, \dots, f_{k,m_k}),$$

where each monomial $f_{k,i}$, for $1 \leq i \leq m_k$, has weighted degree $\frac{1}{2}$ with respect to the weight Λ_1 . Next, assume that in the second step, the ideal $\mathcal{I}_{P_{1,k}}$ is the same as the ideal $\mathcal{I}_{P_{2,k}}$. We know that the entries corresponding to each variable in the monomial $f_{k,i}$, for $1 \leq i \leq m_k$, are the same in both weights Λ_1 and Λ_2 . Therefore, each monomial from the generator $\sum_{i=1}^{m_k} f_{k,i}$ has weighted degree $\frac{1}{2}$ with respect to Λ_2 as well. Assume

that the principal ideal $\mathcal{I}_{P_{2,k}}$ is generated by the sum $\sum_{i=1}^{m_k} f_{k,i} + \sum_{j=1}^{v_k} f_{k,b_j}$. Since the new sum $\sum_{j=1}^{v_k} f_{k,b_j}$ corresponds to the new weight Λ_2 , each monomial f_{k,b_j} has weighted degree $\frac{1}{2}$ with respect to Λ_2 . Therefore, every monomial in the generator of the ideal $\mathcal{I}_{P_{2,k}}$ has weighted degree $\frac{1}{2}$ with respect to Λ_2 .

Next, we assume that for $\ell \geq 2$

$$\mathcal{I}_{P_{\ell,k}} = (f_{k,a_1} + \cdots + f_{k,a_{v_k}}),$$

where each monomial f_{k,a_i} , for $1 \leq i \leq v_k$, has weighted degree exactly equal to $\frac{1}{2}$ with respect to the weight Λ_ℓ .

Now, assume that at step $\ell + 1$, the ideal $\mathcal{I}_{P_{\ell+1,k}}$ is the same as the ideal $\mathcal{I}_{P_{\ell,k}}$. Every monomial in the generator of $\mathcal{I}_{P_{\ell+1,k}}$ also has weighted degree equal to $\frac{1}{2}$ with respect to the weight $\Lambda_{\ell+1}$ because even though Λ_ℓ is not the same as $\Lambda_{\ell+1}$, the weight corresponding to each variable in $f_{k,a_{v_k}}$ is the same in both weights Λ_ℓ and $\Lambda_{\ell+1}$.

Next, assume that at step $\ell + 1$ the sum $\sum_{i=1}^{u_k} f_{k,b_{u_k}}$ is added to the sum $\sum_{i=1}^{v_k} f_{k,a_{v_k}}$ to obtain the generator of the ideal

$$\mathcal{I}_{P_{\ell+1,k}} = (f_{k,a_1} + \cdots + f_{k,a_{v_k}} + f_{k,b_1} + \cdots + f_{k,b_{u_k}}).$$

This implies that each monomial $f_{k,b_{u_k}}$ has weighted degree $\frac{1}{2}$ with respect to the weight $\Lambda_{\ell+1}$. Again, each monomial $f_{k,a_{v_k}}$ is of weighted degree $\frac{1}{2}$ with respect to the weight $\Lambda_{\ell+1}$ since the weight corresponding to each variable in $f_{k,a_{v_k}}$ is the same in both weights Λ_ℓ and $\Lambda_{\ell+1}$. Thus, every monomial from the generator of the ideal $\mathcal{I}_{P_{\ell+1,k}}$ has weighted degree $\frac{1}{2}$ with respect to the weight $\Lambda_{\ell+1}$. $\square$

The example that follows is the ideal restatement of the Kolář algorithm applied to the defining function given in Example 1.

Example 2 Let $M \subset \mathbb{C}^4$ be a sum of squares domain given by the defining function

$$r = 2\operatorname{Re}(z_4) + |z_1 - z_2 + z_3|^2 + |z_1^2 - z_2^2|^2.$$

The associated ideal then becomes

$$\mathcal{I} = (z_1 - z_2 + z_3^2, z_1^2 - z_2^2) = \mathcal{I}_0.$$

The vanishing order here equals one, and so the Bloom-Graham type is 2, which implies that $\mu_1 = \frac{1}{2}$ and $\Lambda_1 = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$. Hence,

$$\mathcal{I}_{P_1} = (z_1 - z_2) = \mathcal{I}_{P_{1,1}}.$$

So $\mathcal{I}_{P_{1,2}}$ is the zero ideal. We consider all Λ_1 -homogeneous transformations and choose

$$\tilde{z}_1 = z_1 - z_2 \quad \tilde{z}_j = z_j, \quad j = 2, 3, 4$$

to make $\mathcal{I}_{P_1}$ independent of the largest number of variables. We shall ignore $\sim$ where no confusion arises. Thus, in the new coordinates,

$$\mathcal{I}_{P_1} = (z_1) = \mathcal{I}_{P_{1,1}}$$

with $d_1 = 2$, and so $\mathcal{I}_0$ becomes $\mathcal{I}_1 = (z_1 + z_3^2, z_1^2 + 2z_1z_2)$. The maximum of W_1 obtained by computing W_1 for each monomial in the generators of $\mathcal{I}_1$ is $\frac{1}{4}$ and $\Lambda_2 = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$. Hence,

$$\mathcal{I}_{P_2} = (z_1 + z_3^2),$$

where $\mathcal{I}_{P_{2,1}} = \mathcal{I}_{P_2}$ and $\mathcal{I}_{P_{2,2}}$ is the zero ideal. Next, consider all Λ_2 -homogeneous transformations and choose

$$\tilde{z}_1 = z_1 + z_3^2, \quad \tilde{z}_j = z_j, \quad j = 2, 3, 4,$$

which makes $\mathcal{I}_{P_2}$ to be dependent on only the variable z_1 . Again, we shall ignore $\sim$ where no confusion arises. In the new coordinates,

$$\mathcal{I}_{P_2} = (z_1)$$

with $d = 2$. The ideal $\mathcal{I}_{P_{2,1}} = \mathcal{I}_{P_2}$, and $\mathcal{I}_{P_{2,2}}$ is the zero ideal. $\mathcal{I}_1$ in the new coordinates becomes

$$\mathcal{I}_2 = (z_1, z_1^2 + 2z_1z_3^2 + z_3^4 + 2z_1z_2 - 2z_2z_3^2).$$

Computing W_2 for each monomial in the generators of $\mathcal{I}_2$ yields the set of numbers $\{\frac{1}{6}, \frac{1}{8}\}$ of which maximum, $\max W_2$, is $\frac{1}{6}$, and so $\Lambda_3 = (\frac{1}{2}, \frac{1}{6}, \frac{1}{6})$. Also,

$$\mathcal{I}_{P_3} = (z_1, -2z_2z_3^2) = (z_1, 2z_2z_3^2).$$

Here $\mathcal{I}_{P_{3,1}} = (z_1)$ and $\mathcal{I}_{P_{3,2}} = (2z_2z_3^2)$. No Λ_3 -homogeneous transformation can make $\mathcal{I}_{P_3}$ to be independent of any variables, and so $d_3 = 0$. Thus, the multitype weight $\Lambda^* = \Lambda_3$, and the final leading polynomial ideal is

$$\mathcal{I}_{P_3} = (z_1, z_2z_3^2).$$

7 Changes of Variables in the Kolář Algorithm

We now give an explicit construction of the polynomial transformations that are performed in the Kolář algorithm. We begin this construction by first relating the weighted homogeneous polynomial transformations to pairs of row-column operations on the Levi matrix of a sum of squares domain. More specifically, we show

that each polynomial transformation corresponds to some finite sequence of row and column operations. Lemma 7.1 below establishes this correspondence.

Throughout this section, we shall assume the following:

Let $M \subset \mathbb{C}^{n+1}$ be the boundary of a sum of squares domain defined by $\{r < 0\}$, where

$$r = 2\operatorname{Re}(z_{n+1}) + \sum_{j=1}^N |f_j(z_1, \dots, z_n)|^2,$$

and $f_1, \dots, f_N$ are holomorphic functions in the neighborhood of the origin. Let

$$r_0 = 2\operatorname{Re}(z_{n+1}) + P(z, \bar{z})$$

be the defining function of the model hypersurface M_0 of M , where $P(z, \bar{z})$ is a polynomial of weighted degree 1 with respect to the multitype weight at the origin Λ^* of M . Let A be the $n \times n$ Levi matrix of the model $M_0 \subset \mathbb{C}^{n+1}$, where we ignore the contribution of the $(n+1)^{\text{st}}$ coordinate as the holomorphic polynomials in the sum of squares do not depend on it as mentioned in the introduction.

Lemma 7.1 *Assume that the D'Angelo 1-type of the hypersurface M in $\mathbb{C}^{n+1}$ at 0 is finite. Let $i \in \{1, \dots, n\}$ be fixed, and let $h \in \mathbb{C}[z]$ for $z = (z_1, \dots, z_n)$ be a nonzero monomial independent of z_i . Let h_ℓ denote the derivative of h with respect to the variable z_ℓ , which is $\partial_{z_\ell} h$ with $\ell \in \{1, \dots, n\} \setminus \{i\}$. Furthermore, let $h_\ell(\tau)$ denote h_ℓ with every factor of z_l replaced by a factor of τ . Performing the elementary row and column operations $R_\ell - h_\ell R_i \rightarrow R_\ell$ and $C_\ell - \bar{h}_\ell C_i \rightarrow C_\ell$ on the Levi matrix A of r_0 for all variables z_ℓ in h corresponds to the polynomial transformation*

$$\tilde{z}_i = z_i + \int_0^{z_\ell} h_\ell(\tau) d\tau = z_i + h; \quad \tilde{z}_\omega = z_\omega \text{ for } \omega \neq i$$

in the sense that the new matrix $\tilde{A}$ obtained after these elementary operations is Hermitian and is the Levi matrix of the new defining function of the sum of squares domain after the change of variables $z_\omega \rightarrow \tilde{z}_\omega$ for $\omega = 1, \dots, n+1$ has taken place.

Remark 7.1.1 The reader should note that while only variables $z_1, \dots, z_n$ play a role in the behavior of the Levi matrix, $\mathbb{C}^{n+1}$ is the underlying space, so all changes of variables described in this section will take place in $\mathbb{C}^{n+1}$ and leave z_{n+1} unchanged.

Proof Suppose that the defining function r_0 of the model hypersurface M_0 is of the form:

$$r_0 = 2\operatorname{Re}(z_{n+1}) + \sum_{t=1}^N |g_t|^2,$$

where $P(z, \bar{z}) = \sum_{t=1}^N |g_t|^2$ and $g_t = \sum_{l=1}^{b_t} m_{t,l}$ is a polynomial consisting of monomials $m_{t,l}$ in the variables $z_1, \dots, z_n$ since $P(z, \bar{z})$ cannot depend on the variable z_{n+1} .

Let $m_{t,l} = C_{t,l} \prod_{\delta=1}^n z_{\delta}^{\alpha_{\delta}^{t,l}}$ with $C_{t,l} \in \mathbb{C}$. For each t and for $l_1, l_2 \in \{1, \dots, b_t\}$, every monomial from the expansion of $|g_t|^2$ can be written as

$$m_{t,l_1} \bar{m}_{t,l_2} = C_{t,l_1} \bar{C}_{t,l_2} \prod_{\delta=1}^n z_{\delta}^{\alpha_{\delta}^{t,l_1}} \bar{z}_{\delta}^{\hat{\alpha}_{\delta}^{t,l_2}}.$$

By writing each term $m_{t,l_1} \bar{m}_{t,l_2}$ for all t in the new coordinates, we obtain $P(z, \bar{z})$ in the new coordinates. Hence, it suffices to show that applying the specified elementary row and column operations to the Levi matrix of the monomial $m_{t,l_1} \bar{m}_{t,l_2}$ corresponds to the polynomial transformation $\tilde{z}_i = z_i + h$; $\tilde{z}_{\omega} = z_{\omega}$ for $\omega \neq i$.

Denote by D the (i, j, k, u) submatrix of the Levi matrix of the monomial $m_{t,l_1} \bar{m}_{t,l_2}$, and let $D = (d_{e\bar{k}})_{e,\kappa=i,j,k,u}$, where $d_{e\bar{k}}$ is given by

$$d_{e\bar{k}} = C_{t,l_1} \bar{C}_{t,l_2} \alpha_e^{t,l_1} \hat{\alpha}_{\kappa}^{t,l_2} z_e^{\alpha_e^{t,l_1}-1} \bar{z}_{\kappa}^{\hat{\alpha}_{\kappa}^{t,l_2}-1} \prod_{\substack{\delta=1 \\ \delta \neq e,\kappa}}^n z_{\delta}^{\alpha_{\delta}^{t,l_1}} \bar{z}_{\delta}^{\hat{\alpha}_{\delta}^{t,l_2}}.$$

Let $h = C z_{a_1}^{\beta_{a_1}}, \dots, z_{a_s}^{\beta_{a_s}}$, where $C \in \mathbb{C}$, $\beta = (\beta_1, \dots, \beta_n)$ is a multiindex, and $a_1, \dots, a_s \in \{1, \dots, n\} \setminus \{i\}$. Now, assume that $j, k \in \{a_1, \dots, a_s\}$ with $j \neq k$ and $u \notin \{a_1, \dots, a_s\}$. Perform the elementary operations $R_{\ell} - h_{\ell} R_i \rightarrow R_{\ell}$ and $C_{\ell} - \bar{h}_{\ell} C_i \rightarrow C_{\ell}$ for all variables z_{ℓ} in h on D to get

$$\begin{pmatrix} d_{i\bar{i}} & d_{i\bar{j}} - \bar{h}_j d_{i\bar{i}} & d_{i\bar{k}} - \bar{h}_k d_{i\bar{i}} & d_{i\bar{u}} \\ d_{j\bar{i}} - h_j d_{i\bar{i}} & d_{j\bar{j}} - h_j d_{i\bar{j}} - \bar{h}_j d_{j\bar{i}} + |h_j|^2 d_{i\bar{i}} & d_{j\bar{k}} - h_j d_{i\bar{k}} - \bar{h}_k d_{j\bar{i}} + h_j \bar{h}_k d_{i\bar{i}} & d_{j\bar{u}} - h_j d_{i\bar{u}} \\ d_{k\bar{i}} - h_k d_{i\bar{i}} & d_{k\bar{j}} - \bar{h}_j d_{k\bar{i}} - h_k d_{i\bar{j}} + \bar{h}_j h_k d_{i\bar{i}} & d_{k\bar{k}} - h_k d_{i\bar{k}} - \bar{h}_k d_{k\bar{i}} + |h_k|^2 d_{i\bar{i}} & d_{k\bar{u}} - h_k d_{i\bar{u}} \\ d_{u\bar{i}} & d_{u\bar{j}} - \bar{h}_j d_{u\bar{i}} & d_{u\bar{k}} - \bar{h}_k d_{u\bar{i}} & d_{u\bar{u}} \end{pmatrix}.$$

The matrix above is the Levi matrix for the monomial in the new coordinates

$$\tilde{m}_{l_1} \bar{\tilde{m}}_{l_2} = C_{t,l_1} \bar{C}_{t,l_2} (\tilde{z}_i - h)^{\alpha_e^{t,l_1}} (\bar{\tilde{z}}_i - \bar{h})^{\hat{\alpha}_{\kappa}^{t,l_2}} \prod_{\delta \neq i} z_{\delta}^{\alpha_{\delta}^{t,l_1}} \bar{z}_{\delta}^{\hat{\alpha}_{\delta}^{t,l_2}},$$

which is obtained after applying the polynomial transformation $\tilde{z}_i = z_i + h$; $\tilde{z}_{\omega} = z_{\omega}$ for $\omega \neq i$ to $m_{t,l_1} \bar{m}_{t,l_2}$.

We now prove that the matrix $\tilde{A}$ is Hermitian. Since the matrix A is Hermitian, we will show that applying the operation $R_{\ell} - h_{\ell} R_i \rightarrow R_{\ell}$ and $C_{\ell} - \bar{h}_{\ell} C_i \rightarrow C_{\ell}$ to A gives a matrix that is Hermitian as well. Let $A = (a_{k\bar{l}})_{1 \leq k, l \leq n}$ be the Levi matrix. Apart from row ℓ and column ℓ , there is no change to A , which is Hermitian. Let $a_{\ell\bar{k}}$ be an entry in row ℓ . Then the entry $a_{k\bar{\ell}}$ is in column ℓ and satisfies the property that $a_{\ell\bar{k}} = \bar{a}_{k\bar{\ell}}$. Performing the elementary operations $R_{\ell} - h_{\ell} R_i \rightarrow R_{\ell}$ and $C_{\ell} - \bar{h}_{\ell} C_i \rightarrow C_{\ell}$ on A gives a new matrix $\tilde{A}$ with $a_{\ell\bar{k}} - h_{\ell} a_{i\bar{k}}$ in row ℓ and $a_{k\bar{\ell}} - \bar{h}_{\ell} a_{k\bar{i}}$ in column ℓ . Now

$$a_{\ell\bar{k}} - h_{\ell} a_{i\bar{k}} = \bar{a}_{k\bar{\ell}} - \bar{h}_{\ell} \bar{a}_{k\bar{i}} = \overline{a_{k\bar{\ell}} - \bar{h}_{\ell} a_{k\bar{i}}}.$$

Thus, the new matrix $\tilde{A}$ is Hermitian as well. $\square$

Proposition 1.3 stated in the introduction provides a general condition for the existence of an allowable polynomial transformation via the elementary row and column operations performed on the Levi matrix of a leading polynomial at some step of the Kolář algorithm. It turns out that the restrictive definition of dependency mentioned in the introduction is a necessary and sufficient condition for the existence of allowable polynomial transformations. At this point, we give the proof of Proposition 1.3 and remark that the proof is constructive in the sense that we will show the allowable polynomial transformation on P_j arises as a composition of polynomial transformations corresponding to elementary row and column operations on A_{P_j} .

Proof of Proposition 1.3 Let k be given, and denote by R_k and C_k the k th row and k th column of the matrix A_{P_j} , respectively. Suppose that the k th row of A_{P_j} is dependent. This implies that C_k must also be dependent since A_{P_j} is Hermitian. Hence, we can write both R_k and C_k , respectively, as follows:

$$R_k = \sum_{\substack{l=1 \\ l \neq k}}^n \beta^l R_l \text{ and } C_k = \sum_{\substack{l=1 \\ l \neq k}}^n \bar{\beta}^l C_l, \quad (7.1)$$

where $\beta^l \in \mathbb{C}[z]$ for every l , $1 \leq l \leq n$. As proven in Theorem 1.1, the leading polynomial P_j is a sum of squares, and so we write $P_j = \sum_{s=1}^m |\phi^s|^2$, where each ϕ^s is a nonzero polynomial with vanishing order greater than or equal to 1. Denote by $a_{k\bar{i}}$ the entry in the $(k, \bar{i})$ position of the matrix A_{P_j} , for $t = 1, \dots, n$. Therefore, for $k \neq l$ the entries in R_k and R_l are given by

$$a_{k\bar{i}} = \sum_{s=1}^m \phi_k^s \bar{\phi}_i^s \text{ and } a_{l\bar{i}} = \sum_{s=1}^m \phi_l^s \bar{\phi}_i^s, \quad (7.2)$$

and the entries in C_k and C_l are given by

$$a_{i\bar{k}} = \sum_{s=1}^m \phi_i^s \bar{\phi}_k^s \text{ and } a_{i\bar{l}} = \sum_{s=1}^m \phi_i^s \bar{\phi}_l^s \quad (7.3)$$

respectively, where $\phi_j^s = \frac{\partial \phi^s}{\partial z_j}$.

We will show that there exist some elementary row and column operations that make R_k to be identically zero and also make every monomial in A_{P_j} to be independent of the variable z_k .

To that end, we see from (7.1) that every pair $R_k - \beta^l R_l$; $C_k - \bar{\beta}^l C_l$ for which the polynomial β^l is nonzero requires corresponding elementary row and column operations in the exact form expressed in the pair. If $\beta^l = 0$ or $R_l \equiv 0$, then no elementary row or column operation is required. Therefore, assume that β^l is nonzero and that R_l is not identically zero. Let $\beta^l(\zeta_k)$ be β^l with each factor of z_k replaced by

a factor of ζ_k . Then $\int_0^{z_k} \beta^l(\zeta_k) d\zeta_k$ must contain the variable z_k together with all the other variables in β^l . Recall that $\beta^l \in \mathbb{C}[z]$, so β^l does not depend on any variable $\bar{z}_v$. Hence, we shall investigate all monomials in A_{P_j} containing the variable z_k .

We recall at this point that by applying the operator $\partial_{z_k} \partial_{\bar{z}_t}$ to P_j , we obtain in row R_k the derivatives of all monomials containing the variable z_k . Let $a_{k\bar{t}}$ be an entry in row R_k . Now, because R_k is dependent, every monomial u in $a_{k\bar{t}}$ arises as the product of a monomial p in β^l for some l with a monomial q in entry $a_{l\bar{t}}$. So $u = pq$, but since u comes from differentiation by $\partial_{z_k} \partial_{\bar{z}_t}$, P_j must contain a monomial $m = uz_k \bar{z}_t$. If $u \in \mathbb{C}$, then no entries in A_{P_j} , except for those in R_k , contain derivatives from m . If u has positive vanishing order, then u depends on at least one variable z_v or $\bar{z}_v$ for some v . Since P_j is real valued, it contains both m and $\bar{m}$. Therefore, without loss of generality, we can assume u depends on z_v ; otherwise, we work with $\bar{u}$. Since u depends on z_v , the entry $a_{v\bar{t}}$ in the v th row R_v contains the monomial $\partial_{z_v} \partial_{\bar{z}_t} m \neq 0$, which has at least one factor of z_k . We seek to eliminate all such monomials containing variable z_k from the matrix A_{P_j} .

Set $\gamma^l = \int_0^{z_k} \beta^l(\zeta_k) d\zeta_k$, and let $\gamma^l = \sum_{b=1}^e m^{l,b}$, where $m^{l,b}$ is a nonzero monomial containing the variable z_k for all $b \geq 1$. We recall from Lemma 7.1 that for any nonzero monomial m in the leading polynomial, if we perform the pair of elementary operations $R_v - \partial_{z_v} m R_\ell \rightarrow R_v$ and $C_v - \partial_{\bar{z}_v} \bar{m} C_\ell \rightarrow C_v$ for all variables z_v in m , then this pair corresponds to the polynomial transformation $\tilde{z}_\ell = z_\ell + \int_0^{z_v} \partial_\tau m(\tau) d\tau = z_\ell + m$; $\tilde{z}_\omega = z_\omega$ for $\omega \neq \ell$, where $\partial_\tau m(\tau)$ is $\partial_{z_v} m$ with each factor of z_v replaced by a factor of τ .

Now, for each monomial $m^{l,b}$ in γ^l , $b = 1, \dots, e$, we perform the elementary row and column operations $R_v - \partial_{z_v} m^{l,b} R_l \rightarrow R_v$ and $C_v - \partial_{\bar{z}_v} \bar{m}^{l,b} C_l \rightarrow C_v$ for every variable z_v in $m^{l,b}$. The composition of all of these polynomial transformations $\mathcal{S}$ is given by $\tilde{z}_l = z_l + \gamma^l$ for every l such that $\beta^l \neq 0$ and $\tilde{z}_\omega = z_\omega$ for all $\omega \neq l$, where $1 \leq \omega \leq n+1$. Note that (7.1) implies that γ^l has the same weight as z_l in Λ_j because P_j only contains terms of weight 1 with respect to Λ_j , so $\mathcal{S}$ is Λ_j -homogeneous as needed.

After all the elementary row and column operations corresponding to the polynomial transformation $\mathcal{S}$ have taken place, the entries in R_k are

$$a'_{k\bar{t}} = \sum_{s=1}^m \phi_k^s \bar{\phi}_t^s - \sum_{l=1}^n \beta^l \left(\sum_{s=1}^m \phi_l^s \bar{\phi}_t^s \right) = a_{k\bar{t}} - \sum_{l=1}^n \beta^l a_{l\bar{t}} \equiv 0 \quad (7.4)$$

as a consequence of (7.1). A similar argument holds for the entries in C_k , which we denote by $a'_{t\bar{k}}$, namely (7.1) implies that $a'_{t\bar{k}} \equiv 0$. Therefore, all entries in the k th row and column of the matrix A_{P_j} are identically zero after the change of variables $\mathcal{S}$ has been performed. Now, assume that the leading polynomial P_j still contains the variable z_k after the given change of variables has been performed on it and some cancellation occurs. Since the leading polynomial P_j is a sum of squares, in the expansion of P_j besides the cross terms, which could possibly cancel each other, we would have at least

two squares of monomials containing z_k . From the above discussion, it is clear that by performing these elementary row and column operations on A_{P_j} , all monomials containing the variable z_k in any of its entries will have been eliminated including any contribution from those squares. Thus, P_j could not possibly have contained the variable z_k , so $\mathcal{S}$ is an allowable polynomial transformation with respect to the variable z_k .

Conversely, suppose that there exists an allowable polynomial transformation on P_j with respect to the variable z_k , and let $\mathcal{T}$ be this polynomial transformation, which we shall express as follows:

$$\tilde{z}_i = z_i + \gamma^i, \quad (7.5)$$

for $i = 1, \dots, n+1$, where some of the γ^i may be zero. We note here that the transformation $\mathcal{T}$ is a Λ_j -homogeneous transformation, and so γ^i has the same weight with respect to Λ_j as z_i . Furthermore, we note that any Λ_j -homogeneous transformation can be written in this form.

We will prove that the k th row R_k is dependent by showing that it satisfies the condition given in (7.1). Assume that the variable z_k is contained in γ^i for some $i \in \{1, \dots, d\}$ with $d \leq n$. We know that each $\tilde{z}_i$ corresponds to the row and column operations:

$$R_k - \gamma_k^i R_i \rightarrow R_k \quad \text{and} \quad C_k - \tilde{\gamma}_k^i C_i \rightarrow C_k, \quad (7.6)$$

respectively, for $i = 1, \dots, d$, where $\gamma_k^i = \partial_{z_k} \gamma^i \neq 0$. Let $\tilde{P}_j$ be the leading polynomial P_j after the polynomial transformation $\mathcal{T}$ is applied to it. Since $\tilde{P}_j$ does not contain the variable z_k , the entries $\tilde{h}_{k\bar{i}}$ of R_k and $\tilde{h}_{i\bar{k}}$ of C_k of the matrix $A_{\tilde{P}_j}$ for all $t = 1, \dots, n$ are zero entries.

Now, by simply reversing the signs involved in the elementary operations in (7.6), we can restore R_k and C_k to their previous forms before the transformation $\mathcal{T}$ was applied to P_j . Hence, by performing the elementary row and column operations

$$R_k + \gamma_k^i R_i \rightarrow R_k \quad \text{and} \quad C_k + \tilde{\gamma}_k^i C_i \rightarrow C_k$$

for all $i = 1, \dots, d$ on the matrix $A_{\tilde{P}_j}$, the entries in R_k and C_k become

$$h_{k\bar{i}} = \sum_{i=1}^d \gamma_k^i h_{i\bar{i}} \quad \text{and} \quad h_{i\bar{k}} = \sum_{i=1}^d \tilde{\gamma}_k^i h_{i\bar{i}}, \quad (7.7)$$

where $h_{i\bar{i}}$ and $h_{i\bar{i}}$ are the entries in the i th row R_i and i th column C_i , respectively. Finally, we obtain that

$$h_{k\bar{i}} = \sum_{i=1}^n \gamma_k^i h_{i\bar{i}} \quad \text{and} \quad h_{i\bar{k}} = \sum_{i=1}^n \tilde{\gamma}_k^i h_{i\bar{i}}, \quad (7.8)$$

where $\gamma_k^i = 0$ for all $i = d + 1, \dots, n$. Thus, both R_k and C_k are dependent, proving the statement. $\square$

It is important to note that for any k , if the diagonal $(k, \bar{k})$ entry of A_{P_j} is the only nonzero entry in its k th row, then the k th row cannot be dependent, where A_{P_j} is the Levi matrix of the leading polynomial P_j . This statement holds because we are working with a sum of squares domain and A_{P_j} is Hermitian.

Lemma 7.2 *Let Γ be the set of $n \times n$ matrices with coefficients in the ring $\mathbb{C}[z, \bar{z}]$. Let $H \in \Gamma$ be Hermitian. For some given i and k , let B be the matrix obtained from H after the elementary row and column operations $R_k + \alpha R_i \rightarrow R_k$ and $C_k + \bar{\alpha} C_i \rightarrow C_k$, for some $\alpha \in \mathbb{C}[z]$, are performed on it. Then $\det(B) = \det(H)$.*

Proof Let E be the matrix obtained from H by the elementary row operation $R_k + \alpha R_i \rightarrow R_k$. Then B is the matrix obtained from E by the elementary column operation $C_k + \bar{\alpha} C_i \rightarrow C_k$. It is obvious from the properties of the determinant that $\det(H) = \det(E)$ and that $\det(E) = \det(B)$. $\square$

Lemma 7.3 *Assume that the D'Angelo 1-type of $M \subset \mathbb{C}^{n+1}$ at 0 is finite. At step j of the Kolář algorithm for the computation of the multitype at 0, let P_j be the leading polynomial and let Q_j be the leftover polynomial.*

If the determinant of A_{P_j} is nonzero, then P_j is independent of the largest number of variables, and no polynomial transformation needs to be performed on it before the next step in the Kolář algorithm.

Proof We shall give a proof of the contrapositive of the statement of this lemma, which states that if there exists an allowable transformation, and hence, one of the rows of A_{P_j} is dependent by Proposition 1.3, then the determinant of A_{P_j} is zero. Suppose that $\mathcal{R} = \{R_1, \dots, R_n\}$, the set of all rows of the matrix A_{P_j} , is dependent and that none of the rows is identically equal to zero. Thus, for some k , we can write

$$R_k = \sum_{\substack{l=1 \\ l \neq k}}^n \alpha_l R_l,$$

where $\alpha_l \in \mathbb{C}[z]$ and R_l is the l th row of A_{P_j} . From the proof of Proposition 1.3, we know that there must exist some elementary row and column operations that transform A_{P_j} into the matrix $\tilde{A}_{P_j}$ of which k th row and column have all zero entries. Since the matrix $\tilde{A}_{P_j}$ has at least one row with all entries equal to zero, its determinant is equal to zero. From Lemma 7.2, we know that $\det(\tilde{A}_{P_j}) = \det(A_{P_j})$.

Thus, $\det(A_{P_j}) = 0$, which is the result we need. $\square$

Given the way the leading polynomial P_j and its Levi matrix A_{P_j} are constructed, it is possible that P_j could be independent of at least one of the variables. If that is the case, then the determinant of the Levi matrix corresponding to the leading polynomial P_j will always be zero since it will have at least one row that is identically zero. Therefore, we need a way to determine when a subset of all nonzero rows of A_{P_j} is independent. To address this situation, we shall consider the following:

Let $A_{P_j} = (a_{i\bar{l}})_{1 \leq i, l \leq n}$ be the Levi matrix of the leading polynomial P_j at step j of the Kolář algorithm. Let m be the number of nonzero rows of the matrix A_{P_j} . Denote by $A_{P_j|m}$ the principal submatrix obtained from A_{P_j} by removing all zero rows and columns to get precisely m rows and columns, for some $m \leq n$. Put differently, $A_{P_j|m}$ is the submatrix consisting only of all nonzero rows and columns of A_{P_j} . If $m = n$, then none of the rows and columns are identically zero.

Also, via the elementary row and column operations, $A_{P_j|m}$ can be transformed into a leading principal submatrix of A_{P_j} , where the first m rows and columns are the ones that remain. In this case, $A_{P_j|m} = (a_{i\bar{l}})_{1 \leq i, l \leq m}$.

Let $\mathcal{R} = \{R_1, \dots, R_n\}$ be the set of all rows of the matrix A_{P_j} and let $\mathcal{S} = \{R_{b_1}, \dots, R_{b_m}\}$ be a subset of $\mathcal{R}$, for $b_e \in \{1, \dots, n\}$, $e = 1, \dots, m$ such that each element R_{b_e} is not identically zero. Then $\mathcal{S}$ is the set of all non zero rows of the submatrix $A_{P_j|m}$.

We now restate Proposition 1.3 and Lemma 7.3 as follows:

Proposition 7.4 *Assume that the D'Angelo 1-type of $M \subset \mathbb{C}^{n+1}$ at 0 is finite. At step j of the Kolář algorithm for the computation of the multitype at 0, let P_j be the leading polynomial, and let Q_j be the leftover polynomial.*

There exists an allowable polynomial transformation on P_j with respect to the variable z_k via the elementary row and column operations if and only if the k th row of $A_{P_j|m}$ is dependent.

Lemma 7.5 *Assume that the D'Angelo 1-type of $M \subset \mathbb{C}^{n+1}$ at 0 is finite. At step j of the Kolář algorithm for the computation of the multitype at 0, let P_j be the leading polynomial, and let Q_j be the leftover polynomial.*

If the determinant of $A_{P_j|m}$ is nonzero, then P_j is independent of the largest number of variables, and no polynomial transformation needs to be performed on it before the next step in the Kolář algorithm.

The proofs of Proposition 7.4 and Lemma 7.5 are identical to the proofs given for Proposition 1.3 and Lemma 7.3, respectively, since the latter do not depend on rows being identically equal to zero. We also note here that the converses of Lemmas 7.3 and 7.5 do not hold. The reason is that the notion of dependency given in (7.1) is more restrictive than the standard notion of dependency in linear algebra, so there might not exist a row that is dependent according to our definition, but the set of rows may satisfy the standard notion of dependency, in which case the determinant of the Levi matrix would be identically equal to zero.

Now, the natural question to ask at this point is this: Given a Levi matrix of a leading polynomial with zero determinant, how can we tell whether or not it has dependent rows? Also, if there exist dependent rows, how can we identify such rows in order to determine the allowable polynomial transformations corresponding to these dependent rows? The answers to these questions lie in the formulation of an algorithm, which we will describe in the next subsection.

7.1 Jacobian Modules and Gradient Ideals

A study of the elementary row and column operations on the Levi matrix reveals that a row (column) operation on the Levi matrix is performed by a multiplication on the left (right) of the Levi matrix by an elementary row (column) matrix. The Levi matrix of a sum of squares domain can always be decomposed as the product of the complex Jacobian matrix of the holomorphic functions that generate the domain and its conjugate transpose. Therefore, every row operation on the Levi matrix could be performed on the complex Jacobian matrix, while every column operation on the Levi matrix is performed on the conjugate transpose of the complex Jacobian matrix. Let A be an $n \times n$ Levi matrix of a domain given by the sum of squares of N holomorphic functions. Then elementary matrices are $n \times n$ matrices, while the complex Jacobian matrix and its conjugate transpose must be $n \times N$ and $N \times n$ matrices, respectively.

In our study of elementary row and column operations performed on the Levi determinant of a sum of squares domain, one particular property of the Levi matrix of a square of one of the generators drew our attention: All entries of any given column (row) have the same anti-holomorphic (holomorphic) parts, and so a study of the relationship between these entries narrows down to a study of the relationship between their holomorphic (anti-holomorphic) parts. In other words, we expect that the study of the Levi matrix will be much easier if we transition from the sum of squares to the underlying ideal of holomorphic functions that generate the domain as we already saw was the case for the computation of the multitype.

We start with a couple of definitions that we specialize to complex polynomials since those are the objects that appear in the Kolář algorithm when it is applied to a sum of squares domain instead of the full holomorphic generators:

Definition 7.1 Let $f \in \mathbb{C}[z_1, \dots, z_n]$ be polynomial in the variables $z_1, \dots, z_n$ with coefficients in $\mathbb{C}$. As in [15], we define the *gradient ideal of f* as the ideal generated by the partial derivatives of f :

$$\mathcal{I}_{\text{grad}}(f) = \langle \nabla f \rangle = \left\langle \frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_n} \right\rangle. \quad (7.9)$$

Definition 7.2 Given the ideal $\langle f \rangle = \langle f_1, \dots, f_N \rangle \subset \mathbb{C}[z_1, \dots, z_n]$, we define the *Jacobian module of f* as

$$\mathfrak{J}_{\langle f \rangle} = \left[\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_n} \right], \quad (7.10)$$

where each $\frac{\partial f}{\partial z_j}$ is a vector. $\mathfrak{J}_{\langle f \rangle}$ is a module over the polynomial ring $\mathbb{C}[z_1, \dots, z_n]$.

To every Jacobian module $\mathfrak{J}_{\langle f \rangle}$, we associate the complex Jacobian matrix $J(f)$ given by

$$J(f) = \begin{pmatrix} \frac{\partial f_1}{\partial z_1} & \frac{\partial f_2}{\partial z_1} & \cdots & \frac{\partial f_N}{\partial z_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_1}{\partial z_n} & \frac{\partial f_2}{\partial z_n} & \cdots & \frac{\partial f_N}{\partial z_n} \end{pmatrix}. \quad (7.11)$$

Likewise, to each gradient ideal $\mathcal{I}_{\text{grad}}(f_i) = \langle \frac{\partial f_i}{\partial z_1}, \dots, \frac{\partial f_i}{\partial z_n} \rangle$ of the generator $f_i \in \mathbb{C}[z_1, \dots, z_n]$ of $\langle f \rangle$, we associate the i th column of $J(f)$ for $1 \leq i \leq n$. The reader should note that row operations on $J(f)$ are precisely operations on the module $\mathfrak{J}_{\langle f \rangle}$.

Now, let $\langle f \rangle = \langle f_1, \dots, f_N \rangle \subset \mathbb{C}[z_1, \dots, z_n]$ be the leading polynomial ideal at some step of the Kolář algorithm. Then we are particularly interested in simplifying the Jacobian module $\mathfrak{J}_{\langle f \rangle}$ such that it is generated by the minimal number of generators. Every generator that is eliminated is a linear combination of some partial derivatives of f with coefficients in $\mathbb{C}[z_1, \dots, z_n]$. Since every generator of the Jacobian module represents a row of the complex Jacobian matrix, every eliminated generator represents a dependent row in the complex Jacobian matrix. Owing to this connection, from every eliminated generator, we can construct a sequence of elementary row operations that corresponds to the linear combination of some partial derivatives of f as described in Proposition 1.3. Hence, we obtain polynomial transformations corresponding to these row operations.

It is easy to observe this relationship if $N = 1$. Then reducing the number of generators of $\mathfrak{J}_{\langle f \rangle}$, if possible, reduces the number of nonzero rows of the associated complex Jacobian matrix. Thus, the minimal number of generators required to generate the Jacobian module is precisely the number of independent rows of the complex Jacobian matrix, which is the same as the number of variables on which the corresponding leading polynomial ideal $\langle f \rangle$ is dependent by Proposition 1.3 after the corresponding change of variables. Hence, the number d_j at step j of the Kolář algorithm is given by $d_j = n - \#f$, where $\#f$ is the minimal number of generators generating the Jacobian module $\mathfrak{J}_{\langle f \rangle}$, n is the number of variables in the polynomial ring $\mathbb{C}[z_1, \dots, z_n]$, and d_j is the largest number of variables of which the leading polynomial at step j is independent. This gives an algebraic characterization of the number d_j in the Kolář algorithm.

In the more general case where $N > 1$, reducing the number of generators of the Jacobian module implies reducing the generators of all gradient ideals by the same operations. Thus, $\frac{\partial f}{\partial z_\ell}$, for some ℓ , is a generator that is eliminated in the Jacobian module $\mathfrak{J}_{\langle f \rangle}$ if and only if the generator $\frac{\partial f_i}{\partial z_\ell}$ of the gradient ideal $\mathcal{I}_{\text{grad}}(f_i)$ for all $i = 1, \dots, N$ is eliminated, namely reduced to 0.

Clearly, if there exists at least one gradient ideal $\mathcal{I}_{\text{grad}}(f_i)$ with minimal number of generators equal to n , then the Jacobian module $\mathfrak{J}_{\langle f \rangle}$ cannot have fewer than n generators, i.e., no reduction via row operations is possible.

In this setup, all elementary pairs of row-column operations on the Levi matrix reduce to just elementary row operations on the complex Jacobian matrix.

7.2 Row Reduction Algorithm

We shall now devise an algorithm that constructs explicitly the polynomial transformations required at each step of the Kolář algorithm when applied to the complex Jacobian matrix of the leading polynomial ideal.

The algorithm gives the conditions for characterizing the required elementary row operations that correspond to the polynomial transformations needed in the Kolář algorithm. The application of the algorithm to the complex Jacobian matrix corresponding to a given leading polynomial ideal will eliminate all dependent rows, if they exist, from the complex Jacobian matrix.

Now, for any $j \geq 1$, let J_{P_j} be the complex Jacobian matrix corresponding to the leading polynomial P_j at step j of the Kolář algorithm.

Let $P_j = \sum_{i=1}^N |h_i|^2$. Then the leading polynomial ideal is given by $\mathcal{I}_{P_j} := \langle h \rangle = \langle h_1, \dots, h_N \rangle$, and the gradient ideal of h_i is $\mathcal{I}_{\text{grad}}(h_i) = \left\langle \frac{\partial h_i}{\partial z_1}, \dots, \frac{\partial h_i}{\partial z_n} \right\rangle$, $i = 1, \dots, N$. Note that the complex Jacobian matrix of P_j is given by $J_{P_j} = J(\mathcal{I}_{P_j}) = J(h)$ and the Levi matrix A_{P_j} of P_j is the product of the complex Jacobian matrix of P_j with its conjugate transpose $J^*(h) : A_{P_j} = J(h)J^*(h)$. We shall reduce, if possible, the number of generators of each gradient ideal one at a time and control the changes that occur in other gradient ideals as a result of these reduction operations. By control, we mean setting appropriate conditions on the reduction operations used.

If $\det A_{P_j} = \det (J(h)J^*(h))$ is nonzero, then no change of variables is required by Lemma 7.3. Thus, assume that $\det (J(h)J^*(h)) = 0$. Then:

1. We begin the process by first considering the gradient ideal $\mathcal{I}_{\text{grad}}(h_i) = \left\langle \frac{\partial h_i}{\partial z_1}, \dots, \frac{\partial h_i}{\partial z_n} \right\rangle$ for any i . Simplify the gradient ideal $\mathcal{I}_{\text{grad}}(h_i)$ such that it consists of the minimal number of generators. Assume that

$$\frac{\partial h_i}{\partial z_k} = \sum_{u=1}^v \gamma_{c_u} \frac{\partial h_i}{\partial z_{c_u}}, \quad (7.12)$$

for some k , $c_u \neq k$, $v < n$, and γ_{c_u} a nonzero polynomial in $\mathbb{C}[z_1, \dots, z_n]$. Then perform the following elementary row operations on the complex Jacobian matrix $J(h)$:

$$R_\ell - \frac{\partial \zeta_{c_u}}{\partial z_\ell} R_{c_u} \rightarrow R_\ell, \quad (7.13)$$

for all $u = 1, \dots, v$ and for all variables z_ℓ in $\zeta_{c_u} = \int_0^{z_k} \gamma_{c_u}(t) dt$. By Lemma 7.1, the row operations in (7.13) correspond to the polynomial transformation given by

$$\tilde{z}_{c_u} = z_{c_u} + \int_0^{z_k} \gamma_{c_u}(t) dt; \quad \tilde{z}_\omega = z_\omega, \quad (7.14)$$

for all $\omega \neq c_u$ and for all $u = 1, \dots, v$. The generator $\frac{\partial h_i}{\partial z_k}$ vanishes in $\mathcal{I}_{\text{grad}}(h_i)$ after the row operations in (7.13) are performed on $J(h)$.

We say row R_{c_u} is used as a *central row* in the sequence of row operations and the generator $\frac{\partial h_i}{\partial z_{c_u}}$ is used as a *central generator* in the simplification of the gradient ideal $\mathcal{I}_{\text{grad}}(h_i)$ for all i . We remark here that for all subsequent row operations performed on the complex Jacobian matrix, the row R_{c_u} cannot be used as a central row and $\frac{\partial h_e}{\partial z_{c_u}}$ cannot be used as a central generator in the simplification of any other gradient ideal $\mathcal{I}_{\text{grad}}(h_e)$ for $e \neq i$. This condition is imposed due to Proposition 1.3, which is an equivalence. Reusing a central row or a central generator might reintroduce a variable that has been eliminated from the leading polynomial.

- Next, consider another gradient ideal $\mathcal{I}_{\text{grad}}(h_s) = \left\langle \frac{\partial h_s}{\partial z_1}, \dots, \frac{\partial h_s}{\partial z_n} \right\rangle$ for $s \neq i$. Clearly, the k th generator of this gradient ideal is

$$\frac{\partial h_s}{\partial z_k} - \sum_{u=1}^v \gamma_{c_u} \frac{\partial h_s}{\partial z_{c_u}} \quad (7.15)$$

due to the row operations given in (7.13). We simplify the ideal $\mathcal{I}_{\text{grad}}(h_s)$ such that it has the minimal number of generators while ensuring that the generators $\frac{\partial h_s}{\partial z_{c_u}}$ for $u = 1, \dots, v$ are not used as central generators in the simplification of $\mathcal{I}_{\text{grad}}(h_s)$. Perform the related row operations.

- Proceed similarly by considering other gradient ideals different from the previous ones. Since there are only finitely many gradient ideals and finitely many generators that generate each of them, the process will terminate after a finite number of steps.

We will show in the lemma that follows that the polynomial transformation in (7.14) corresponding to the row operations given in (7.13) is Λ_j -homogeneous. Thus, we state the following:

Lemma 7.6 *Assume that the D'Angelo 1-type of $M \subset \mathbb{C}^{n+1}$ at 0 is finite. At step j of the Kolář algorithm for the computation of the multitype at 0, let Λ_j be the weight, P_j the leading polynomial, and $\mathcal{I}_{P_j}$ the corresponding leading polynomial ideal. Let $\mathcal{I}_{\text{grad}}(\psi)$ be the gradient ideal of some generator ψ of the ideal $\mathcal{I}_{P_j}$. Assume that*

$$\frac{\partial \psi}{\partial z_k} = \sum_{u=1}^v \gamma_{c_u} \frac{\partial \psi}{\partial z_{c_u}}, \quad (7.16)$$

for some k , where $k \neq c_u$, $v < n$, and γ_{c_u} is a nonzero polynomial in $\mathbb{C}[z_1, \dots, z_n]$. Let $\zeta_{c_u} = \int_0^{z_k} \gamma_{c_u}(t) dt$.

Then the polynomial transformation given by $\tilde{z}_{c_u} = z_{c_u} + \zeta_{c_u}$; $\tilde{z}_\omega = z_\omega$ for all $\omega \neq c_u$ corresponding to the elementary row operations $R_\ell - \frac{\partial \zeta_{c_u}}{\partial z_\ell} R_{c_u} \rightarrow R_\ell$ for all variables z_ℓ in ζ_{c_u} and for all $u = 1, \dots, v$ performed on the complex Jacobian matrix J_{P_j} is Λ_j -homogeneous for all $u = 1, \dots, v$.

Proof We start the proof by recalling from Theorem 1.1 that the leading polynomial is a sum of squares at every step of the Kolář algorithm. Hence, P_j is a sum of squares. Let $\Lambda_j = (\lambda_1, \dots, \lambda_n)$. Since variables are not ordered in increasing weight order, we let $\phi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ be the bijection $\phi = (\phi_1, \dots, \phi_n)$ such that the variable z_i , $1 \leq i \leq n$, has weight λ_{ϕ_i} .

We will show that the term γ_{c_u} in the polynomial transformation is of weighted degree $(\lambda_{\phi_{c_u}} - \lambda_{\phi_k})$. Let ν be the weighted degree of γ_{c_u} with respect to the weight Λ_j . By our hypothesis, the weighted degrees of $\frac{\partial \psi}{\partial z_{c_u}}$ and $\frac{\partial \psi}{\partial z_k}$ are $\frac{1}{2} - \lambda_{\phi_{c_u}}$ and $\frac{1}{2} - \lambda_{\phi_k}$, respectively, since all generators of the leading polynomial ideal $\mathcal{I}_{P_j}$ are of weighted degree $\frac{1}{2}$ with respect to Λ_j . The weighted degree of the right hand side of the expression given in (7.16) is $\nu + \frac{1}{2} - \lambda_{\phi_{c_u}}$. Hence solving the equation in (7.16) for ν gives $\nu = \lambda_{\phi_{c_u}} - \lambda_{\phi_k}$. Thus, the weighted degree of $\zeta_{c_u} = \int_0^{z_k} \gamma_{c_u}(t) dt$ is $\lambda_{\phi_{c_u}}$ as required. $\square$

Remark 7.6.1 The polynomial γ_{c_u} cannot depend on the variable z_{c_u} because if it were to depend on z_{c_u} , then its weighted degree would satisfy $\nu \geq \lambda_{\phi_{c_u}}$, but $\nu = \lambda_{\phi_{c_u}} - \lambda_{\phi_k}$, which gives a contradiction because $\lambda_{\phi_k} > 0$.

Lemma 7.7 Assume that the D'Angelo 1-type of M in $\mathbb{C}^{n+1}$ at 0 is finite. At step j of the Kolář algorithm for the computation of the multitype at 0, let P_j be the leading polynomial, and let $\mathcal{I}_{P_j} = \langle h \rangle \subset \mathbb{C}[z_1, \dots, z_n]$ be the corresponding leading polynomial ideal.

If the Row Reduction algorithm is applied to the complex Jacobian matrix $J(h)$, then every dependent row of $J(h)$ vanishes. In other words, the leading polynomial ideal $\mathcal{I}_{P_j}$ is independent of the largest number of variables after the Row Reduction algorithm is applied to $J(h)$.

Proof From Theorem 1.1 the leading polynomial P_j is a sum of squares, and so let $P_j = \sum_{i=1}^N |h_i|^2$. Then the leading polynomial ideal $\mathcal{I}_{P_j}$ is $\langle h \rangle = \langle h_1, \dots, h_N \rangle$, and let the gradient ideal of each generator h_i be $\mathcal{I}_{\text{grad}}(h_i) = \langle \frac{\partial h_i}{\partial z_1}, \dots, \frac{\partial h_i}{\partial z_n} \rangle$.

Now, assume that R_k , the k th row of $J(h)$ is dependent. We will show that the generator $\frac{\partial h}{\partial z_k}$ of the Jacobian module given by $\mathfrak{J}(h) = \left[\frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_n} \right]$ vanishes after applying the Row Reduction algorithm on the complex Jacobian matrix $J(h)$. Since R_k is dependent, we can write the generator $\frac{\partial h}{\partial z_k}$ as

$$\frac{\partial h}{\partial z_k} = \sum_{u=1}^v \gamma_{c_u} \frac{\partial h}{\partial z_{c_u}}, \quad (7.17)$$

for some k , $c_u \neq k$, $v < n$, and γ_{c_u} a nonzero polynomial in $\mathbb{C}[z_1, \dots, z_n]$ for every u . Hence, every generator $\frac{\partial h_i}{\partial z_k}$ of the gradient ideal $\mathcal{I}_{\text{grad}}(h_i)$ can be written as

$$\frac{\partial h_i}{\partial z_k} = \sum_{u=1}^v \gamma_{c_u} \frac{\partial h_i}{\partial z_{c_u}}, \quad (7.18)$$

for $i = 1, \dots, N$ and the same polynomial coefficients γ_{cu} . Thus, it suffices to show that $\frac{\partial h_i}{\partial z_k}$ vanishes at the termination of the algorithm for every $i = 1, \dots, N$. Consider the ideal $\mathcal{I}_{\text{grad}}(h_i)$ for some $i \in \{1, \dots, N\}$. If the generator $\frac{\partial h_i}{\partial z_k}$ is zero, then there is nothing to be done, and so we move to a different gradient ideal. Hence, suppose that $\frac{\partial h_i}{\partial z_k}$ is nonzero. Then at least one of the generators $\frac{\partial h_i}{\partial z_{cu}}$ is nonzero for some u . Suppose that $\frac{\partial h_i}{\partial z_{cu}} \neq 0$ for all $u \in \{1, \dots, w\}$ for $w \leq v$. Since it satisfies the condition in (7.18), we perform the elementary row operations $R_\ell - \frac{\partial \zeta_{cu}}{\partial z_\ell} R_{cu} \rightarrow R_\ell$, for all variables z_ℓ in $\zeta_{cu} = \int_0^{z_k} \gamma_{cu}(t) dt$ and for all $u = 1, \dots, w$. This eliminates the term $\sum_{u=1}^w \gamma_{cu} \frac{\partial h_i}{\partial z_{cu}}$, from the ideal $\mathcal{I}_{\text{grad}}(h_i)$. The generator $\frac{\partial h_i}{\partial z_k}$ of the Jacobian module becomes

$$\frac{\partial h_i}{\partial z_k} = \sum_{u=w+1}^v \gamma_{cu} \frac{\partial h_i}{\partial z_{cu}} \quad (7.19)$$

after the row operations have been performed on $J(h)$. Note here that the generator $\frac{\partial h_e}{\partial z_{cu}}$, for all $e \neq i$ and $u = 1, \dots, w$ cannot be central in any simplification process in the gradient ideal $\mathcal{I}_{\text{grad}}(h_e)$ after the row operations.

Next, consider another gradient ideal $\mathcal{I}_{\text{grad}}(h_e)$ for $e \neq i$. Then its k th generator after the reduction operation is

$$\frac{\partial h_e}{\partial z_k} - \sum_{u=1}^w \gamma_{cu} \frac{\partial h_e}{\partial z_{cu}} = \sum_{u=w+1}^v \gamma_{cu} \frac{\partial h_e}{\partial z_{cu}}. \quad (7.20)$$

If the expression in (7.20) equals zero, then there is nothing left to be done. If the expression in (7.20) is nonzero, then $\frac{\partial h_e}{\partial z_{cu}} \neq 0$ for some $u = w+1, \dots, q$ with $q \leq v$. Perform the elementary row operations $R_\ell - \frac{\partial \zeta_{cu}}{\partial z_\ell} R_{cu} \rightarrow R_\ell$, for all variables z_ℓ in $\zeta_{cu} = \int_0^{z_k} \gamma_{cu}(t) dt$ and for all $u = w+1, \dots, q$ to eliminate the term $\sum_{u=w+1}^q \gamma_{cu} \frac{\partial h_e}{\partial z_{cu}}$, where $q \leq v$. We follow this process through in each of the distinct gradient ideals until all gradient ideals have been considered. The expression in (7.19) becomes zero at some point; otherwise, we get a contradiction to R_k being dependent. $\square$

Example 3 Let the hypersurface $M \subseteq \mathbb{C}^5$ be given by $r = 0$ with

$$r = 2\text{Re}(z_5) + |(z_1 + z_2 z_4)^2 + z_2^4|^2 + |(z_1 + z_2 z_3^2)^2|^2 + |z_2^9|^2 + |z_3^{10}|^2 + |z_4^{12}|^2.$$

Let $B = 2(z_1 + z_2 z_4)$, $C = 2(z_1 + z_2 z_3^2)$, and let the ideal associated to the domain M be $\langle h \rangle = \langle (z_1 + z_2 z_4)^2 + z_2^4, (z_1 + z_2 z_3^2)^2 + z_2^9, z_3^{10}, z_4^{12} \rangle$. The complex Jacobian matrix is given by

$$J(h) = \begin{pmatrix} B & C & 0 & 0 & 0 \\ z_4 B + 4z_2^3 & z_3^2 C & 9z_2^8 & 0 & 0 \\ 0 & 2z_2 z_3 C & 0 & 10z_3^9 & 0 \\ z_2 B & 0 & 0 & 0 & 12z_4^{11} \end{pmatrix}.$$

The Bloom-Graham type is 4, $\mathcal{I}_{P_1} = \langle z_1^2, z_1^2 \rangle$, and $\Lambda_1 = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$. The maximum $W_1 = \frac{1}{8}$, the leading polynomial ideal $\mathcal{I}_{P_2} = \langle (z_1 + z_2 z_4)^2 + z_2^4, z_1^2 \rangle$, and $\Lambda_2 = (\frac{1}{4}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8})$. The maximum $W_2 = \frac{1}{16}$, $\mathcal{I}_{P_3} = \langle (z_1 + z_2 z_4)^2 + z_2^4, (z_1 + z_2 z_3^2)^2 \rangle$, and $\Lambda_3 = (\frac{1}{4}, \frac{1}{8}, \frac{1}{8}, \frac{1}{16})$. The complex Jacobian matrix corresponding to $\mathcal{I}_{P_3}$ is given by

$$J(h_1, h_2) = \begin{pmatrix} B & C \\ z_4 B + 4z_2^3 & z_2^3 C \\ 0 & 2z_2 z_3 C \\ z_2 B & 0 \end{pmatrix}.$$

Consider the gradient ideal $\mathcal{I}_{\text{grad}}(h_1) = \langle B, z_4 B + 4z_2^3, 0, z_2 B \rangle$. Since $\frac{\partial h_1}{\partial z_4} = z_2 \frac{\partial h_1}{\partial z_1}$, its simplification is $\mathcal{I}_{\text{grad}}(h_1) = \langle B, 4z_2^3 \rangle$. We perform the elementary row operations $R_2 - z_4 R_1 \rightarrow R_2$ and $R_4 - z_2 R_1 \rightarrow R_4$ on the matrix $J(h)$. The matrices above become

$$J(h) = \begin{pmatrix} B & C & 0 & 0 & 0 \\ 4z_2^3 & (z_3^2 - z_4)C & 9z_2^8 & 0 & 0 \\ 0 & 2z_2 z_3 C & 0 & 10z_3^9 & 0 \\ 0 & -z_2 C & 0 & 0 & 12z_4^{11} \end{pmatrix} \quad \text{and} \quad J(h_1, h_2) = \begin{pmatrix} B & C \\ 4z_2^3 & (z_3^2 - z_4)C \\ 0 & 2z_2 z_3 C \\ 0 & -z_2 C \end{pmatrix}.$$

These operations correspond to the polynomial transformation $\tilde{z}_1 = z_1 + z_2 z_4 : \tilde{z}_\omega = z_\omega$ for $\omega \neq 1$, and now $B = 2\tilde{z}_1$ and $C = 2(\tilde{z}_1 + \tilde{z}_2(\tilde{z}_3^2 - \tilde{z}_4))$. The generator $\frac{\partial h_2}{\partial z_1}$ cannot be a central generator in the simplification of $\mathcal{I}_{\text{grad}}(h_2)$, and row 1 cannot be used as a central row in any subsequent elementary row operations. Consider the next gradient ideal $\mathcal{I}_{\text{grad}}(h_2) = \langle C, (\tilde{z}_3^2 - \tilde{z}_4)C, 2\tilde{z}_2 \tilde{z}_3 C, -\tilde{z}_2 C \rangle$. Here the generator $-\tilde{z}_2 C$ in the fourth component is the only central generator in $\mathcal{I}_{\text{grad}}(h_2)$. Thus, $\frac{\partial h_2}{\partial z_3} = -2\tilde{z}_3 \frac{\partial h_2}{\partial z_4}$, and its simplification is $\langle C, (\tilde{z}_3^2 - \tilde{z}_4)C, -\tilde{z}_2 C \rangle$.

We perform the elementary row operations $R_3 + 2z_3 R_4 \rightarrow R_3$ on the matrix $J(h)$. The matrices become

$$J(h) = \begin{pmatrix} B & C & 0 & 0 & 0 \\ 4z_2^3 & (z_3^2 - z_4)C & 9z_2^8 & 0 & 0 \\ 0 & 0 & 0 & 10z_3^9 & 24z_3 z_4^{11} \\ 0 & -z_2 C & 0 & 0 & 12z_4^{11} \end{pmatrix} \quad \text{and} \quad J(h_1, h_2) = \begin{pmatrix} B & C \\ 4z_2^3 & (z_3^2 - z_4)C \\ 0 & 0 \\ 0 & -z_2 C \end{pmatrix}.$$

This operation corresponds to the polynomial transformation $\dot{z}_4 = \tilde{z}_4 - \tilde{z}_3^2$; $\dot{z}_\omega = \tilde{z}_\omega$ for $\omega \neq 4$, and now $B = 2\dot{z}_1$ and $C = 2(\dot{z}_1 - \dot{z}_2 \dot{z}_4)$. The leading polynomial ideal $\mathcal{I}_{P_3} = \langle \dot{z}_1^2 + \dot{z}_2^4, (\dot{z}_1 - \dot{z}_2 \dot{z}_4)^2 \rangle$. The maximum $W_3 = \frac{1}{20}$, and $\Lambda_4 = (\frac{1}{4}, \frac{1}{8}, \frac{1}{8}, \frac{1}{20})$.

Acknowledgements Special thanks goes to the author's supervisor Andreea Nicoara for all her valuable remarks and guidance during our many discussions. The author would also like to thank Dmitri Zaitsev for his useful comments and Berit Stensønes for her encouragement.

Funding Open access funding provided by NTNU Norwegian University of Science and Technology (incl St. Olavs Hospital - Trondheim University Hospital)

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bloom, T., Graham, I.: On "type" conditions for generic real submanifolds of C^n . *Invent. Math.* **40**(3), 217–243 (1977)
2. Catlin, D.: Boundary invariants of pseudoconvex domains. *Ann. Math. (2)* **120**(3), 529–586 (1984)
3. Catlin, D., Cho, J.-S.: Sharp estimates for the $\bar{\partial}$ -Neumann problem on regular coordinate domains. Preprint (2008). [arXiv:08110830v1\[math.CV\]](https://arxiv.org/abs/08110830v1)
4. Catlin, D., D'Angelo, J.P.: Subelliptic estimates. In *Complex Analysis. Trends in Mathematics*, pp. 75–94. Birkhäuser/Springer Basel AG, Basel (2010)
5. D'Angelo, J.P.: Real hypersurfaces, orders of contact, and applications. *Ann. Math. (2)* **115**(3), 615–637 (1982)
6. D'Angelo, J.P.: Several complex variables and the geometry of real hypersurfaces. In: *Studies in Advanced Mathematics*. CRC Press, Boca Raton (1993)
7. Fassina, M.: Singularities and Multiplier Algorithms for Real Hypersurfaces. PhD Thesis, Illinois University, Urbana (2020)
8. Khanh, T.V., Zampieri, G.: Precise subelliptic estimates for a class of special domains. *J. Anal. Math.* **123**, 171–181 (2014)
9. Kim, S.-Y., Zaitsev, D.: Jet vanishing orders and effectivity of Kohn's algorithm in dimension 3. *Asian J. Math.* **22**(3), 545–568 (2018)
10. Kim, S.-Y., Zaitsev, D.: Triangular resolutions and effectiveness for holomorphic subelliptic multipliers. *Adv. Math.* **387**, 1–36 (2021). [arXiv:2003.06482](https://arxiv.org/abs/2003.06482)
11. Kohn, J.J.: Boundary behavior of $\bar{\partial}$ on weakly pseudo-convex manifolds of dimension two. *J. Differ. Geom.* **6**, 523–542. Collection of articles dedicated to S. S. Chern and D. C. Spencer on their sixtieth birthdays (1972)
12. Kohn, J.J.: Subellipticity of the $\bar{\partial}$ -Neumann problem on pseudo-convex domains: sufficient conditions. *Acta Math.* **142**(1–2), 79–122 (1979)
13. Kolář, M.: The Catlin multitype and biholomorphic equivalence of models. *Int. Math. Res. Not. (IMRN)* **18**, 3530–3548 (2010)
14. Kolář, M., Meylan, F., Zaitsev, D.: Chern–Moser operators and polynomial models in CR geometry. *Adv. Math.* **263**, 321–356 (2014)
15. Nie, J., Demmel, J., Sturmfels, B.: Minimizing polynomials via sum of squares over the gradient ideal. *Math. Program.* **106**(3, Ser. A), 587–606 (2006)
16. Siu, Y.-T.: Effective termination of Kohn's algorithm for subelliptic multipliers. *Pure Appl. Math. Q.* **6**(4, Special Issue: In honor of Joseph J. Kohn. Part 2), 1169–1241 (2010)
17. Siu, Y.-T.: New procedure to generate multipliers in complex Neumann problem and effective Kohn algorithm. *Sci. China Math.* **60**(6), 1101–1128 (2017)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.