



# Global vs Blow-Up Solutions and Optimal Threshold for Hyperbolic ODEs with Possibly Singular Nonlinearities

Daniele Cassani<sup>1,2</sup> · Tosiya Miyasita<sup>3</sup>

Received: 15 August 2023 / Accepted: 29 October 2023 / Published online: 7 December 2023  
© The Author(s) 2023

## Abstract

We consider a hyperbolic ordinary differential equation perturbed by a nonlinearity which can be singular at a point and in particular this includes MEMS type equations. We first study qualitative properties of the solution to the stationary problem. Then, for small value of the perturbation parameter as well as initial value, we establish the existence of a global solution by means of the Lyapunov function and we show that the omega limit set consists of a solution to the stationary problem. For strong perturbations or large initial values, we show that the solution blows up. Finally, we discuss the relationship between upper bounds of the perturbation parameter for the existence of time-dependent and stationary solutions, for which we establish an optimal threshold.

**Keywords** Lyapunov functions · Global solutions · Omega limit set · Blow-up · Dynamical threshold · MEMS

**Mathematics Subject Classification** 34A12 · 34G20 · 37L25

---

✉ Daniele Cassani  
daniele.cassani@uninsubria.it

Tosiya Miyasita  
miyasita.t@yamato-u.ac.jp

<sup>1</sup> Dipartimento di Scienza e Alta Tecnologia, Università degli Studi dell'Insubria, Varese, Italy

<sup>2</sup> RISM–Riemann International School of Mathematics, Villa Toeplitz, Via G.B. Vico, 46, 21100 Varese, Italy

<sup>3</sup> Division of Mathematical Science, Department of Science and Engineering, Faculty of Science and Engineering, Yamato University, 2-5-1, Katayama-cho, Suita-shi, Osaka 564-0082, Japan

### 1 Introduction

In this paper, we study the following ordinary differential equation:

$$\begin{cases} u_{tt} + \alpha f(u_t) + \beta u = \lambda g(u) \text{ for } t > 0, \\ u(0) = u_0, \\ u_t(0) = v_0, \end{cases} \tag{1}$$

where  $\lambda > 0, \alpha \geq 0, \beta > 0, u_0 \in \mathbb{R}$  and  $v_0 \in \mathbb{R}$ . Under appropriate assumptions on  $f$  and  $g$ , we discuss the global existence and blow-up of the solutions to (1). For  $\alpha \geq 0$  and the initial value  $(u_0, v_0) \in \mathbb{R}^2$ , we denote by  $\lambda^*(u_0, v_0)(\alpha)$  the dynamical threshold for the existence of a global solution of (1). Namely, the solution of (1) exists globally in time for  $0 < \lambda < \lambda^*(u_0, v_0)(\alpha)$  and blows up for  $\lambda > \lambda^*(u_0, v_0)(\alpha)$ . At the same time, let us denote by  $\lambda^*$  the stationary threshold, see Theorem 1 below, for the existence of solutions to

$$\beta\phi = \lambda g(\phi). \tag{2}$$

From the point of view of applications to Micro Electro Mechanical Systems, the value  $\lambda^*$  plays an important role as it is connected to the so-called ‘‘pull-in instability’’, see [1, 2] and references therein.

**Assumption 1** We impose the following assumptions on the damping term  $f$ :

- (f1)  $f \in C^1(\mathbb{R})$ ;
- (f2)  $f(0) = 0$ ;
- (f3)  $f(v)v > 0$  for  $v \in \mathbb{R} \setminus \{0\}$ ;
- (f4) there exist  $\eta > 0$  and  $\theta \geq 0$  such that  $|f(v)| \leq \eta |v|^{\theta+1}$  for  $v \in \mathbb{R}$ .

**Remark 1** Note that under these assumptions one has  $f'(0) \geq 0$ .

**Assumption 2** Let  $b \in (0, +\infty]$  and  $I = (-\infty, b)$ . We impose the following assumptions on the nonlinearity  $g$ :

- (g1)  $g(u) \in C^2(I)$ ;
- (g2)  $g(u) > 0$  for  $u \in I$ . Moreover, we normalize  $g(u)$  by  $g(0) = 1$ ;
- (g3)  $g'(0) \geq 0$ ;
- (g4)  $g''(u) > 0$  for  $u \in I$ ;
- (g5) If  $I = \mathbb{R}$ , for sufficiently large  $R > 0$ , there exists  $\tau > 0$  such that  $g''(u) > \tau$  for  $u > R$ .

If  $I = (-\infty, b), b < +\infty$ , the following hold:

$$\begin{aligned} \lim_{u \rightarrow b^-} g(u) = \lim_{u \rightarrow b^-} g'(u) = \lim_{u \rightarrow b^-} g''(u) = +\infty, \\ \lim_{u \rightarrow b^-} \int_0^u g(s) \, ds = +\infty \quad \text{and} \quad \lim_{u \rightarrow b^-} \left( \int_0^u g(s) \, ds - \frac{u}{2} g(u) \right) < 0. \end{aligned}$$

**Remark 2** Note that under these assumptions solutions to (2) are positive.

So far, a huge literature has been devoted to (1) and related problems from the theoretical point of view as well as from the point of view of applications, see for instance [6, 9, 12] and references therein.

In [5], Flores studies the following problem

$$\begin{cases} u_{tt} + \alpha u_t + u = \frac{\lambda}{(1-u)^2}, & \text{for } t > 0, \\ u(0) = u_0 \in [0, 1), \\ u_t(0) = v_0. \end{cases}$$

He proves that  $0 < \lambda(0, 0)(\alpha) < \lambda^*$  for  $\alpha \geq 0$  and that  $\lim_{\alpha \rightarrow +\infty} \lambda(0, 0)(\alpha) = \lambda^*$ . In [7], Haraux considers the following

$$u_{tt} + c|u_t|^\alpha u_t + |u|^\beta u = 0, \quad \text{for } c > 0, \alpha > 0, \beta > 0$$

and studies the existence of sign-changing solutions, number of the zeros and the decay estimates for solutions depending on the value of the parameters  $c, \alpha, \beta$ . Moreover, the results are generalized to

$$u_{tt} + c|u_t|^\alpha u_t + |u|^\beta u = f(t), \quad \text{for } c > 0, \alpha > 0, \beta > 0$$

for a continuous function  $f(t)$  with some decaying properties. In [11], Souplet studies the following backward equation

$$u_{tt} - |u_t|^\alpha u_t + |u|^\beta u = 0 \quad \text{for } \alpha > 0, \beta > 0$$

and proves the existence of unbounded global solutions, unbounded oscillatory solutions, as well as their blow-up rate and asymptotic behaviour. In [3, 10], the authors consider

$$u_{tt} - |u_t|^\alpha + \lambda |u|^\beta = 0, \quad \text{for } \lambda > 0, \alpha > 0, \beta > 0.$$

They investigate for which value of parameters one has existence of solutions, derive their asymptotic behaviour and classify the ground state solutions according to the value of parameters.

Here we aim at extending the result in [5] to more general nonlinearities  $f$  and  $g$ . Henceforth, we consider (1) and (2) under Assumptions 1 and 2 unless otherwise stated. We consider first the stationary problem (2) for which we obtain the bifurcation diagram of the solution set  $\{(\lambda, \phi)\}$ .

**Theorem 1** *There exists a unique  $p > 0$  such that*

$$g(p) - pg'(p) = 0.$$

Let

$$\lambda^* := \frac{\beta p}{g(p)}.$$

The following hold:

- (i) For any  $\lambda < \lambda^*$ , there are two solutions  $\phi_1 = \phi_1(\lambda)$  and  $\phi_2 = \phi_2(\lambda)$  of (2) with  $0 < \phi_1 < p < \phi_2 < b$  and

$$\lim_{\lambda \searrow 0} (\phi_1, \phi_2) = (0, b) \quad \text{and} \quad \lim_{\lambda \nearrow \lambda^*} (\phi_1, \phi_2) =: (\phi_1^*, \phi_2^*) = (p, p).$$

Moreover,  $\phi_1(\lambda)$  and  $\phi_2(\lambda)$  are respectively increasing and decreasing with respect to  $\lambda$ ;

- (ii) For  $\lambda = \lambda^*$ , there exists a unique solution  $\phi_1 = \phi_2 = p$  of (2);
- (iii) For  $\lambda > \lambda^*$ , do not exist solutions to (2).

In order to investigate the dynamical behaviour, we find the solution  $(\bar{\lambda}, \bar{\phi}_2)$  with the following properties:

**Theorem 2** *There exists a unique solution  $(\lambda, \phi_2(\lambda)) = (\bar{\lambda}, \bar{\phi}_2)$  of (2) satisfying the following:*

$$0 < \bar{\lambda} < \lambda^*, \quad p < \bar{\phi}_2 < b \quad \text{and} \quad \bar{\lambda} = \frac{\beta}{2} \frac{\bar{\phi}_2^{-2}}{\int_0^{\bar{\phi}_2} g(s) \, ds}.$$

Next, on the one hand we consider the time-dependent equation (1) and derive the conditions for the existence of global bounded solutions. To state the theorem, we define the functional

$$J_\lambda(u) := \frac{\beta}{2} u^2 - \lambda \int_0^u g(s) \, ds.$$

**Theorem 3** *For  $\lambda < \lambda^*$  and  $\alpha \geq 0$ , let*

$$D_0 := \left\{ (u, v) \in \mathbb{R}^2 \mid l(\lambda) < u < \phi_2(\lambda), \ v^2 < J_\lambda(\phi_2) - J_\lambda(u_0) \right\},$$

where  $l(\lambda)$  is a constant depending only on  $\lambda, \beta, g$  and  $u_0$ . If  $(u_0, v_0) \in D_0$ , then (1) has a unique global solution  $u \in W^{2,\infty}([0, \infty))$ . Moreover, if  $\alpha > 0$ , then the following holds

$$\lim_{t \rightarrow +\infty} \left( |u(t) - \phi_1(\lambda)| + |u_t(t)| \right) = 0. \tag{3}$$

On the other hand, we obtain the following blow-up result for  $\lambda > \lambda^*$  or  $\phi_2 < u_0 < b$ . Let  $T_\infty \in (0, +\infty]$ , be the maximal time of existence of a solution to (1). Then, we have the following

**Theorem 4** *Let  $\lambda > \lambda^*, \alpha \geq 0$  and  $u_0, v_0 \geq 0$ . The solution  $(u, u_t)$  of (1) blows up to  $(+\infty, +\infty)$  for  $b = +\infty$  and quenches to  $(b, +\infty)$  for  $0 < b < +\infty$  as  $t \rightarrow T_\infty$ , where  $T_\infty < +\infty$  if  $\alpha = 0$  or  $0 < b < +\infty$ .*

**Theorem 5** Let  $0 < \lambda \leq \lambda^*$  and  $\alpha \geq 0$ . For any  $\phi_2(\lambda) < u_0 < b$  and  $v_0 \geq 0$ , the same conclusion of Theorem 4 holds.

The main result of this paper is concerned with establishing an optimal dynamical threshold, indeed we have

**Theorem 6** Let  $f(v) = v$ . The function  $\lambda(0, 0)(\alpha)$  is continuous and monotone increasing with respect to  $\alpha \geq 0$  and satisfies:

- (i)  $\bar{\lambda} \leq \lambda(0, 0)(\alpha) < \lambda^*$  for  $\alpha \geq 0$ ;
- (ii)  $\lim_{\alpha \rightarrow 0^+} \lambda(0, 0)(\alpha) = \bar{\lambda}$ ;
- (iii)  $\lim_{\alpha \rightarrow +\infty} \lambda(0, 0)(\alpha) = \lambda^*$ .

This paper is organized as follows: In Sect. 2, we consider the stationary problem. Thanks to the assumptions on  $g$ , we get at most two positive solutions. In Sect. 3, we settle preliminary lemmas, involving energy and dynamical system, in order to investigate the dynamical behaviour. On the one hand, in Sect. 4, we establish the existence of a global solution and periodic orbit under some appropriate conditions. On the other hand, for large values of the perturbation parameter or large initial values, we prove that the solution blows up. In Sect. 5, we discuss qualitative properties of the orbit such as openness, monotonicity and continuity. In Sect. 6, we prove our main result, namely Theorem 6.

## 2 The Stationary Problem

Here we study the solution set of the function equation (2) and obtain the upper bound  $\lambda^*$  for the existence of solutions. We regard  $\phi = \phi(\lambda)$  as a function of the parameter  $\lambda$ .

**Proof of Theorem 1** Since solutions are positive, we consider in (2)  $\phi \geq 0$ . By Assumption 2,

$$F(u) := \frac{u}{g(u)}$$

is well-defined for  $u \geq 0$  and the following holds

$$F'(u) = \frac{1}{g(u)^2} (g(u) - ug'(u)).$$

From  $F(0) = F(b) = 0$  and  $F'(0) > 0$ , there exists  $p > 0$  such that  $F'(p) = 0$ . Next we show the uniqueness of such  $p$ . Set  $G(u) := g(u) - ug'(u)$ , and consider the sign of  $G(u)$ . We have  $G(0) = 1$ ,

$$G'(u) = -ug''(u) < 0$$

for all  $0 < u < b$ ,  $G'(0) = 0$  and  $G'(b) = -\infty$ , which implies that  $G(u) > 0$  for  $u < p$  and  $G(u) < 0$  for  $u > p$  and thus such  $p$  is uniquely determined. The statement follows by drawing the graph of  $y = \beta F(u)$  and  $y = \lambda$ .  $\square$

Define

$$H_\lambda(u) := \beta u - \lambda g(u).$$

Then  $\beta F(u) = \lambda$  is equivalent to  $H_\lambda(u) = 0$  and the sign of  $H_\lambda(u)$  is given by the value of  $\lambda$ .

**Corollary 1** For  $\lambda < \lambda^*$ , there exist  $\phi_1, \phi_2$  with  $\phi_1 < \phi_2 < b$  such that

- (i)  $H_\lambda(\phi_1) = H_\lambda(\phi_2) = 0$ ;
- (ii)  $H_\lambda(u) < 0$ , for  $u < \phi_1, \phi_2 < u < b$ ;
- (iii)  $H_\lambda(u) > 0$ , for  $\phi_1 < u < \phi_2$ .

Furthermore,  $H_\lambda(u)$  is increasing in  $(-\infty, q)$  and decreasing in  $(q, b)$ , where  $q$  satisfies  $\beta = \lambda g'(q)$  and  $p < q < \phi_2 < b$ .

**Proof** Noting that  $H_\lambda(\phi_1) = H_\lambda(\phi_2) = 0$ ,  $H'_\lambda(u) = \beta - \lambda g'(u)$  and that  $H''_\lambda(u) = -\lambda g''(u) < 0$ , we find  $q \in (\phi_1, \phi_2)$  uniquely satisfying  $H'_\lambda(q) = 0$ . Moreover, from

$$H'_\lambda(p) = \beta - \lambda g'(p) = \lambda^* \frac{g(p)}{p} - \lambda \frac{g(p)}{p} = (\lambda^* - \lambda) \frac{g(p)}{p} > 0,$$

one has  $p < q$ . □

**Remark 3** For  $\lambda = \lambda^*$ , we have  $\phi_1(\lambda^*) = \phi_2(\lambda^*) = p$  and  $H_{\lambda^*}(u) < 0$ , for  $u \neq p$ ,  $H_{\lambda^*}(p) = 0$ . Furthermore,  $H_{\lambda^*}(u)$  is increasing in  $(-\infty, p)$  and decreasing in  $(p, b)$ .

**Corollary 2** For  $\lambda > \lambda^*$ , we have

$$H_\lambda(u) < 0 \quad \text{for } u < b.$$

In particular, there exists  $\xi > 0$  such that

$$H_\lambda(u) \leq -\xi < 0$$

for all  $u \geq 0$ , where  $\xi$  depends only on  $\lambda, \beta$  and  $g$ .

**Proof** Set  $\xi = \lambda - \lambda^*$ , to have

$$\xi = \lambda - \beta F(p) \leq \lambda - \beta F(u) = -\frac{1}{g(u)} H_\lambda(u) \leq -H_\lambda(u)$$

as  $1 = g(0) \leq g(u)$  holds for all  $u \geq 0$ . □

**Lemma 1** For  $\lambda < \lambda^*$ , one has

$$H'_\lambda(\phi_1(\lambda)) = \sup_{\phi_1 \leq u \leq \phi_2} \frac{H_\lambda(u)}{u - \phi_1(\lambda)}.$$

**Proof** Noting that  $H_\lambda(\phi_1) = 0$ , we apply the mean value theorem to obtain the conclusion, as  $H'_\lambda(u) < 0$  for  $\phi_1 \leq u \leq \phi_2$ . □

**Proof of Theorem 2** Let us define

$$I(u) := 2 \int_0^u g(s) \, ds - ug(u), \quad 0 \leq u < b.$$

We claim that  $I(u_0) = 0$  for some  $p < u_0 < b$ . As a consequence, we find a solution  $(\lambda, \phi_2) = (\beta u_0/g(u_0), u_0)$  of (2) by Theorem 1 and

$$\lambda = \frac{\beta u_0}{g(u_0)} = \frac{\beta u_0^2}{2 \int_0^{u_0} g(s) \, ds} \in (0, \lambda^*).$$

Let us prove the claim. First we have

$$\begin{aligned} I(p) &= 2 \int_0^p g(s) \, ds - pg(p) \\ &= 2 \int_0^p g(s) \, ds - \frac{\beta}{\lambda^*} p^2 \\ &= \frac{2}{\lambda^*} \int_0^p (\lambda^* g(s) - \beta s) \, ds \\ &= -\frac{2}{\lambda^*} \int_0^p H_{\lambda^*}(s) \, ds > 0 \end{aligned}$$

by Remark 3. Observe that  $I'(u) = G(u) < 0$  and that  $I''(u) = G'(u) < 0$  for  $p < u < b$ , where  $G$  is defined as in the proof of Theorem 1. By the monotonicity of  $I$ , we need only to show that  $I(b) < 0$ . In the case of  $b < +\infty$ ,  $I(b) < 0$  follows from Assumption 2. In the case  $b = +\infty$ , since  $I''(+\infty) = G'(+\infty) = -\infty$  holds, we have  $I'(+\infty) = G(+\infty) = -\infty$  and finally  $I(+\infty) = -\infty$ , which proves the theorem. □

Let us give a few examples where  $(\lambda^*, p)$  and  $(\bar{\lambda}, \bar{\phi}_2)$  are explicitly known.

**Example 1** If  $\beta = 1$  and  $g(u) = e^u$ , Assumption 2 is satisfied with  $b = +\infty$ . We have

$$(\lambda^*, p) = \left( \frac{1}{e}, 1 \right)$$

and  $(\bar{\lambda}, \bar{\phi}_2)$  satisfies

$$\bar{\phi}_2 e^{\bar{\phi}_2} - 2e^{\bar{\phi}_2} + 2 = 0, \quad \bar{\lambda} = \bar{\phi}_2 e^{-\bar{\phi}_2}, \quad \bar{\phi}_2 \in (1, 2).$$

**Example 2** If  $\beta = 1$  and  $g(u) = 1 + u^{2k}$  for  $k \in \mathbb{N}$ , Assumption 2 is satisfied with  $b = +\infty$ . In particular, in the case  $k = 1$ , we have

$$(\lambda^*, p) = \left(\frac{1}{2}, 1\right) \quad \text{and} \quad (\bar{\lambda}, \bar{\phi}_2) = \left(\frac{\sqrt{3}}{4}, \sqrt{3}\right).$$

**Example 3** If  $\beta = 1$  and  $g(u) = 1/(1 - u)^p$  for  $p > 1$ ,  $g(u)$  has a singularity at  $u = 1$ . However, Assumption 2 is satisfied with  $b = 1$ . In particular, in the case of  $p = 2$  we have

$$(\lambda^*, p) = \left(\frac{4}{27}, \frac{1}{3}\right) \quad \text{and} \quad (\bar{\lambda}, \bar{\phi}_2) = \left(\frac{1}{8}, \frac{1}{2}\right).$$

### 3 A Dynamical System

Consider the following energy functional

$$E_\lambda(u, v) := \frac{\beta}{2}u^2 + \frac{1}{2}v^2 - \lambda \int_0^u g(s) \, ds.$$

Then,  $E_\lambda(u(t), u_t(t))$  turns out to be the Lyapunov function for (1). In fact, we have

$$\frac{d}{dt} E_\lambda(u(t), u_t(t)) = -\alpha f(u_t(t))u_t(t) \leq 0,$$

which yields

$$E_\lambda(u(t), u_t(t)) + \alpha \int_0^t f(u_t(r))u_t(r) \, dr = E_\lambda(u_0, v_0). \tag{4}$$

Hence, by use of

$$J_\lambda(u) := \frac{\beta}{2}u^2 - \lambda \int_0^u g(s) \, ds,$$

the energy inequality

$$J_\lambda(u) \leq J_\lambda(u_0) + \frac{1}{2}v_0^2 \tag{5}$$

holds by (4). Every local solution satisfies (5) as long as it exists. To extend the solution globally in time, we consider some properties of  $J_\lambda(u)$  in the following two lemmas:



**Lemma 2** For  $\lambda < \lambda^*$ ,  $J_\lambda(u)$  has a local minimum at  $u = \phi_1(\lambda)$  and a local maximum  $u = \phi_2(\lambda)$ . Moreover,  $J_\lambda(\phi_1(\lambda)) < 0$  holds.

**Proof** Since  $J'_\lambda(u) = \beta u - \lambda g(u) = H_\lambda(u)$  and  $J''_\lambda(u) = H'_\lambda(u)$ , we have  $J'_\lambda(\phi_1) = J'_\lambda(\phi_2) = 0$  and  $J''_\lambda(\phi_2) < 0 < J''_\lambda(\phi_1)$  by Corollary 1. Then  $J_\lambda(\phi_1(\lambda)) < J_\lambda(0) = 0$  for all  $\lambda \in (0, \lambda^*)$ , which completes the proof.  $\square$

**Lemma 3** The following hold:

- (i)  $J_\lambda(\phi_2(\lambda)) > 0$  for  $0 < \lambda < \bar{\lambda}$ ;
- (ii)  $J_\lambda(\phi_2(\lambda)) = 0$  for  $\lambda = \bar{\lambda}$ ;
- (iii)  $J_\lambda(\phi_2(\lambda)) < 0$  for  $\bar{\lambda} < \lambda < \lambda^*$ .

**Proof** From

$$\frac{d}{d\lambda} J_\lambda(\phi_2(\lambda)) = H_\lambda(\phi_2(\lambda)) \frac{d}{d\lambda} \phi_2(\lambda) - \int_0^{\phi_2(\lambda)} g(s) ds = - \int_0^{\phi_2(\lambda)} g(s) ds < 0$$

by Corollary 1,  $J_\lambda(\phi_2(\lambda))$  is monotone decreasing in  $\lambda$ . It follows from Theorem 2 that  $J_{\bar{\lambda}}(\phi_2(\bar{\lambda})) = 0$ , which yields the conclusions.  $\square$

In the next section, we consider dynamical properties of the solution of (1), which can be written in the following form

$$\frac{d}{dt} \begin{bmatrix} u \\ u_t \end{bmatrix} = \begin{bmatrix} u_t \\ -\alpha f(u_t) - H_\lambda(u) \end{bmatrix}. \tag{6}$$

Now under Assumptions 1 and 2, we obtain a local solution. Next, we establish the existence of a global solution exploiting Lemmas 2 and 3. For this purpose, we consider the stability of the equilibrium point. At the equilibrium point

$$(u, u_t) = (\phi_i(\lambda), 0)$$

for  $i = 1, 2$ , the linearized equation is given by

$$\frac{d}{dt} \begin{bmatrix} U \\ V \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -H'_\lambda(\phi_i) & -\alpha f'(0) \end{bmatrix} \begin{bmatrix} U \\ V \end{bmatrix}$$

and the eigenvalues  $\mu_i^\pm$  of coefficient matrix are given respectively by

$$\mu_i^\pm = \frac{-\alpha f'(0) \pm \sqrt{\{\alpha f'(0)\}^2 - 4H'_\lambda(\phi_i)}}{2}.$$

Along with Corollary 1, we have the following two lemmas:

**Lemma 4** For  $\lambda < \lambda^*$ , the equilibrium point  $(u, u_t) = (\phi_1(\lambda), 0)$  is a stable focus for  $0 < \alpha f'(0) < 2\sqrt{H'_\lambda(\phi_1(\lambda))}$ , a stable node for  $\alpha f'(0) \geq 2\sqrt{H'_\lambda(\phi_1(\lambda))}$  and it is a centre for  $\alpha f'(0) = 0$ .

**Lemma 5** For  $\lambda < \lambda^*$ , the equilibrium point  $(u, u_t) = (\phi_2(\lambda), 0)$  is a saddle for all  $\alpha \geq 0$ .

### 4 The Time-Dependent Problem

In this Section, for small parameters and small initial values, we establish the existence of a global solution by means of the Lyapunov function method. For the dissipative case, that is  $\alpha > 0$ , we show that the global solution converges to the stationary solution. For the conservative case, namely  $\alpha = 0$ , we consider the periodic orbit starting at  $(0, 0)$ . Note that a trivial periodic orbit does not exist by  $u_{tt}(0) = \lambda g(0) = \lambda > 0$ . Finally, we show that the solution becomes unbounded for large parameters as well as for large initial values.

**Proof of Theorem 3** We define by  $l(\lambda) \in (-\infty, \phi_1(\lambda))$  the point satisfying  $J_\lambda(u) = J_\lambda(\phi_2(\lambda))$ . Then, we have  $l(\lambda) < 0$ ,  $l(\lambda) = 0$  and  $l(\lambda) > 0$  for  $\lambda \in (0, \bar{\lambda})$ ,  $\lambda = \bar{\lambda}$  and  $\lambda \in (\bar{\lambda}, \lambda^*)$ , respectively. In fact, by  $J'_\lambda(u) = H_\lambda(u)$  and Corollary 1, we have  $J_\lambda(\phi_1) < J_\lambda(u_1) < 0 = J_\lambda(0) < J_\lambda(u_2)$  for  $u_2 < 0 < u_1 < \phi_1$ . For  $\lambda \in (0, \bar{\lambda})$ ,  $J_\lambda(u)$  attains a local maximum  $J_\lambda(\phi_2) > 0$  by Lemma 3. Hence we have  $l(\lambda) < 0$ . Other cases follows easily according to the sign of  $J_\lambda(\phi_2)$ . Note that  $J_\lambda(u_0) < J_\lambda(\phi_2)$  holds for  $l(\lambda) < u_0 < \phi_2(\lambda)$  by Lemma 2. Then, for  $(u_0, v_0) \in D_0$ , (5) yields

$$J_\lambda(u) < \frac{1}{2} (J_\lambda(\phi_2) + J_\lambda(u_0)) < J_\lambda(\phi_2)$$

and moreover

$$l(\lambda) < u(t) < J_\lambda^{-1} \left( \frac{J_\lambda(\phi_2) + J_\lambda(u_0)}{2} \right) < \phi_2, \tag{7}$$

where  $J_\lambda^{-1}$  is the inverse function of  $J_\lambda$  defined at  $(\phi_1, \phi_2)$ . Finally (4) implies that

$$u_t^2 \leq 2E_\lambda(u_0, v_0) + 2\lambda \int_0^u g(s) ds < 2E_\lambda(u_0, v_0) + 2\lambda g(\phi_2)\phi_2$$

and that  $(u(t), u_t(t))$  is uniformly bounded in  $\mathbb{R}^2$  for all  $t \geq 0$ . Therefore, since  $u_{tt}$  is also bounded for  $t \geq 0$  by (1), we have  $u(t) \in W^{2,\infty}([0, \infty))$ . If  $\alpha > 0$ , the Lyapunov function  $E_\lambda(u(t), u_t(t))$  is strictly decreasing in  $t$ . Thus it follows from Theorem 5.1.8 and Corollary 8.5.1 in [8] that the omega limit set  $\omega(u_0, v_0)$  is connected in  $\mathbb{R}^2$  and included in the solution set of (2). Since  $\bigcup_{t \geq 0} (u(t), u_t(t)) \notin \{(\phi_2, 0)\}$  by (7), we obtain (3). □

**Remark 4** Since  $l(\lambda) < \phi_1$  and  $J_\lambda(u) < J_\lambda(\phi_2)$  for all  $\lambda < \lambda^*$  and  $\phi_1 < u < \phi_2$ , we have  $(u_0, 0) \in D_0$  with  $\phi_1 < u_0 < \phi_2$ . In other words, the solution exists globally in time for the initial value  $(u_0, v_0)$  with  $\phi_1 < u_0 < \phi_2$  and  $v_0 = 0$ .

**Proof of Theorem 4** Let us divide the proof into four cases:

**I. The case**  $\alpha = 0$  and  $I = (-\infty, b)$ . Integrating (1) and thanks to Corollary 2, we have

$$u_t \geq v_0 + \xi t \geq 0 \quad \text{and} \quad u \geq u_0 + v_0 t + \frac{\xi}{2} t^2 \geq 0, \tag{8}$$

which implies that  $u(t)$  reaches  $b$  for finite  $T_\infty < \infty$ . Then

$$u_t^2(T_\infty) = 2E_\lambda(u_0, v_0) + 2\lambda \int_0^b g(s) \, ds - \beta b^2 = +\infty$$

by (4) and Assumption 2.

**II. The case**  $\alpha = 0$  and  $I = \mathbb{R}$ . Assume by contradiction that  $T_\infty = \infty$ . By (8), there exists sufficiently large  $T > 0$  such that

$$g''(u(t)) > \tau \tag{9}$$

for all  $t > T$ , where  $\tau$  is given in Assumption 2 and  $T$  can be taken as  $T = \sqrt{2\xi^{-1}R}$ . Then, integrating inequality (9) twice, with respect to  $u$ , over  $[u(T), u]$  we get

$$g(u(t)) \geq g(u(T)) + \frac{\tau}{2} (u(t) - u(T))^2$$

as  $g'(u) \geq 0$  for  $u \geq 0$ . Thus we have

$$\begin{aligned} u_{tt}(t) &= \lambda g(u(t)) - \beta u(t) \\ &\geq \lambda \left\{ g(u(T)) + \frac{\tau}{2} (u(t) - u(T))^2 \right\} - \beta u(t) \\ &= -H_\lambda(u(T)) + \frac{\tau\lambda}{2} (u(t) - u(T))^2 - \beta (u(t) - u(T)) \\ &> \frac{\tau\lambda}{2} (u(t) - u(T))^2 - \beta (u(t) - u(T)) \end{aligned} \tag{10}$$

for all  $t > T$ . Since  $u'(t) > 0$  holds for all  $t > T$  by (8), we have

$$u_t u_{tt} > \frac{\tau\lambda}{2} (u(t) - u(T))^2 u_t - \beta (u(t) - u(T)) u_t$$

and

$$\begin{aligned} \{u_t(t)\}^2 &> \{u_t(T)\}^2 + \frac{\tau\lambda}{3} (u(t) - u(T))^3 - \beta (u(t) - u(T))^2 \\ &> \left\{ \frac{\tau\lambda}{3} (u(t) - u(T)) - \beta \right\} (u(t) - u(T))^2. \end{aligned}$$

Take  $T_1 \in (T, +\infty)$  such that

$$\frac{\tau\lambda}{3} (u(t) - u(T)) - \beta > \frac{\tau\lambda}{6} (u(t) - u(T))$$

holds for all  $t > T_1$ . For instance, we can take  $T_1$  as follows

$$T_1 = \sqrt{\frac{2}{\xi}} \sqrt{\frac{6\beta}{\tau\lambda} + u(T)} > \sqrt{\frac{2u(T)}{\xi}} \geq \sqrt{\frac{2R}{\xi}} = T.$$

Then we have

$$u_t(t) > \sqrt{\frac{\tau\lambda}{6}} (u(t) - u(T))^{\frac{3}{2}}$$

and

$$(u(t) - u(T))^{-\frac{3}{2}} u_t(t) > \sqrt{\frac{\tau\lambda}{6}}$$

for all  $t > T_1$ . Integrating this inequality over  $[T_1, t]$ , we have

$$0 < \frac{2}{\sqrt{u(t) - u(T)}} < \frac{2}{\sqrt{u(T_1) - u(T)}} - \sqrt{\frac{\tau\lambda}{6}} (t - T_1),$$

which implies that  $\lim_{t \rightarrow T_2} u(t) = +\infty$ , where

$$T_2 = T_1 + \sqrt{\frac{6}{\tau\lambda} \frac{2}{\sqrt{u(T_1) - u(T)}}} < +\infty,$$

contradicting the maximality of  $T_\infty$  and hence necessarily  $T_\infty < +\infty$ . Thanks to (4), we also have

$$\xi u + E_\lambda(u_0, v_0) \leq - \int_0^u H_\lambda(s) ds + E_\lambda(u_0, v_0) = \frac{1}{2} u_t^2, \tag{11}$$

which implies that both  $u$  and  $u_t$  blow up to  $+\infty$  as  $t \rightarrow T_\infty$ .

**III. The case  $\alpha \neq 0$  and  $I = (-\infty, b)$ .** If  $v_0 = 0$ , we have

$$u_{tt}(0) = -H_\lambda(u_0) \geq \xi > 0. \tag{12}$$

Hence for  $v_0 \geq 0$ , we have  $u_t(t) > 0$  for sufficiently small  $t > 0$ . If there exists  $T_3 \in (0, T_\infty)$  such that  $u_t(t) > 0$  for all  $t \in (0, T_3)$  and  $u_t(T_3) = 0$ , then we have  $u_{tt}(T_3) > 0$  similarly to (12), which contradicts the positivity of  $u_t$ . Hence, if necessary, we retake the initial value as  $u_0 = u(T_4)$  and  $v_0 = v(T_4)$  for some  $T_4 > 0$  so that  $u_0 > 0, v_0 > 0, u(t) > 0$  and  $u_t(t) > 0$  hold for all  $t \in (0, T_\infty)$ . We estimate  $u_t(t)$ . First if  $u_{tt}(0) > 0$  holds, we have  $u_t(t) \geq v_0$  for sufficiently small  $t > 0$ . On the other hand, if  $u_{tt}(t) \leq 0$  holds for some  $t \geq 0$ , we have

$$\alpha \eta u_t^{\theta+1} \geq u_{tt} + \alpha f(u_t) = -H_\lambda(u) \geq \xi > 0$$

and then

$$u_t \geq \left(\frac{\xi}{\alpha\eta}\right)^{\frac{1}{\theta+1}}.$$

Thus we have

$$u_t \geq \min \left\{ v_0, \left(\frac{\xi}{\alpha\eta}\right)^{\frac{1}{\theta+1}} \right\} \equiv C_1 > 0 \tag{13}$$

for all  $t \in (0, T_\infty)$ , which yields

$$u(t) > u_0 + C_1 t \tag{14}$$

for all  $t \in (0, T_\infty)$ , which brings back to the same situation of case I above.

**IV. The case  $\alpha \neq 0$  and  $I = \mathbb{R}$ .** By estimates carried out in the case III, we have that (13) and (14) hold. Hence,  $(u, u_t)$  is unbounded in  $\mathbb{R}^2$  for  $t \in (0, T_\infty)$ , where  $T_\infty \leq +\infty$ . In the case of  $T_\infty < +\infty$ , both  $u$  and  $u_t$  blow up to  $+\infty$  as  $t \rightarrow T_\infty$  similarly to (11). Next let us consider the case  $T_\infty = +\infty$  and let us prove that  $u_t(t) \rightarrow +\infty$ , as  $t \rightarrow +\infty$ . Now suppose that there exists a constant  $C_2 > 0$  satisfying

$$C_1 < u_t < C_2$$

for all  $t \geq T$ . Then, buying the line of (10) we obtain the following differential inequality

$$\begin{aligned} u_{tt} + \alpha\eta u_t^{\theta+1} &\geq u_{tt} + \alpha f(u_t) \\ &> \frac{\tau\lambda}{2} (u(t) - u(T))^2 - \beta (u(t) - u(T)) \\ &= \frac{\tau\lambda}{2} \left(u(t) - u(T) - \frac{\beta}{\tau\lambda}\right)^2 - \frac{\beta^2}{2\tau\lambda} \end{aligned}$$

for all  $t \geq T$ . Thus for sufficiently large  $t > T$ , we have

$$u_{tt} \geq \frac{\tau\lambda}{2} \left(C_1 t + u_0 - u(T) - \frac{\beta}{\tau\lambda}\right)^2 - \frac{\beta^2}{2\tau\lambda} - \alpha\eta C_2^{\theta+1},$$

which yields  $u_t(t) \rightarrow +\infty$ , as  $t \rightarrow +\infty$  by integration and contradicting the boundedness of  $u_t$ . Hence, both  $u$  and  $u_t$  blow up to  $+\infty$  as  $t \rightarrow T_\infty$ . □

**Proof of Theorem 5** By hypothesis,  $\phi_2 < u(t) < b$  holds for sufficiently small  $t > 0$ . If  $v_0 = 0$ , we have

$$u_{tt}(0) = -H_\lambda(u_0) > -H_\lambda(\phi_2) = 0$$

by Corollary 1 and Remark 3. Hence we may assume that  $\phi_2 < u_0 < u(t) < b$  and  $u_t(t) > 0$  holds for all  $t \in (0, T_\infty)$ . Now set  $\xi = -H_\lambda(u_0) > 0$  and proceed as in the proof of Theorem 4.  $\square$

Let us denote by  $\gamma(t; \lambda, \alpha)$  the orbit of solution of (1) starting at  $(0, 0)$ . We note that from Lemma 4, one has  $\alpha f'(0) = 0$  if and only if  $(\phi_1(\lambda), 0)$  is a centre.

**Proposition 1** *Let  $\alpha = 0$ . We have that  $0 < \lambda < \bar{\lambda}$  is satisfied if and only if the orbit  $\gamma(t; \lambda, 0)$  is periodic. If  $\bar{\lambda} < \lambda < \lambda^*$ , then the orbit is unbounded.*

**Proof** In the conservative case  $\alpha = 0$ , we have

$$E_\lambda(u, u_t) = E_\lambda(0, 0) \iff J_\lambda(u) + \frac{1}{2}u_t^2 = 0 \tag{15}$$

by (4).  $(u_t)_t = -H_\lambda(u)$  is positive either for  $u < \phi_1$  or  $\phi_2 < u < b$  and negative for  $\phi_1 < u < \phi_2$  by Corollary 1. If  $0 < \lambda < \bar{\lambda}$ ,  $\gamma(t; \lambda, 0)$  exists globally and never passes  $(\phi_2, 0)$  by Theorem 3, (15) and Lemma 3. Hence  $\gamma(t; \lambda, 0)$  crosses the  $u$ -axis at some point  $(u_1, 0)$  for  $0 < u_1 < \phi_2$  by Theorem 5 and Remark 4, which implies that the orbit is periodic along with (15). Conversely, if the orbit is periodic, the orbit passes  $(u_1, 0)$  for  $u_1 \in (\phi_1, \phi_2)$  satisfying (15). Thus  $J_\lambda(u)$  has a zero in  $u \in (\phi_1, \phi_2)$ , which is equivalent to  $\lambda \in (0, \bar{\lambda})$  by the monotonicity of  $J_\lambda(u)$  with respect to  $u$  in Corollary 1, Lemmas 2 and 3. If  $\bar{\lambda} < \lambda < \lambda^*$ , there does not exist such  $u_1$ , which means that  $\phi_2 < u < b$  and  $u_t > 0$  for sufficiently large  $t$ . Finally apply Theorem 5 to conclude.  $\square$

**Remark 5** Since  $\lambda < \bar{\lambda}$  is equivalent to  $l(\lambda) < 0$ , we have  $(0, 0) \in D_0$ . Thus by Theorem 3, the orbit  $\gamma(t; \lambda, \alpha)$  for  $\alpha > 0$  exists globally in time and converges to  $(\phi_1(\lambda), 0)$ , as  $t \rightarrow +\infty$ . Hence along with Proposition 1, we have  $\bar{\lambda} \leq \lambda(0, 0)$  ( $\alpha$ ) for  $\alpha > 0$  and  $\bar{\lambda} = \lambda(0, 0)$  ( $0$ )  $< \lambda^*$ .

### 5 Properties of the Dissipative Orbit

In this Section, we study the properties of the orbit starting at the origin for the dissipative case, that is,  $\alpha > 0$ . Moreover, we also assume that  $f(v) = v$ . Then,  $(\phi_1(\lambda), 0)$  is a hyperbolic sink for all  $\lambda \in (\bar{\lambda}, \lambda^*)$ . The argument proceeds in the same way as in Sect. 3 of [5].

Let us first define a few sets which will be used in the sequel:

$$\begin{aligned} \Gamma &:= \left\{ (\lambda, \alpha) \in \mathbb{R}^2 \mid \bar{\lambda} < \lambda < \lambda^*, \alpha > 0 \right\}; \\ \Gamma_1 &:= \{ (\lambda, \alpha) \in \Gamma \mid \gamma(t; \lambda, \alpha) \rightarrow (\phi_1(\lambda), 0), \text{ as } t \rightarrow +\infty \}; \\ \Gamma_2 &:= \{ (\lambda, \alpha) \in \Gamma \mid \gamma(t; \lambda, \alpha) \rightarrow (\phi_2(\lambda), 0), \text{ as } t \rightarrow +\infty \}; \\ \Gamma_3 &:= \{ (\lambda, \alpha) \in \Gamma \mid \gamma(t; \lambda, \alpha) \text{ becomes unbounded as } t \rightarrow T_\infty \}, \end{aligned}$$

where  $\gamma(t; \lambda, \alpha)$  is the orbit of the solution  $(u, v) = (u, u_t)$  of (1) starting at  $(0, 0)$ . In what follows we will also use for convenience the following equivalent notations

$$\gamma(t) = \gamma(t; \lambda, \alpha) = \begin{bmatrix} u(t; \lambda, \alpha) \\ v(t; \lambda, \alpha) \end{bmatrix} = \begin{bmatrix} u(t) \\ v(t) \end{bmatrix}$$

for any fixed  $\alpha, \lambda > 0$ .

**Lemma 6**  $\Gamma_1$  and  $\Gamma_3$  are open subsets of  $\Gamma$ .

**Proof** If we take  $(\lambda_0, \alpha_0) \in \Gamma_1$ , then  $(u, v) = (\phi_1(\lambda_0), 0)$  is a hyperbolic sink by Lemma 4. Hence, there exists  $r > 0$  depending only on  $\lambda_0, \alpha_0, \beta$  and  $g$  such that we can find an invariant disk  $B_r(\phi_1(\lambda_0), 0)$  of radius  $r$  centred at  $(\phi_1(\lambda_0), 0)$  for the dynamical system induced by (6). In other words, there exists  $T > 0$  such that  $\gamma(t; \lambda_0, \alpha_0) \subset B_r(\phi_1(\lambda_0), 0)$  for all  $t > T$ . If necessary, we can take  $r > 0$  so small such that

$$B_r(\phi_1(\lambda_0), 0) \subset \left\{ (u, v) \in \mathbb{R}^2 \mid u \leq p \right\},$$

where  $p$  is defined in Theorem 1 and satisfies  $\phi_1(\lambda_0) < p < \phi_2(\lambda_0)$ . For  $(\lambda, \alpha) \in \Gamma$  sufficiently close to  $(\lambda_0, \alpha_0)$ ,  $\gamma(t; \lambda, \alpha) \subset B_r(\phi_1(\lambda_0), 0)$  for sufficiently large  $t$  thanks to the continuous dependence on parameters. Since  $(\phi_1(\lambda), 0)$  is also a hyperbolic sink,  $\gamma(t; \lambda, \alpha)$  converges  $(\phi_1(\lambda), 0)$  as  $t \rightarrow +\infty$ . Hence we have  $(\lambda, \alpha) \in \Gamma_1$ . Now take  $(\lambda_0, \alpha_0) \in \Gamma_3$  and define

$$\mathcal{U}_{\lambda, \alpha} := \left\{ (u, v) \in \mathbb{R}^2 \mid \phi_2(\lambda) < u < b, 0 < v < -\frac{1}{\alpha} H_\lambda(u) \right\}.$$

For sufficiently large  $t > 0$ , we have  $\gamma(t; \lambda_0, \alpha_0) \subset \mathcal{U}_{\lambda_0, \alpha_0}$  by a phase plane analysis of (6) together with  $u_{tt}(0) = \lambda_0 > 0$  and Remark 4. Again, by continuous dependence, for large  $t > 0$  we have  $\gamma(t; \lambda, \alpha) \subset \mathcal{U}_{\lambda_0, \alpha_0}$  for  $(\lambda, \alpha) \in \Gamma$  sufficiently close to  $(\lambda_0, \alpha_0)$ . Since we have  $\phi_2(\lambda) < u(t; \lambda, \alpha) < b$  and  $u_t(t; \lambda, \alpha) > 0$  for sufficiently large  $t$ ,  $\gamma(t; \lambda, \alpha)$  becomes unbounded by Theorem 5, which proves that  $(\lambda, \alpha) \in \Gamma_3$ . □

**Proposition 2** We have  $\Gamma = \bigcup_{i=1}^3 \Gamma_i$ .

**Proof** Note that every bounded orbit for  $\alpha > 0$  converges to  $\phi_1$  or  $\phi_2$  as  $t \rightarrow +\infty$  by Corollary 8.5.1 in [8]. If an unbounded orbit  $\gamma(t; \lambda, \alpha)$  exists, then  $u(t) \geq 0$  and  $u_t(t) \geq 0$  hold for sufficiently small  $t \geq 0$  by  $u_{tt}(0) = \lambda > 0$  and the orbit  $\gamma(t; \lambda, \alpha)$  enters  $\mathcal{U}_{\lambda, \alpha}$  by Remark 4. Then we have  $(\lambda, \alpha) \in \Gamma_3$  as in the proof of Theorem 5. □

Let  $m$  be a negative constant to be determined later. Let us define the line segments  $s_i$  for  $i = 1, 2, 3$  and the triangular region  $\mathcal{T}$  as follows:

$$\begin{aligned} s_1 &:= \{(u, v) = (\phi_2(\lambda), v) \mid m(\phi_2(\lambda) - \phi_1(\lambda)) \leq v \leq 0\}; \\ s_2 &:= \{(u, v) = (u, 0) \mid \phi_1(\lambda) \leq u \leq \phi_2(\lambda)\}; \end{aligned}$$

$$s_3 := \{(u, v) = (u, m(u - \phi_1(\lambda))) \mid \phi_1(\lambda) \leq u \leq \phi_2(\lambda)\};$$

$$\mathcal{T} := \{(u, v) \in \mathbb{R}^2 \mid \phi_1(\lambda) \leq u \leq \phi_2(\lambda), m(u - \phi_1(\lambda)) \leq v \leq 0\}.$$

**Proposition 3** For  $\alpha > 2\sqrt{H'_\lambda(\phi_1)}$ , there exists a heteroclinic orbit from  $(\phi_2, 0)$  to  $(\phi_1, 0)$ .

*Proof.* Since the vector field on  $s_1$  and  $s_2$  defined by (6) points inward  $\mathcal{T}$ , we can choose  $m < 0$  such that the vector field on  $s_3$  also points inward  $\mathcal{T}$ . Denote by  $N$  the normal vector on  $s_3$

$$N = \begin{bmatrix} m \\ -1 \end{bmatrix}$$

and by  $V$  the vector on  $s_3$  defined as follows

$$V = \begin{bmatrix} u_t \\ -\alpha u_t - H_\lambda(u) \end{bmatrix} = \begin{bmatrix} m(u - \phi_1) \\ -\alpha m(u - \phi_1) - H_\lambda(u) \end{bmatrix}$$

for  $\phi_1 < u < \phi_2$ . Let us compute the inner product between  $N$  and  $V$  to obtain

$$\begin{aligned} N \cdot V &= m^2(u - \phi_1) + \alpha m(u - \phi_1) + H_\lambda(u) \\ &= (u - \phi_1) \left( m^2 + \alpha m + \frac{H_\lambda(u)}{u - \phi_1} \right) \\ &\leq (u - \phi_1) \left( m^2 + \alpha m + H'_\lambda(\phi_1) \right) \end{aligned}$$

by Lemma 4. Take  $m$  such that  $N \cdot V < 0$ , that is,  $\mu_1^- < m < \mu_1^+$ , where  $\mu_1^\pm$  are defined in Sect. 3 with  $f'(0) = 1$ . Now the branch of the unstable manifold of  $(\phi_2, 0)$  that points inside the region

$$\{(u, v) \in \mathbb{R}^2 \mid u < \phi_2(\lambda), v < 0\}$$

enters  $\mathcal{T}$  and does not leave it. Hence this bounded orbit in  $\mathcal{T}$  converges to  $(\phi_1, 0)$  as  $t \rightarrow +\infty$ . □

**Remark 6** A typical example for  $f$  is given by  $f(v) = |v|^\gamma$ , where  $\gamma \geq 1$ . However, we consider the case of  $f(v) = v$ . Indeed, if we take  $f(v) = |v|^2$ , Proposition 3 does not hold because

$$N \cdot V = m^2(u - \phi_1) + \alpha m^2(u - \phi_1)^2 + H_\lambda(u) > 0$$

for any  $m \in \mathbb{R}$  and  $u \in (\phi_1, \phi_2)$ .



In order to prove the monotonicity of the orbit, let us introduce the following notation

$$\gamma_i(t) = \gamma(t; \lambda_i, \alpha) = \begin{bmatrix} u(t; \lambda_i, \alpha) \\ v(t; \lambda_i, \alpha) \end{bmatrix} = \begin{bmatrix} u_i(t) \\ v_i(t) \end{bmatrix}$$

for any fixed  $\alpha > 0$  and  $0 < \lambda_i < \lambda^*$ ,  $i = 1, 2$ . We next prove the monotonicity of the orbit in  $\lambda$  for fixed  $\alpha$ .

**Proposition 4** *For fixed  $\alpha > 0$ ,  $v(t; \lambda, \alpha)$  is increasing with respect to  $\lambda$  as long as  $v(t; \lambda, \alpha) > 0$ . Moreover, if there exists  $\lambda_0 \in (0, \lambda^*)$  such that  $(\lambda_0, \alpha) \in \Gamma_2 \cup \Gamma_3$ , then we have  $(\lambda, \alpha) \in \Gamma_3$  for all  $\lambda > \lambda_0$ .*

**Proof** Since we have  $v_i(0) = 0$  and  $(v_i)_t(0) = \lambda_i$ ,  $v_1(t) < v_2(t)$  holds for sufficiently small  $t > 0$ . For the second component of vector field defined in (6), we have

$$\frac{d}{d\lambda} (-\alpha v - H_\lambda(u)) = g(u) > 0.$$

Let us prove the second statement. If  $(\lambda_0, \alpha) \in \Gamma_3$ , the monotonicity yields  $(\lambda, \alpha) \in \Gamma_3$  for all  $\lambda > \lambda_0$ . If  $(\lambda_0, \alpha) \in \Gamma_2$ , the monotonicity of  $\phi_i(\lambda)$  and that of the orbit in  $\lambda$  imply  $(\lambda, \alpha) \in \Gamma_3$  by Theorem 5. □

As stated after the proof of Proposition 6 in [5], the stable local manifold of the saddle  $(\phi_2(\lambda), 0)$  plays a crucial role in determining the threshold of the parameter  $\lambda$ . Now for fixed  $\lambda \in (\bar{\lambda}, \lambda^*)$ , we regard the behaviour of the local stable manifold as a function of  $\alpha$ . We shall prove that the manifold crosses the positive  $u$ -axis for small  $\alpha > 0$ . Then the orbit  $\gamma(t)$  cannot approach the stationary points, which implies that  $(\lambda, \alpha) \in \Gamma_3$ . On the other hand, for large  $\alpha > 0$ , we prove that the manifold crosses the negative  $u$ -axis. In this case, the solution  $(u, v)$  is bounded for all  $t \geq 0$  and  $(\lambda, \alpha) \notin \Gamma_2$ , which means that  $(\lambda, \alpha) \in \Gamma_1$ . Finally, we uniquely determine  $\alpha^*(\lambda) > 0$  such that the manifold crosses the  $u$ -axis at  $u = 0$ . Then we establish the continuity and monotonicity of  $\alpha^*(\lambda) > 0$  with respect to  $\lambda$  and define  $\lambda^*(0, 0) (\alpha)$  by the inverse function of  $\alpha^*(\lambda)$ . In order to analyze the stable manifold of  $(\phi_2(\lambda), 0)$ , we perform the following change of variables in (1)

$$\begin{cases} t = -s, \\ U(s) = \phi_2(\lambda) - u(t), \\ V(s) = v(t) \end{cases}$$

and consider

$$\frac{d}{ds} \begin{bmatrix} U \\ V \end{bmatrix} = \begin{bmatrix} V \\ \alpha V + H_\lambda(\phi_2(\lambda) - U) \end{bmatrix} \tag{16}$$

for  $s < 0$  with  $U(-\infty) = V(-\infty) = 0$ . As we have seen in Sect. 3, the eigenvalues  $\eta^\pm$  of the linearized operator at  $(U, V) = (0, 0)$  corresponding to (16) are given by

$$\eta^\pm = \eta^\pm(\alpha) = \frac{\alpha \pm \sqrt{\alpha^2 - 4H'_\lambda(\phi_2)}}{2}.$$

The branch of the local unstable manifold can be expressed by the graph  $V = \Phi(U; \lambda, \alpha)$  as long as  $V(s) > 0$  since  $V(s) = U_s(s) > 0$ . First, it is clear that  $\Phi(0; \lambda, \alpha) = 0$ . Since we concentrate on the unstable manifold, we have  $(d\Phi/dU)(0; \lambda, \alpha) = \eta^+$  by  $H'_\lambda(\phi_2) < 0$ . Finally, we have

$$\frac{d\Phi}{dU}(U; \lambda, \alpha) = \frac{V_s}{U_s} = \alpha + \frac{H_\lambda(\phi_2(\lambda) - U)}{\Phi(U; \lambda, \alpha)}. \tag{17}$$

We denote  $\Phi_i(U) = \Phi(U; \lambda, \alpha_i)$  for fixed  $\lambda \in (\bar{\lambda}, \lambda^*)$  and  $\alpha_i > 0$  with  $i = 1, 2$  and establish the monotonicity of  $\Phi(U; \lambda, \alpha)$  with respect to  $\alpha$ .

**Proposition 5** *For fixed  $\lambda \in (0, \lambda^*)$ , let  $0 < \alpha_1 < \alpha_2$ . The graph  $V = \Phi_2(U)$  stays above the graph  $V = \Phi_1(U)$ . Moreover, the graph  $V = \Phi_2(U)$  and  $V = \Phi_1(U)$  never intersect each other as long as they are defined.*

**Proof** Since we have  $\Phi_2(0) - \Phi_1(0) = 0$  and  $(d\Phi_2/dU)(0) - (d\Phi_1/dU)(0) = \eta^+(\alpha_2) - \eta^+(\alpha_1) > 0$ , we obtain  $\Phi_2(U) - \Phi_1(U) > 0$  for sufficiently small  $U > 0$ . Thanks to (17),  $V = \Phi_2(U)$  and  $V = \Phi_1(U)$  can not intersect each other.  $\square$

For fixed  $\lambda \in (\bar{\lambda}, \lambda^*)$ , let

$$K(\lambda) := \{\alpha \geq 0 \mid \text{there exists } P_\alpha > 0 \text{ such that } \Phi(P_\alpha; \lambda, \alpha) = 0\}.$$

We are interested in the set  $L(\lambda)$  of all points  $P_\alpha$  defined by

$$L(\lambda) := \{P_\alpha \mid \alpha \in K(\lambda)\}.$$

$L(\lambda)$  consists of the points where the unstable manifold intersects the positive  $U$ -axis. We shall show that  $K(\lambda)$  is a non-empty interval and that  $L(\lambda)$  is an unbounded interval. For this purpose, let us define two lines parallel to the  $V$ -axis as follows:

$$\begin{aligned} M(\lambda) &:= \{(\phi_2(\lambda) - \phi_1(\lambda), V) \mid V \geq 0\}; \\ M_\Phi(\lambda) &:= \{(\phi_2(\lambda) - \phi_1(\lambda), \Phi(\phi_2(\lambda) - \phi_1(\lambda); \lambda, \alpha)) \mid \alpha \geq 0\}. \end{aligned}$$

Thanks to the continuity of the intersection point with respect to  $\alpha$  proved in [4],  $M_\Phi(\lambda)$  is an interval in  $M(\lambda)$ .

**Lemma 7**  $M_\Phi(\lambda) = \{\phi_2(\lambda) - \phi_1(\lambda)\} \times [\Phi(\phi_2(\lambda) - \phi_1(\lambda); \lambda, 0), +\infty)$  for  $\lambda \in (\bar{\lambda}, \lambda^*)$ .

**Proof** Noting that  $H_\lambda(\phi_2 - U) \geq 0$  for  $U \in [0, \phi_2 - \phi_1]$ , we have

$$\frac{d\Phi}{dU}(U; \lambda, \alpha) \geq \alpha$$

for any  $\alpha \geq 0$  and  $U \in [0, \phi_2 - \phi_1]$  by (17). Hence integrating this inequality over  $[0, \phi_2 - \phi_1]$ , we obtain

$$\Phi(\phi_2 - \phi_1; \lambda, \alpha) \geq \alpha(\phi_2 - \phi_1) \rightarrow +\infty, \text{ as } \alpha \rightarrow +\infty.$$

□

Next we establish a few properties of  $K(\lambda)$  and  $L(\lambda)$ , for which we set  $A(\lambda) := \sup K(\lambda)$  and  $P(\lambda) := \sup L(\lambda)$ .

**Lemma 8**  $K(\lambda)$  and  $L(\lambda)$  are nonempty intervals for  $\lambda \in (\bar{\lambda}, \lambda^*)$ .

**Proof** There exists some  $u \in (0, \phi_1)$  such that  $J_\lambda(u) = J_\lambda(\phi_2)$  holds. From the arguments of phase plane analysis in Theorem 3, Remark 4 and Proposition 1,  $0 \in K(\lambda)$  holds. Hence  $K(\lambda) \neq \emptyset$  follows for  $\lambda \in (\bar{\lambda}, \lambda^*)$ . Finally, as mentioned in Proposition 5, the continuity and monotonicity of  $\Phi$  yield the conclusion. □

We see that  $K(\lambda)$  and  $L(\lambda)$  are intervals. Next we prove that the right endpoint of  $K(\lambda)$  is open.

**Lemma 9**  $K(\lambda) = [0, A(\lambda))$  for  $\lambda \in (\bar{\lambda}, \lambda^*)$ .

**Proof** We may assume  $A(\lambda) < +\infty$ . Suppose by contradiction that  $K(\lambda) = [0, A(\lambda)]$ . The orbit  $\Phi(U; \lambda, A(\lambda))$  intersects the line  $M(\lambda)$ . Then, by definition we can find  $P_{A(\lambda)} > 0$  such that  $\Phi(P_{A(\lambda)}; \lambda, A(\lambda)) = 0$ . Next, the orbit starting at  $(P_{A(\lambda)}, 0)$  enters the region  $\{(U, V) \in \mathbb{R}^2 \mid V < 0\}$  with the  $U$ -coordinate decreasing. Since the set

$$\{(\phi_2(\lambda) - \phi_1(\lambda), \Phi(\phi_2(\lambda) - \phi_1(\lambda); \lambda, \alpha)) \mid \alpha \in [0, A(\lambda) + \varepsilon]\}$$

for any  $\alpha \geq 0$  and sufficiently small  $\varepsilon > 0$  describes a closed interval in the line  $M(\lambda)$  by the continuity and monotonicity of  $\Phi$  with respect to  $\alpha$  and Lemma 7, the behaviour of the orbit  $\Phi(U; \lambda, \alpha)$  is the same as that of  $\Phi(U; \lambda, A(\lambda))$  for  $\alpha \in (A(\lambda), A(\lambda) + \varepsilon)$  with sufficiently small  $\varepsilon > 0$  by the continuous dependence on the parameter and initial value. Hence we obtain  $P_\alpha$  satisfying  $\Phi(P_\alpha; \lambda, \alpha) = 0$  for  $\alpha \in (A(\lambda), A(\lambda) + \varepsilon)$ , which contradicts the definition of  $A(\lambda)$ . □

Finally, we prove that  $L(\lambda)$  is an unbounded interval.

**Lemma 10**  $P(\lambda) = +\infty$  for  $\lambda \in (\bar{\lambda}, \lambda^*)$ .

**Proof** First, we deal with the case of  $A(\lambda) < +\infty$ . Assume by contradiction that  $P(\lambda) < +\infty$ . The orbit  $\Phi(U; \lambda, \alpha)$  intersects the line  $M(\lambda)$  for any  $\alpha \in (0, A(\lambda))$ . In addition, due to (17),  $(d\Phi)/(dU)(U; \lambda, \alpha) < +\infty$  on every finite interval for

any  $\alpha \in (0, A(\lambda))$  as long as  $\Phi(U; \lambda, \alpha) > 0$ . Note that  $A(\lambda) \notin K(\lambda)$ . Thus the continuous dependence on the parameter yields  $\Phi(U; \lambda, A(\lambda)) > 0$  for all  $U > 0$ . Because of the continuous dependence, for any  $\alpha \in (A(\lambda) - \varepsilon, A(\lambda))$  with sufficiently small  $\varepsilon > 0$ , we have  $\Phi(U; \lambda, \alpha) > 0$  for all  $U \geq 0$ , which contradicts  $\alpha \in K(\lambda)$ . Next, we treat the case of  $A(\lambda) = +\infty$ . For all  $\alpha \geq 0$ , there exists  $P_\alpha \in (\phi_2 - \phi_1, +\infty)$  such that we have  $\Phi(P_\alpha; \lambda, \alpha) = 0$ . The intersection point of  $V = \Phi(U; \lambda, \alpha)$  and  $M(\lambda)$  lies in the region  $\{(U, V) \mid V' > 0\}$ . The graph of  $V = \Phi(U; \lambda, \alpha)$  must leave the region at the point  $(U_\alpha, V_\alpha)$  satisfying

$$V_\alpha = -\frac{1}{\alpha} H_\lambda(\phi_2 - U_\alpha), \quad \phi_2 - \phi_1 < U_\alpha < P_\alpha \quad \text{and} \quad \Phi(\phi_2 - \phi_1; \lambda, \alpha) < V_\alpha.$$

Then we have

$$H_\lambda(\phi_2 - U_\alpha) < -\alpha \Phi(\phi_2 - \phi_1; \lambda, \alpha) \rightarrow -\infty$$

as  $\alpha \rightarrow +\infty$  by Lemma 7, which implies that  $b > p > \phi_1 > \phi_2 - U_\alpha \rightarrow -\infty$  as  $\alpha \rightarrow +\infty$ , where  $p$  is defined in Theorem 1. Eventually we obtain  $P_\alpha > U_\alpha \rightarrow \infty$ , as  $\alpha \rightarrow +\infty$ . □

### 6 Proof of Theorem 6

The strategy to prove our main result is the following. For fixed  $\lambda \in (\bar{\lambda}, \lambda^*)$ , there exists  $\alpha^*(\lambda)$  such that the solution of (1) exists globally in time for  $\alpha^*(\lambda) < \alpha < +\infty$  and that is unbounded for  $0 < \alpha < \alpha^*(\lambda)$ . Then, the function  $\alpha^*(\lambda)$  turns out to be monotone increasing and continuous with respect to  $\lambda$ . Finally, we have  $\alpha^*(\lambda) \rightarrow +\infty$ , as  $\lambda \rightarrow \lambda^*$ . We denote the inverse function of  $\alpha^*(\lambda)$  by  $\lambda(0, 0)(\alpha)$  which we prove to have the desired properties as in Theorem 6.

**Lemma 11** *For fixed  $\lambda \in (\bar{\lambda}, \lambda^*)$ , there exists  $\alpha^*(\lambda) > 0$  such that the solution of (1) exists globally in time for  $\alpha^*(\lambda) < \alpha < +\infty$  and that becomes unbounded for  $0 < \alpha < \alpha^*(\lambda)$ .*

**Proof** For any  $\alpha \in (0, A(\lambda))$ , we have  $P_\alpha > P_0$  such that  $\Phi(P_\alpha; \lambda, \alpha) = 0$  by Proposition 5. Let

$$x_\alpha = \phi_2 - P_\alpha.$$

In other words,  $x_\alpha$  is an intersection point between the stable manifold of the saddle  $(\phi_2(\lambda), 0)$  and the  $u$ -axis. Since the point  $P_0$  is determined by the orbit of the conservative case, we have  $x_0 > 0$ , that is,  $P_0 < \phi_2$  by Proposition 1 for the case  $\lambda \in (\bar{\lambda}, \lambda^*)$ . By Lemma 10, the monotonicity and continuity of the intersection point, there exists  $\alpha^*(\lambda) > 0$  uniquely determined such that  $x_{\alpha^*(\lambda)} = 0$ , or  $P_{\alpha^*(\lambda)} = \phi_2$ . First, we consider the case of  $0 < \alpha < \alpha^*(\lambda)$ . Since we have

$$\phi_2 - \phi_1 < P_0 < P_\alpha < \phi_2 \iff 0 < x_\alpha < x_0 < \phi_1,$$

$(0, 0)$  is not in the domain of attraction of  $(\phi_1(\lambda), 0)$  for the orbit  $\gamma(t; \lambda, \alpha)$ . Hence it enters

$$Z_1 := \left\{ (u, v) \in \mathbb{R}^2 \mid 0 < u < b, v > 0, v_t > 0 \right\}$$

and

$$Z_2 := \left\{ (u, v) \in \mathbb{R}^2 \mid 0 < u < b, v > 0, v_t < 0 \right\}.$$

Then, we can find  $T > 0$  such that  $\phi_2(\lambda) < u(T) < b, v(T) > 0$  and  $v_t(T) = 0$  by Proposition 5. Hence the solution blows up by Theorem 5.

Next, we consider the case  $\alpha > \alpha^*(\lambda)$ . Similarly, we obtain

$$\phi_2 < P_\alpha \iff x_\alpha < 0.$$

Therefore,  $\gamma(t; \lambda, \alpha)$  enters  $Z_1$  and then  $Z_2$ . Since  $\gamma(t; \lambda, \alpha)$  stays below the stable manifold of  $(\phi_2(\lambda), 0)$ , it necessarily intersects the  $u$ -axis between  $\phi_1(\lambda)$  and  $\phi_2(\lambda)$ . Hence the solution exists globally in time by Remark 4. □

**Lemma 12**  $\alpha^*(\lambda)$  is monotone increasing for  $\lambda \in (\bar{\lambda}, \lambda^*)$ .

*Proof* Let  $\bar{\lambda} < \lambda_1 < \lambda_2 < \lambda^*$ . We have  $(\lambda_1, \alpha^*(\lambda_1)) \in \Gamma_2$  by Lemmas 6, 11 and Proposition 2. Thus we have  $(\lambda_2, \alpha^*(\lambda_1)) \in \Gamma_3$  by Proposition 4. Hence, from Lemma 11,  $\alpha^*(\lambda_1) < \alpha^*(\lambda_2)$  follows. □

**Lemma 13**  $\alpha^*(\lambda)$  is continuous for  $\lambda \in (\bar{\lambda}, \lambda^*)$ .

*Proof* Let  $\lambda_0 \in (\bar{\lambda}, \lambda^*)$  be fixed. Then we have  $(\lambda_0, \alpha^*(\lambda_0) - \varepsilon) \in \Gamma_3$  and  $(\lambda_0, \alpha^*(\lambda_0) + \varepsilon) \in \Gamma_1$  for any chosen  $\varepsilon \in (0, \alpha^*(\lambda_0))$ . By Lemma 6, there exists  $\delta > 0$  such that  $(\lambda, \alpha^*(\lambda_0) - \varepsilon) \in \Gamma_3$  and  $(\lambda, \alpha^*(\lambda_0) + \varepsilon) \in \Gamma_1$  hold for  $|\lambda - \lambda_0| < \delta$ . Lemma 11 implies that  $\alpha^*(\lambda_0) - \varepsilon < \alpha^*(\lambda) < \alpha^*(\lambda_0) + \varepsilon$ , and in turn  $|\alpha^*(\lambda) - \alpha^*(\lambda_0)| < \varepsilon$ . □

**Lemma 14** We have  $\lim_{\lambda \nearrow \lambda^*} \alpha^*(\lambda) = +\infty$ .

*Proof* Assume that for  $\lambda = \lambda^*$  there exists  $\alpha_0 > 0$  such that  $\gamma(t; \lambda^*, \alpha_0) \rightarrow (p, 0)$  as  $t \rightarrow +\infty$ , where  $\phi_1(\lambda^*) = \phi_2(\lambda^*) = p$ . We will show that this assumption leads us to a contradiction. Indeed, in this case we have  $(\lambda, \alpha) \in \Gamma_1$  for all  $0 < \lambda < \lambda^*$  and  $\alpha \geq \alpha_0$  by  $\alpha^*(\lambda^*) \leq \alpha_0$  and Lemma 12. Hence  $\gamma(t; \lambda^*, \alpha) \rightarrow (p, 0)$  as  $t \rightarrow +\infty$  for all  $\alpha \geq \alpha_0$ . Now, again through the transformation

$$\begin{cases} t = -s, \\ U(s) = p - u(t), \\ V(s) = v(t), \end{cases}$$

(6) is equivalent to

$$\frac{d}{ds} \begin{bmatrix} U \\ V \end{bmatrix} = \begin{bmatrix} V \\ \alpha V + H_{\lambda^*}(p - U) \end{bmatrix}$$

for  $s < 0$  with  $U(-\infty) = V(-\infty) = 0$ . We have one-parameter family of unstable manifold of  $(U, V) = (0, 0)$  with a branch which enters

$$W_1 := \left\{ (U, V) \in \mathbb{R}^2 \mid U > 0, V > \Psi_{\alpha, \lambda^*}(U) \right\},$$

$$W_2 := \left\{ (U, V) \in \mathbb{R}^2 \mid U > 0, 0 < V < \Psi_{\alpha, \lambda^*}(U) \right\}$$

and approaches  $(U, V) = (p, 0)$ , where

$$\Psi_{\alpha, \lambda}(U) \equiv -\frac{1}{\alpha} H_\lambda(p - U).$$

It is clear that  $V_s > 0$  in  $W_1$  and that  $V_s < 0$  in  $W_2$ . Each of these branches of the unstable manifolds for  $\alpha \geq \alpha_0$  is the graph of a function  $V = \Phi(U; \lambda^*, \alpha)$  defined for  $U \in [0, p]$ . Moreover, similarly to (17), we have

$$\frac{d\Phi}{dU}(U; \lambda^*, \alpha) = \alpha + \frac{H_{\lambda^*}(p - U)}{\Phi(U; \lambda^*, \alpha)}$$

and

$$\frac{d\Phi}{dU}(0; \lambda^*, \alpha) = \eta^+(\alpha) = \alpha.$$

As in Proposition 5, the graph  $V = \Phi(U; \lambda^*, \alpha_2)$  stays above the graph  $V = \Phi(U; \lambda^*, \alpha_1)$  for  $\alpha_0 \leq \alpha_1 < \alpha_2$ . Let  $(U_0, V_0)$  be the intersection point of the branch  $V = \Phi(U; \lambda^*, \alpha_0)$  of the unstable manifold with the graph  $V = \Psi_{\alpha_0, \lambda^*}(U)$ . Then we have  $U_0 > 0, V_0 > 0$  and

$$V_0 = \Phi(U_0; \lambda^*, \alpha_0).$$

By monotonicity and continuity, there exists  $\delta > 0$  such that

$$\frac{1}{2} V_0 = \Phi(U_0 - \delta; \lambda^*, \alpha_0).$$

Let consider the value

$$\alpha > \max\left(\alpha_0, \frac{4\lambda^*}{V_0}\right).$$

Noting that  $U_0 < p$  and that  $0 > H_{\lambda^*}(p - U) \geq H_{\lambda^*}(0) = -\lambda^*$  for  $U \in [U_0 - \delta, U_0]$ , we have

$$\begin{aligned} \frac{d\Phi}{dU}(U; \lambda^*, \alpha) &= \alpha + \frac{H_{\lambda^*}(p - U)}{\Phi(U; \lambda^*, \alpha)} \\ &\geq \alpha + \frac{2H_{\lambda^*}(p - U)}{V_0} \end{aligned}$$

$$\begin{aligned} &> \alpha - \frac{2\lambda^*}{V_0} \\ &> \frac{1}{2}\alpha \end{aligned}$$

for  $U \in [U_0 - \delta, U_0]$ . Apply the mean value theorem to obtain

$$\begin{aligned} \Phi(U; \lambda^*, \alpha) &= \Phi(U; \lambda^*, \alpha) - \Phi(U_0 - \delta; \lambda^*, \alpha) + \Phi(U_0 - \delta; \lambda^*, \alpha) \\ &> \frac{1}{2}\alpha \{U - (U_0 - \delta)\} + \frac{1}{2}V_0 \end{aligned}$$

for  $U \in [U_0 - \delta, U_0]$  and in particular

$$\Phi(U_0; \lambda^*, \alpha) > \frac{1}{2}\alpha\delta + \frac{1}{2}V_0.$$

If necessary, we can take  $\alpha$  larger and satisfying

$$\frac{1}{2}\alpha\delta + \frac{1}{2}V_0 > \frac{\lambda^*}{\alpha} = \Psi_{\alpha, \lambda^*}(p)$$

so that

$$\Phi(U_0; \lambda^*, \alpha) > \Psi_{\alpha, \lambda^*}(p).$$

Finally we obtain

$$\{(U, V) \in \mathbb{R}^2 \mid V = \Phi(U; \lambda^*, \alpha), U_0 < U < p\} \subset W_1,$$

and thus

$$V_0 < \Phi(p; \lambda^*, \alpha) = 0,$$

which contradicts the fact  $V_0 > 0$ .  $\square$

**Proof of Theorem 6**  $\alpha^*(\lambda)$  is the desired threshold with respect to  $\alpha$  by Lemma 11. Since  $\alpha^*(\lambda)$  is strictly increasing and continuous by Lemmas 12 and 13 respectively, the inverse function of  $\alpha^*(\lambda)$ , denoted by  $\lambda(0, 0)(\alpha)$ , is well-defined for  $\alpha \geq 0$  and inherits the properties of monotonicity and continuity: Remark 5 and Lemma 14 complete the proof.  $\square$

**Acknowledgements** This work was motivated and started while the second named author was visiting the Riemann International School of Mathematics at the University of Insubria in Varese - Italy, during September 2017. He would like to express his deepest gratitude for the warm hospitality.

**Funding** Open access funding provided by Università degli Studi dell'Insubria within the CRUI-CARE Agreement.

**Data availability** Not applicable.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Cassani, D., Kaltenbacher, B., Lorenzi, A.: Direct and inverse problems related to MEMS. *Inverse Problems* **25**(10), 105002 (2009)
2. Cassani, D., Fattorusso, L., Tarsia, A.: *Nonlocal Dynamic Problems with Singular Nonlinearities and Applications to MEMS. Progress in Nonlinear Differential Equations and their Applications*, vol. 85, pp. 187–206. Birkhäuser, Boston (2014)
3. Chipot, M., Weissler, F.B.: Some blowup results for a nonlinear parabolic equation with a gradient term. *SIAM J. Math. Anal.* **20**(4), 886–907 (1989)
4. Conley, C.C.: *Isolated Invariant Sets and the Morse Index*. CBMS Regional Conference Series in Mathematics, vol. 38. American Mathematical Society, Providence (1978)
5. Flores, G.: On the dynamic pull-in instability in a mass-spring model of electrostatically actuated MEMS devices. *J. Differ. Equ.* **262**(6), 3597–3609 (2017)
6. Guo, J.-S., Kavallaris, N.I.: On a nonlocal parabolic problem arising in electrostatic MEMS control. *Discret. Contin. Dyn. Syst.* **32**(5), 1723–1746 (2012)
7. Haraux, A.: Sharp decay estimates of the solutions to a class of nonlinear second order ODE's. *Anal. Appl. (Singap.)* **9**(1), 49–69 (2011)
8. Haraux, A., Jendoubi, M.A.: *The Convergence Problem for Dissipative Autonomous Systems*. SpringerBriefs in Mathematics. Springer/BCAM Basque Center for Applied Mathematics, Cham/Bilbao (2015)
9. Kavallaris, N.I., Lacey, A.A., Nikolopoulos, C.V., Tzanetis, D.E.: On the quenching behaviour of a semilinear wave equation modelling MEMS technology. *Discret. Contin. Dyn. Syst.* **35**(3), 1009–1037 (2015)
10. Peletier, L.A., Serrin, J., Zou, H.: Ground states of a quasilinear equation. *Differ. Integral Equ.* **7**(3–4), 1063–1082 (1994)
11. Souplet, P.: Critical exponents, special large-time behavior and oscillatory blow-up in nonlinear ODE's. *Differ. Integral Equ.* **11**(1), 147–167 (1998)
12. Versaci, M., Di Barba, P., Morabito, F.C.: Curvature-dependent electrostatic field as a principle for modelling membrane-based MEMS devices. A review. *Membranes* **10**(11), 361 (2020)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.