Correction to: Natural habitat and vegetation types of river gravel bars in the Caucasus Mountains, Georgia

Veronika Kalníková · Kryštof Chytrý · Pavel Novák · Dominik Zukal · Milan Chytrý

Published online: 8 November 2020
© Institute of Botany, Czech Academy of Sciences 2020

Correction to: Folia Geobot (2020) 55:41–62
<https://doi.org/10.1007/s12224-020-09364-6>

Please be informed that the wrong table was inadvertently published in the following paper: "Natural habitat and vegetation types of river gravel bars in the Caucasus Mountains, Georgia", DOI number 10.1007/s12224-020-09364-6 by Veronika Kalníková Kryštof Chytrý, Pavel Novák, Dominik Zukal & Milan Chytrý, published in Folia Geobotanica Volume 55, issue 1 (June 2020). Please find the correct table 2 below.

Springer wishes to apologise for any inconvenience caused.

The online version of the original article can be found at
<https://doi.org/10.1007/s12224-020-09364-6>

V. Kalníková (✉) · K. Chytrý · P. Novák · D. Zukal ·
M. Chytrý
Department of Botany and Zoology, Masaryk University,
Kotlářská 2, CZ-611 37 Brno, Czech Republic
e-mail: v.kalnikova@seznam.cz

V. Kalníková
Beskydy Protected Landscape Area Administration, Nádražní 36,
CZ-756 61 Rožnov pod Radhoštěm, Czech Republic

Vegetation type	Tus-Cal	Epi	Pet-Cre	Sal-M	Sal-Hip
Number of relevés	8	23	14	19	6
<i>Tussilagini farfarae-Calamagrostietum pseudophragmitae</i>					
<i>Equisetum arvense</i>	50	9	7	5	.
<i>Juncus articulatus</i>	38	.	.	5	17
<i>Calamagrostis pseudophragmites</i>	100	70	.	89	50
<i>Plantago major</i>	50	4	14	37	.
<i>Epilobietum colchici</i>					
<i>Senecio leucanthemifolius</i> subsp. <i>caucasicus</i>	.	30	.	5	.
<i>Poa alpina</i>	25	65	.	42	.
<i>Racomitrium canescens</i> (B)	13	52	21	16	.
<i>Rumex acetosella</i>	.	48	14	11	17
<i>Pinus sylvestris</i>	.	26	.	11	.
<i>Pilosella officinarum</i>	.	26	.	11	.
<i>Petrorhagia saxifraga-Crepis foetida</i> community					
<i>Trifolium arvense</i>	.	9	64	16	17
<i>Plantago lanceolata</i>	13	4	100	42	83
<i>Crepis foetida</i>	.	9	79	26	50
<i>Silene compacta</i>	.	17	50	11	.
<i>Petrorhagia saxifraga</i>	.	13	86	32	67
<i>Arenaria serpyllifolia</i> agg.	.	22	79	37	33
<i>Barbula unguiculata</i> (B)	.	4	43	.	17
<i>Daucus carota</i>	.	.	29	5	.
<i>Petrorhagia prolifera</i>	.	.	21	.	.
<i>Echium vulgare</i>	.	.	57	21	33
<i>Syntrichia ruralis</i> (B)	.	4	21	.	.
<i>Salici purpureae-Myricarietum germanicae</i>					
<i>Myricaria germanica</i>	.	22	.	89	.
<i>Trisetum rigidum</i>	.	22	.	58	.
<i>Gypsophila elegans</i>	.	13	.	47	.
<i>Equisetum variegatum</i>	.	.	.	32	.
<i>Silene ruprechtii</i>	.	4	.	26	.
<i>Lotus corniculatus</i>	13	.	14	42	.
<i>Vicia sosnowskyi</i>	.	.	.	16	.
<i>Cirsium echinus</i>	.	13	14	37	.
<i>Salvia verticillata</i>	.	4	7	26	.
<i>Salici incanae-Hippophaëtum rhamnoidis</i>					
<i>Hippophaë rhamnoides</i>	.	9	7	21	100
<i>Sonchus oleraceus</i>	.	4	.	.	33
<i>Paracynoglossum glochidiatum</i>	.	9	21	.	50
<i>Verbascum sessiliflorum</i>	.	.	7	.	33
<i>Elytrigia repens</i>	.	.	7	.	33
<i>Hypochaeris radicata</i>	.	.	7	.	33
<i>Poa pratensis</i>	.	.	7	5	33
<i>Catapodium rigidum</i>	.	.	14	.	33
<i>Alnus glutinosa</i>	13	.	7	.	33
<i>Erigeron canadensis</i>	13	9	64	11	67
<i>Epilobium colchicum</i>	25	87	7	74	.

Vegetation type	Tus-Cal	Epi	Pet-Cre	Sal-M	Sal-Hip
Number of relevés	8	23	14	19	6
Other species occurring in at least 20% of relevés					
<i>Setaria viridis</i>	.	.	29	.	17
<i>Salix caprea</i>	.	13	14	26	.
<i>Medicago minima</i>	.	.	29	.	33
<i>Euphorbia maculata</i>	.	.	29	.	33
<i>Bidens frondosus</i>	25	.	.	.	17
<i>Phalaroides arundinacea</i>	25	.	.	.	17
<i>Anisantha tectorum</i>	.	4	29	.	17
<i>Achillea millefolium</i> agg.	.	9	21	21	.
<i>Sedum spurium</i>	13	22	14	11	.
<i>Tussilago farfara</i>	25	17	7	21	.
<i>Alnus incana</i>	.	17	.	21	.
<i>Sedum pallidum</i>	.	26	29	5	.
<i>Ceratodon purpureus</i> (B)	.	17	29	11	.
<i>Vulpia myuros</i>	.	.	21	.	17
<i>Bromus japonicus</i>	.	4	29	11	17
<i>Papaver fugax</i>	.	17	.	21	.
<i>Seseli transcaucasicum</i>	.	13	.	21	.
<i>Prunella vulgaris</i>	.	13	14	26	17
<i>Agrostis stolonifera</i> agg.	25	4	43	37	17
<i>Bryum caespitium</i> (B)	13	17	14	26	17
<i>Sedum album</i>	.	4	21	.	17
<i>Ranunculus repens</i>	25	.	7	16	17
<i>Bryum argenteum</i> (B)	.	22	29	.	17
<i>Equisetum ramosissimum</i>	25	.	7	.	33
<i>Tanacetum parthenium</i>	.	13	36	16	17
<i>Erigeron acris</i>	.	39	7	37	.
<i>Cerastium fontanum</i>	.	26	7	11	17
<i>Trifolium repens</i>	38	9	36	42	33
<i>Ambrosia artemisiifolia</i>	.	.	21	.	33
<i>Trifolium pratense</i>	13	17	7	37	.
<i>Poa compressa</i>	.	.	21	26	.
<i>Leucanthemum vulgare</i>	13	4	21	53	33
<i>Tortella inclinata</i> (B)	.	9	.	21	.
<i>Medicago lupulina</i>	25	17	57	63	17
<i>Poa nemoralis</i>	13	35	.	16	.
<i>Artemisia absinthium</i>	.	13	7	32	17
<i>Salix alba</i>	25	9	14	47	33
<i>Polytrichum piliferum</i> (B)	.	22	.	11	.
<i>Tripleurospermum caucasicum</i>	13	26	.	5	.
<i>Salix purpurea</i>	38	13	7	42	50
<i>Brachythecium rivulare</i> (B)	25	.	7	.	33