Mesanthemum tuberosum Lecomte resurrected from *M. prescottianum* (Bong.) Körn. (Eriocaulaceae), variation and lectotypification

Sylvia M. Phillips¹, Fatoumata Fofana² & Martin Cheek¹

Summary. *Mesanthemum prescottianum* is a West African species occurring in the uplands extending from Guinea to Ivory Coast. Variation within the species is discussed and the synonym *M. tuberosum* Lecomte is raised to specific status. The conservation status of the first species is assessed using the IUCN 2012 standard as Least Concern, and the second as Vulnerable. Lectotypes are selected for both names. The type location of *M. prescottianum*, claimed to be Brazil, is shown to be an error.

Key Words. Brazil, lectotype, taxonomy, West Africa.

Mesanthemum is a genus of c. 15 species occurring in Africa, with two in Madagascar (Phillips 2011). It is relatively uniform, the majority of species being perennials from a tough rootstock or rhizome, and is more vigorous than all except the largest species of *Eriocaulon* in Africa. *Eriocaulon* is a pantropical genus with many more species in Africa than *Mesanthemum*. *Mesanthemum* lacks the fenestrate leaves found in *Eriocaulon*, and can also be readily distinguished from nearly all African species of that genus by its hairy leaves and scapes (Phillips 2011). An early division in keys to *Mesanthemum* is the easily observed character as to whether the inner involucre bracts radiate beyond the central disc of flowers, thus resembling the ray florets of a member of the Asteraceae (Jacques-Félix 1947; Meikle 1968; Phillips 2010). This character is well expressed in the West African species *M. prescottianum* (Bong.) Körn., where the white capitula are often described as “daisy-like”. The only other species with similar daisy-like capitula is *M. bennae* Jacq.-Fél., also from Guinea but known only from cliffs of Mt Benna, at Kounounkan. It is easily distinguished from both *M. prescottianum* and *M. tuberosum* by its stouter habit, much broader leaves (c. 20 mm), broader velutinous involucre bracts and pilose floral bracts.

A recent field trip to the Fouta Djallon upland region of Guinea (Conakry) by the third author as part of the Tropical Important Plant Areas programme (Darbyshire, continuously updated; Darbyshire *et al.* 2017) has resulted in a reassessment of variation in *Mesanthemum prescottianum*. This species

is confined to uplands in the West African countries of Guinea, Sierra Leone, Liberia and Ivory Coast. It is known to be rather variable, as noted by Jacques-Félix (1947: 144), who reduced *M. tuberosum* Lecomte from Fouta Djallon to synonymy within *M. prescottianum*. This treatment was followed by Meikle in *Flora of West Tropical Africa* (1968: 64).

Mesanthemum prescottianum is a distinctive species in the field due to its conspicuous shining white “daisy” capitula with sharply acute, spreading involucre bracts. Material collected in Fouta Djallon looks different in the field, with duller, whitish capitula, narrower, obtuse involucre bracts and scapes arising from well-developed underground corms (Fig. 1), at first leading to the suspicion that that it could be an undescribed species. This population was subsequently found to match the type of *M. tuberosum*. A detailed examination of all the material in the Kew herbarium showed the Fouta Djallon material to be sufficiently distinct from *M. prescottianum*, both morphologically and in habitat preference, to merit specific status. *Mesanthemum tuberosum* is therefore resurrected here. As the *Flora of West Tropical Africa* does not include descriptions, a full description of both species is given.

This paper is part of a series of studies on African *Eriocaulaceae* by the first author. Recent discoveries of W African members of the family include Phillips & van der Burgt (2012) and Phillips & Mesterházy (2015).

Following decades of relative inactivity in Republic of Guinea, in the last seven years there has been a dramatic rise in the publication of new discoveries in other families. Examples are: *Brachystephanus*

Fig. 1. *Mesanthemum tuberosum* A habit, small plant showing corm; B habit, tall plant; C scape indumentum; D leaf indumentum, adaxial surface, left margin inrolled; E inflorescence, side-view, hydrated, from A; F outer bract; G inner bract; H transitional bract; J floral bract; K receptacular hair showing slightly clavate distal portion; L detail of 2-celled leaf hair; M young male flower, lateral view; N older male flower, lateral view, all anthers fallen; P female flower, lateral view; Q male flower sepals, apices split, midvein present; R female flower sepals, left and right are keeled, central sepal is flat; S male corolla cut and opened out to show stamens (part reconstructed); T female corolla, lateral view; U female corolla cut and opened showing pistil. All from Cheek 18489 (K). DRAWN BY ANDREW BROWN.

oreacanthus Champl. (Champluvier & Darbyshire 2009), *Eriocaulon cryptocephalum* S. M. Phillips & Mesterházy (2015), *Eriosema triformum* Burgt (van der Burgt *et al.* 2012), *Gymnosiphon samoritourenus* Cheek (Cheek & van der Burgt 2010), *Inversodicraea pepehabai* Cheek (Cheek & Haba 2016), *Napoleonea alata* Jongkind (Prance & Jongkind 2015), *Psychotria samoritourei* Cheek (Cheek & Williams 2016), *Striga magnibracteata* Eb. Fisch. & I. Darbysh. (Fischer *et al.* 2011) and *Xysmalobium samoritourei* Goyder (2009). Just over the border in Mali, *Calophyllum africanum* Cheek & Q. Luke (2016) was recently found and in Ivory Coast, *Macropodiella cussetiana* Cheek (Cheek & Ameka 2016). Even a new rheophytic genus, *Karima* Cheek & Riina has come to light in Guinea (Cheek *et al.* 2016).

Mesanthemum prescottianum (Bong.) Körn. (Körnische 1863: 472, t. 60 f. 1); Durand & Schinz (1895: 504); N. E. Brown (1901: 261); Ruhland (1903: 118); Meikle (1968: 327).

Eriocaulon prescottianum Bong. (Bongard 1831: 635).

Type: Sierra Leone ("Brasilia", see below), com. Lindley s.n. (sphalm. Lindly), 1828 (LE lectotype [LE 00001102], designated here; K, isoelectotype (K000346197)).

Perennial from a slightly swollen base or with a small corm, flowering towards end of wet season (late July – Oct.); sometimes producing small tubers on horizontal stolons; roots dense, tough, tightly attached to substrate. *Leaves* up to 5, up to 35 cm long, 2.6 – 6 mm wide, linear or rarely ensiform, flat or lightly folded, outwardly curving when short, pilose especially on upper surface and margins towards base, tip acute. *Scapes* 1 – 4, 25 – 50 (– 67) cm high, slender, pilose. *Sheaths* pilose, limb acuminate, glabrous or sparsely pilose within. *Capitulum* shiny white, (17 –) 25 – 35 mm wide, flat-topped, greenish crateriform involucre at base. *Involucral bracts* in several series of increasing length, outer bracts lanceolate-oblong, 2.5 – 7 mm long, leathery with thinner margins, pilose with soft dense hairs, acute, inner bracts shiny white, narrowly lanceolate-oblong, radiating ray-like and much exceeding floral disc, 9 – 18 mm long, thickly papery, outer (lower) surface pubescent, strongly many-veined, veins appearing as furrows on upper surface, acute. *Floral bracts* fusiform above middle, filiform below, upper part fleshy, glabrous, acute. *Receptacle* flat, densely villous, hairs usually colourless, occasionally pale brownish-grey, rarely blackish. *Flowers* 2 – 2.5 mm long. *Male flowers*: sepals narrowly oblong, free, concave, glabrous, broadly obtuse or minutely denticulate; petal-tube shallowly 3-lobed, upper margin densely villous, pilose inside; anthers dark brown or

black. *Female flowers*: sepals resembling male sepals; petals free below middle, connate above, connate portion with a band of hairs outside from upper margins of free petal-bases, some hairs inside around 3 glands, upper margin shallowly 3-lobed and densely villous. Fig. 2A & B

DISTRIBUTION. Guinea (Conakry), Ivory Coast, Liberia, Sierra Leone.

SPECIMENS EXAMINED (Three selected from each country where more are available). **GUINEA (CONAKRY).** Beyla Distr., Simandou Ridge, 8 Nov. 2005, Cheek 12665 (K, HNG); Nzérékoré, Nimba Mts, near pumping station Zougoué R., 24 July 2008, Jongkind 8265 (K, WAG); Macenta Prefecture, Ziamia Mts, Mt Wokou just S of Macenta, 24 Oct. 2008, Pearce 22 (K, HNG, WAG). **IVORY COAST.** Mt Tonkoui, Aug. 1954, Schnell 6363 (K, P); Nimba Mts, 10 Aug. 1954, Boughey G.C.18053 (K). **LIBERIA.** Central Province, Sanokwele Distr., near Sanokwele, 26 Sept. 1947, Baldwin 9516 (K, NY, S, M). **SIERRA LEONE.** Makeni Hill, 26 Aug. 1951, Jordan 499 (K); Tonkolili Distr., S Sula Mts, NE of Farangbaaya, 12 Sept. 2009, Kanu 6 (FBC, K, SL); N of Makeni on Kalaba road, 21 Aug. 1967, Morton 4534 (K, SL).

HABITAT. Thin peaty soil over exposed rock with seasonal seepages and in rocky crevices, especially on granite outcrops, often growing with grasses and species of Cyperaceae; 630 – 1600 m.

CONSERVATION STATUS. *Mesanthemum prescottianum* is here assessed as Least Concern (LC), using the IUCN (2012) Standard. There are about 30 global locations where it is widespread and frequent, and at most of them it is not threatened. The extent of occurrence was measured as 82,953 km² using Geocat (Bachman *et al.* 2011). At Tonkolili, Simandou and Nimba it is threatened by exploration for exploitation of iron ore by open-cast mining.

NOTES. *Mesanthemum prescottianum* has a restricted distribution in the far west of Africa, where it is known only from thin soil over rock, to which it is very tightly affixed. It varies considerably in vigour, capitulum size, hairiness of the involucral bracts, and the colour of the hairs in the capitulum. However, it is always immediately recognisable by its shiny white daisy-like capitula with long, ray-like, usually pointed involucral bracts. The dark anthers of this species and *M. tuberosum* appear to be unique in the genus, other species having creamy-white or yellowish anthers.

Collections from Sierra Leone and Liberia have capitula with colourless hairs on and around the flowers, so the floral disc is pale. These hairs are light grey-brown in collections seen from Ivory Coast and southern Guinea, giving the floral disc a somewhat darker appearance. A striking variant from the Nimba Mountains in southern Guinea has very large capitula

Fig. 2. A, B *Mesanthemum prescottianum*: A in habitat (Haba 216); B capitulum (Kanu 6). C – E *M. tuberosum*: C capitulum (Cheek 18489); D corm (Cheek 18489); E in habitat (Cheek 18489). PHOTOS: A & B X. VAN DER BURGT, C – E M. CHEEK.

(35 mm wide), almost blackish receptacular hairs and copper-brown sepals, so the capitula have very dark centres (Jongkind 8265 & 8266).

Mesanthemum prescottianum flowers late in the wet season, reaching peak flowering in August and September and seeding in late October and November,

the dried capitula persisting on the plant. It typically grows directly on rock in seasonally wet places, never in permanent standing water or marsh.

TYPIIFICATION. Bongard was based in St Petersburg (LE). He states in the protologue that he received his specimen from Lindley who had it from the herbarium of D. Prescott. There is a good specimen in LE (LE00001102) with a handwritten label, presumably by Bongard, stating “*Eriocaulon prescottianum* Bogn! Brasilia, comm. Lindley 1828”. Bongard’s protologue also includes a reference to a plate (t. 36) but this was never published, as noted by Körnicke (1863) when he transferred the species to *Mesanthemum*. One of the plants on the sheet is the basis of t. 36, as indicated by a pencil note.

In the protologue Bongard (1831) gives the provenance of his collection as Rio de Janeiro and the specimen label states “Brasilia”. This locality information was repeated by Körnicke who was following Bongard. This appears to be a mistake as *Mesanthemum prescottianum* is confined to West Africa. Kew also has a specimen of *M. prescottianum* received from Herb. Lindley, stamped “Herb. Benth.” and with the handwritten name *M. radicans*. It is probably a duplicate of Bongard’s specimen which was sent to Bentham, but has the locality “Sierra Leone”. N. E. Brown (1901: 262) remarks that Sierra Leone is much more likely to be the correct origin rather than Brazil. Ruhland (1903) states that other specimens sent out by Lindley have the Sierra Leone locality.

The specimen in St Petersburg is much better than the one at Kew and is the specimen which Bongard used for his protologue. There is no specimen of *Mesanthemum prescottianum* in either the Cambridge (CGE) or Oxford (OXF) herbaria. The LE duplicate is therefore selected here to be the lectotype, despite the wrong locality information.

Mesanthemum tuberosum Lecomte, *Bull. Soc. Bot. France* 55: 598 (1908 publ. 1909). Type: Guinea (Conakry), Kouroussa to Timbo, June 1907, *Pobéguin* 734 (P lectotype [P00443181], designated here).

Perennial from a broad-based corm buried 2 cm or more deep, flowering at start of wet season (June – July); corm well developed, covered in brown fibres. *Leaves* 1 – 3, 5 – 26 cm long, 2.4 – 10 mm wide, flat or lightly folded, longer leaves linear, shorter leaves narrowly lanceolate and outwardly curving from sheathing base, pilose with patent hairs especially towards base and on margins, tip hard, subacute. *Scapes* 1 – 3, up to 33 cm high, densely pilose with patent hairs. *Sheaths* tightly cylindrical, opening abruptly into flattened acuminate limb. *Capitulum*

dull white, 12 – 25 (– 30) mm wide, flat topped, green or brown crateriform involucre at base. *Involucral bracts* in several series of increasing length, outer bracts broadly lanceolate, 5 – 7 mm long, leathery in lower third, upper part and margins herbaceous, softly pilose, abruptly narrowed above, inner progressively more caudate, becoming broadly linear, extending ray-like beyond floral disc, dull white, fleshy, 3 – 5-veined, obtuse or subacute, innermost shorter and grading into floral bracts, all except outermost subtending flowers. *Floral bracts* broadly linear to clavate, contracted to filiform base, fleshy, white, glabrous, obtuse or subacute. *Receptacle* flattened, flowers and slender bases of floral bracts embedded in dense cushion of grey hairs, fleshy white tips of floral bracts protruding above. *Flowers* 2.5 – 3.5 mm. *Male flowers*: sepals elliptic-oblong to obovate, unequally connate at base, concave, glabrous, obtuse; petal-tube shallowly 3-lobed, upper margin densely villous with grey-brown hairs, glabrous inside; anthers dark brown. *Female flowers*: sepals resembling male sepals; petals free in lower third, connate above, villous in 3 longitudinal bands along seams of upper connate portion, glabrous inside, upper margin shallowly 3-lobed and densely villous with grey-brown hairs. Figs 1, 2C – E

DISTRIBUTION. Guinea (Conakry), Fouta Djallon Highlands, distributed in the highest part of the range in an isosceles triangle which has at its corners the towns of Mali, Mamou and Gaoual. Map. 1.

SPECIMENS EXAMINED. GUINEA (CONAKRY). FOUTA DJALON: Dalaba, 10°37'41.9"N 12°16'46.2"W, 10 June 2016, *Cheek* 18489 & near Forestry House, 21 June 1962, *Adames* 273 (K); Labe, between Labe and Pita on the road c. 15 km from Pita, 11°10'47.9"N 12°22'51.3"W, 12 June 2016, *König* 30 (K, HNG); 5 km S of Sériba, 11°4'N 13°1'W, 28 July 1992, *Porembski* 44 (K); 40 km W of Labé, 11°2'N 12°4'W, 29 July 1992, *Porembski* 64; 20 km N of Dalaba, 10°4'N 12°2'W, 30 July 1992, *Porembski* 84 (K); 20 km S of Mamou, 10°1'N 12°3'W, 31 July 1992, *Porembski* 97 (K); Mamou, Sept. 1954, *Schnell* 6789 (K); Mali, Sept. 1954, *Schnell* 7063 (K).

HABITAT. Sandstone bowal with stepped horizontal flats covered in sandy soil; 900 – 1500 m.

CONSERVATION STATUS. We here assess *Mesanthemum tuberosum* using the categories and criteria of IUCN (2012) as Vulnerable, VU B2ab(iii), since nine threat-based locations exist (see Map 1), the area of occupancy is 36 km² using the preferred IUCN 4 km² cell size, and because threats have been observed at several locations from cattle grazing and trampling to modification of habitat for cultivation (pers. obs. Cheek June 2016). A poster campaign in the Fouta Djallon is intended to raise awareness of the

Map 1. Distribution of *Mesanthemum tuberosum* in Guinea (Conakry).

importance of conserving this threatened rare and endemic species.

NOTES. In dissected specimens the female flowers were few in the axils of the involucre bracts, flowers of the central floral disc being apparently all male.

This taxon appears to be an extreme variant from the *Mesanthemum prescottianum* genepool, confined to the Fouta Djallon area of Guinea. It was described by Lecomte as a new species *M. tuberosum*, but sunk into synonymy by later authors. It differs from

M. prescottianum by the characters set out in Table 1 below. It also occupies a different habitat, which reinforces the decision here to recognise it again at specific rank.

TYPIIFICATION. Lecomte (1909) cites three specimens in his protologue, all from the Fouta Djallon upland in Guinea (Conakry): *Pobéguin* 734 (Kouroussa to Timbo); *Caille* in *Chevalier* 15038 (Bilima); *Chevalier* 18389 (between Timbo and Ditinn). The specimen *Pobéguin* 734 [P00443181] is labelled as holotype. It is

Table 1. Comparison of the main differences between *Mesanthemum prescottianum* and *M. tuberosum*.

	<i>Mesanthemum prescottianum</i>	<i>Mesanthemum tuberosum</i>
Habitat	rocky outcrops with seeping surface water in wet season	sandy flats overlying impervious rock, damp in wet season
Flowering season	mid – late wet season late July – Oct.	early wet season June - July
Base of culm	slender, 2 – 3 mm thick, tightly attached to rock by dense network of tough roots	stout, c. 10 mm thick, covered in brown fibres, roots not anchored to rock
Vegetative reproduction	Small white tubers produced on horizontal stolons	Stolons not seen, rosette sometimes dividing
Capitulum	shiny white longest involucre bracts acute	dull white longest involucre bracts obtuse

a good specimen including a label with collection details and a capsule with dissections. It is therefore selected here as lectotype. A second sheet of *Pobéguin* 734 [P00443180] is an isoelectotype.

Acknowledgements

The third author's fieldwork costs for this paper were supported by the Darwin initiative DEFRA project 23–002 “Tropical Important Plant Areas of Guinea”. Charlotte Couch, Natalie König and Fatoumata Fofana supported the fieldwork that led to the rediscovery of *Mesanthemum tuberosum*. The curators of the herbaria in Cambridge (CGE) and Oxford (OXF) are thanked for searching for possible material of *M. prescottianum* originating from Prescott's herbarium. George Gosline produced the map. Saba Rokni generated the Geocat files that allowed calculations for the IUCN assessments.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Bachman, S., Moat, J., Hill, A. V., De la Torre, J. & Scott, B. (2011). Supporting Red List threat assessments with GeoCAT, a geospatial conservation assessment tool. *Zookeys* 150: 117 – 126.
- Bongard, M. (1831). Essai monographique sur les espèces d'*Eriocaulon* du Brésil. *Mém. Acad. Imp. Sci. St.-Petersbourg. Sér. 6, Sci Math* 1: 601 – 655.
- Brown, N. E. (1901). *Eriocaulaceae*. In: W. Thiselton-Dyer (ed.), *Flora of Tropical Africa* Vol. 8: 230 – 264. Lovell Reeve & Co., London.
- van der Burgt, X. M., Haba, P. K., Haba, P. M. & Goman, A. S. (2012). *Eriosema triformum* (Leguminosae: Papilionoideae), a new unifoliate species from Guinea, West Africa. *Kew Bull.* 67 (2): 263 – 271. <https://doi.org/10.1007/s12225-012-9357-5>
- Champluvier, D. & Darbyshire, I. (2009). A revision of the genera *Brachystephanus* and *Oreacanthus* (Acanthaceae) in tropical Africa. *Syst. Geogr. Pl.* 79 (2): 115 – 192.
- Cheek, M. & Ameka, G. (2016). *Macropodiella cussetiana* (Podostemaceae) a new species from Côte d'Ivoire. *Kew Bull.* 71: 21. <https://doi.org/10.1007/S12225-016-9634-9>.
- & van der Burgt, X. (2010). *Gymnosiphon samoritourenus* (Burmanniaceae) a new species from Guinea, with new records of other achlorophyllous heteromycotrophs. *Kew Bull.* 65: 83 – 88. <https://doi.org/10.1007/s12225-010-9180-9>
- , Challen, G., Lebbie, A., Banks, H., Barberá, P. & Riina, R. (2016). Discovering *Karima* (Euphorbiaceae) a New Crotonoid Genus from West Tropical Africa Long Hidden within *Croton*. *PLOS ONE*. <https://doi.org/10.1371/journal.pone.0152110>
- & Haba, P. (2016). *Inversodicraea* Engl. resurrected and *I. pepehabai* sp. nov. (Podostemaceae), a submontane forest species from the Republic of Guinea. *Kew Bull.* 71: 55. <https://doi.org/10.1007/S12225-016-9673-2>
- & Luke, Q. (2016). *Calophyllum* (Clusiaceae – Guttiferae) in Africa. *Kew Bull.* 71: 20. <https://doi.org/10.1007/S12225-016-9637-6>
- & Williams, T. (2016). *Psychotria samoritourei* (Rubiaceae), a new liana species from Loma-Man in Upper Guinea, West Africa. *Kew Bull.* 71: 19. <https://doi.org/10.1007/S12225-016-9638-5>
- Darbyshire, I. (continuously updated). Tropical Important Plant Areas. <http://science.kew.org/strategic-output/tropical-important-plant-areas>
- , Anderson, S., Asatryan, A., Byfield, A., Cheek, M., Clubbe, C., Ghrabi, Z., Harris, T., Heatubun, C. D., Kalema, J., Magassouba, S., McCarthy, B., Milliken, W., Montmollin, B. de, Nic Lughadha, E., Onana, J. M., Saidou, D., Sarbu, A., Shrestha, K. & Radford, E. A. (2017). Important Plant Areas: revised selection criteria for a global approach to plant conservation. *Biodivers. Conserv.* 26: 1767 – 1800. <https://doi.org/10.1007/s10531-017-1336-6>.
- Durand, T. & Schinz, H. (1895). *Conspectus Florae Africae* 5. Jardin Botanique de l'Etat, Bruxelles, Berlin.
- Fischer, E., Darbyshire, I. & Cheek, M. (2011). *Striga magnibracteata* (Orobanchaceae) a new species from Guinée and Mali. *Kew Bull.* 66: 441 – 445. <https://doi.org/10.1007/s12225-011-9296-6>
- Goyder, D. J. (2009). *Xysmalobium samoritourei* (Apocynaceae: Asclepiadoideae), a new species from the Guinea Highlands of West Africa. *Kew Bull.* 63: 473 – 475. <https://doi.org/10.1007/s12225-008-9059-1>
- Jacques-Félix, H. (1947). Le genre *Mesanthemum* (Eriocaulaceae). *Bull. Soc. Bot. France* 94: 143 – 151.
- IUCN (2012). *IUCN Red List Categories and Criteria: Version 3.1*. Second edition. IUCN, Gland and Cambridge.
- Körnigke, F. (1863). Eriocaulaceae. In: C. F. Ph. von Martius & A. W. Eichler (eds), *Flora Brasiliensis*, Vol. 3 (1): 274 – 507. Typographia regia, Monachii.
- Lecomte, H. (1909). Eriocaulacées d'Afrique. *Bull. Soc. Bot. France* 55: 594 – 602.

- Meikle, R. D. (1968). *Eriocaulaceae*. In: F. N. Hepper (ed.), *Flora of West Tropical Africa*, Ed. 2, 3: 57 C67. Crown Agents, London.
- Phillips, S. M. (2010). *Eriocaulaceae*. In: J. R. Timberlake & E. S. Martins (eds), *Flora Zambesiaca*, 13 (4): 1 – 151. Royal Botanic Gardens, Kew.
- ____ (2011). *Eriocaulaceae*. In: *Flore du Cameroun* 38: 1 – 35. Ministry of Scientific Research & Innovation, Cameroon.
- ____ & van der Burgt X. M. (2012). Two new species of *Eriocaulon* (Eriocaulaceae) from Sierra Leone. *Kew Bull.* 67: 273 – 280.
- ____ & Mesterházy, A. (2015). Revision of small ephemeral species of *Eriocaulon* (Eriocaulaceae) in West Africa with long involucre bracts. *Kew Bull.* 70: 5 (17 pp). <https://doi.org/10.1007/S12225-014-9557-2>
- Prance, G. T. & Jongkind, C. C. H. (2015). A revision of African Lecythidaceae. *Kew Bull.* 70: 6. <https://doi.org/10.1007/s12225-014-9547-4>.
- Ruhland, W. (1903). *Eriocaulaceae*. In: A. Engler (ed.), *Das Pflanzenreich* Vol. 13: 1 – 294 (IV – 30), Wilhelm Engelmann, Leipzig.