Senecio scapioides (Compositae: Senecioneae: Senecioniinae): a new species from the Departamento de Boyacá, in Andean Colombia

José Aguilar-Cano¹ & D. J. Nicholas Hind²

Summary. Routine use of the virtual herbarium of the collections in COL (Instituto de Ciencias Naturales de la Universidad Nacional de Colombia) revealed the presence of an as yet undescribed species of *Senecio* amongst the material of *Senecio adglacialis* and *S. leucanthemoides*. Following subsequent herbarium studies and fieldwork, *Senecio scapioides* (Compositae: Senecioneae: Senecioniinae), from the Departamento de Boyacá, in Andean Colombia, was recognised and is described and illustrated; it is compared with *S. adglacialis* and *S. leucanthemoides*. Notes on its distribution and habitat, conservation status, phenology, and etymology are also provided, and the distribution of the three species mapped.

Resumen. Revisiones rutinarias del herbario virtual de las colecciones en COL (Instituto de Ciencias Naturales de la Universidad Nacional de Colombia) reveló la presencia de una especie aún no descrita de *Senecio* entre el material de *Senecio adglacialis* y *S. leucanthemoides*. Como resultado del trabajo de herbario y campo subsiguientes, *Senecio scapioides* (Compositae: Senecioneae: Senecioniinae), del Departamento de Boyacá, en los andes colombianos, fue reconocida e ilustrada; la nueva especie se compara con *S. adglacialis* y *S. leucanthemoides*. Adicionalmente se discute su distribución y hábitat, estado de conservación, fenología y etimología, así como el mapa de distribución geográfica de las tres especies.

Key Words. Asteraceae, checklist, digitisation, Flora of Colombia.

Introduction

In 2015, one of the authors (DJNH) reassessed the status of the Compositae in published Floras, checklists and available related databases. During that exercise, the available database for the checklist of the Colombian Flora (Bernal *et al.* 2015) was reassessed for its completeness (of both genera and species), for the Compositae. At that time, 230 genera (together with 7 splits from *Vernonia* Schreb.), including 1274 species, this represented 523 endemic species (41.1% of the total of Compositae), and equivalent to 5.6% of the Flora's angiosperm total. However, a significant number of taxa (both genera and species) were missing from the database; this re-assessment was put on hold until mid-2017. Whilst the number of genera remains the same (with some added and others deleted), the current estimates are 1260 species (excluding purely cultivated species), including 582 endemics (46.3%).

In 2016, the Royal Botanic Gardens, Kew, became a partner in a joint programme, Colombia Bio (Colombia BIO 2016), established by ColCiencias

(the Colombian Administrative Department of Science, Technology and Innovation), and aimed at increasing knowledge of Colombia's biodiversity. This partnership involves the Instituto de Investigación de Recursos Biológicos Alexander von Humboldt (FMB = Herbario Federico Medem Bogotá), and the Universidad Pedagógica y Tecnología de Colombia; Kew's involvement is funded in part by the British Council (via BEIS = Department for Business, Energy & Industrial Strategy, UK). As a subset of Colombia BIO, Boyacá Bio is a regional programme which Kew has become directly involved with. The intention is to provide checklists of some of the more important families of flowering plants (Araceae, Leguminosae, Myrtaceae) where Kew has some expertise, and often in collaboration with Colombian researchers, for the Departamento de Boyacá. The experience of one author (DJNH) in creating annotated systematic checklists (see Hind 2011, for an unpublished version of one for Bolivia), was then used to create a similar style of preliminary checklist of the Compositae for the Departamento de Boyacá, Colombia, en route to a full draft list of the Compositae of Colombia,

Accepted for publication 4 March 2020. Published online 24 August 2020

¹ Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Herbario FMB, Carrera 8 # 15-08, Claustro de San Agustín, Villa de Leyva, Colombia. e-mail: jaguilar@humboldt.org.co

² Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK. e-mail: n.hind@kew.org

supplementing the published Catálogo (Bernal *et al.* 2016a, b).

Whilst the second author (DJNH) constructed the Compositae checklist for the Departamento, constant checks were made against the virtual herbarium of the Instituto de Ciencias Naturales de la Universidad Nacional de Colombia (= COL). These were primarily concerned with a visual confirmation that the determinations of the material were correct, as well as supplying supplementary information to be used in the checklist (e.g. altitude of collections, habit and habitat, and flowering time). At present, following the second author's listing, there are 23 native or native and naturalised genera from the tribe Senecioneae present in Colombia. The two largest genera are the shrubby *Monticalia* C.Jeffrey (45 spp. — c.f. no accepted species in Avila *et al.* 2016; 32 spp. endemic) and lianescent *Pentacalia* Cass. (46 spp. — cf. 103 spp. in Bernal *et al.* 2016a, b; 34 spp. endemic); *Senecio* L. (mostly perennial stoloniferous herbs in Colombia) is the third largest with 37 spp. recorded to date (compared to 43 spp. in Avila *et al.* 2016; 24 spp. endemic), of which an estimated 24 spp. are considered endemic to the country. Within Boyacá only 15 genera have been recorded, although the same genera are foremost, with *Monticalia* the largest (19 spp., 13 spp. endemic) and *Pentacalia* with only 9 spp. (6 spp. endemic), except that *Senecio* is the second largest (17 spp.), of which 11 spp. are endemic.

However, it was during that process that it became clear that within *Senecio* some imaged collections did not belong to the taxa to which they had been determined. Most of the senecios in Boyacá are high altitude species (3,000 – 5,000 m, rarely down to 2,000 m) and, excluding the weedy *S. vulgaris* L. (which is only present in its discoid form in Colombia — e.g. Cuatrecasas 13658, Díaz-Piedrahíta *et al.* 958, Knuth 3379, Velasco 184), 75% of them possess radiate capitula, together with one variety of *S. comosus* Sch.Bip.; half of the radiate species have yellow ray limbs, the others purple or carmine-coloured. All of the discoid-headed taxa possess yellow corollas. Material determined as *S. leucanthemoides* Cuatrec., other than the type collection, Cuatrecasas *et al.* 12050 (and the contemporaneous Cuatrecasas *et al.* 12055), was clearly not conspecific. In addition, material determined as *S. adglacialis* Cuatrec. was clearly not that species, but appeared identical to that mis-determined under *S. leucanthemoides* together with one additional sheet under the easily-recognised *S. formosus* Kunth. This unmatched material is of a distinctive rhizomatous perennial herb with entire to sparsely serrate sessile leaves and solitary, scapiform radiate capitula, with yellow ray limbs.

The two authors of the present paper made contact, via Dr Steven Sylvester (now Associate Professor at NF, Nanjing Forestry University), during

November 2017, and the first author examined the errant herbarium material in COL, as well as the material in FMB, securing a loan of material from FMB to COL, and undertaking fieldwork to check the localities highlighted by the material seen in COL virtual herbarium. It became clear that this material is indeed of a species of *Senecio* not previously recognised for Colombia and it is formally described below.

Description

***Senecio scapioides* Aguilar-Cano & D.J.N.Hind, sp. nov.**
Type: Colombia. Departamento de Boyacá: Municipio de Arcabuco, Cordillera Oriental, Vereda Rupavita, Parque Natural Regional Paramo del Valle, 5°44'45"N, 73°22'7"W, 3696 m, 18 Nov. 2017 (fl.), J. Aguilar-Cano 4434 (holotype FMB, isotype K).

<http://www.ipni.org/urn:lsid:ipni.org:names:77206330-1>

Rhizomatous, single-stemmed *perennial herbs* up 45 cm tall; *basal stem* creeping, decumbent, thin-woody, 1.5 – 3.5 mm diam., unbranched. *Inflorescence* scapiform, erect, 15 – 37 cm tall, purple, indumentum densely arachnoid and sericeous, white, covering stems, abaxial leaf surface and involucre, together with glandular, simple multicellular-uniseriate, whitish hyaline trichomes 1.4 – 2.8 mm long, composed of 22 – 36 (– 42) short-oblong cells, these becoming gradually smaller towards apex, ending in a rounded cell. *Basal leaves* 15 – 20 together in a loose rosette, these present at anthesis, leaf blade narrowly oblanceolate-oblong to oblanceolate-oblong, 22 – 45 (– 50) × 3 – 6 mm, sessile, lamina base abruptly attenuate and not dilated at node, herbaceous, pubescent especially along margins, on apex and veins, abaxially whitish green to purple, with an arachnoid and sericeous indumentum, adaxially green, midrib purple, margins entire to obscurely dentate and moderately revolute, apex acute-apiculate; *cauline leaves* 7 – 15, dissimilar to basal leaves, sparser and gradually smaller towards stem apex, sessile, base dilated but not auriculate, blade narrowly oblong-lanceolate or linear-lanceolate, 16 – 28 × 3 – 5 mm, herbaceous, adaxially green, densely sericeous-pubescent, abaxially completely purple and indumentum densely arachnoid and pubescent, midrib purple, margins dentate and strongly revolute, apex acute-apiculate. *Capitula* heterogamous and radiate, solitary and nodding; involucre campanulate, 10 – 33 × 16 – 43 mm, involucre closely subtended by a calyculus, calycular bracts 10 – 12, linear-lanceolate, 13 – 13 × 0.9 – 1.3 mm, herbaceous, green and apically purplish, densely arachnoid and sericeous, apex acuminate with a tuft of short glandular, unicellular trichomes; phyllaries 15, uniseriate, linear-lanceolate, 14 – 17 × 1.3 – 2.5 mm, herbaceous, green, apically purplish, indumentum arachnoid and sericeous,

margins scarious, 0.4 – 0.8 mm wide, often fimbriate, apex acuminate with a tuft of short, glandular, unicellular trichomes; receptacle weakly convex, 5 – 6 × 9 – 10 mm, surface honeycombed. *Ray florets* 16 – 18, female and fertile, ray limb elliptic-oblong, 15.1 – 16.2 × 6.2 – 7.2 mm, yellow, glabrate, apex finely 3-dentate, 8-veined, corolla tube yellow, cylindrical, 5.9 – 6.4 mm long, densely pubescent towards apex with simple, glandular, multicellular uniseriate trichomes, 0.3 – 0.6 mm long; staminodes present; style yellow, 5.4 – 8.5 mm long, base dilated, style arms with marginal stigmatic lines, 1.3 – 2.1 mm long, apices truncate, covered by regularly distributed papillae. *Achenes* ovate, 1.5 – 1.9 × 0.5 – 0.8 mm, covered with short, glandular, unicellular trichomes; carpogonium symmetrical in a broad ring, 0.2 mm diam.; pappus setae numerous, unequal, biseriate, 4.5 – 7.3 mm long, barbellate, deciduous, white. *Disc florets* numerous, 90 – 200, hermaphrodite, fertile; corollas bright yellow at anthesis (brownish post anthesis), 8.2 – 9.9 mm long, limb funnel-shaped, tube 3.0 – 3.6 mm long, glabrate, throat 4.4 – 5.4 × 1.1 – 1.5 mm, 5-lobed, triangular-ovate, 0.9 – 1.0 × 0.5 – 0.6 mm, abaxially thickened and adaxially with a tuft of papillae; anthers barely exerted, 2.8 – 3.0 × 0.2 – 0.4 mm, apical anther appendages lanceolate, anther bases calcarate, anther collars 0.8 – 1.0 mm long, in front view 0.2 – 0.3 mm and 0.1 – 0.2 mm wide (proximal and distal measurements respectively), in side view distally flattened and proximally elongate, balusterform, composed of 8 – 9 layers of rectangular cells; filaments flat, inserted below apex of corolla tube and adnate throughout; style yellow, 7.9 – 10.6 mm long, glabrous, style arms 1.7 – 2.2 mm long, with marginal stigmatic lines, apices sub-truncate to obtuse with a tuft of irregularly distributed papillae, longer papillae surrounding apex and shorter papillae covering whole surface of apex. *Achenes* cylindrical, 1.7 – 2.9 × 0.5 – 0.8 mm, covered with short, glandular, unicellular setulae; carpogonium a symmetrical broad ring, 0.2 – 0.3 mm in diam.; pappus setae numerous, biseriate, unequal, 6.2 – 9.5 mm long, barbellate, deciduous, white.

RECOGNITION. In habit, leaf shape and yellow radiate capitula *Senecio scapioides* resembles *S. adglacialis* but the new species differs in having a densely arachnoid indumentum and in being sericeous pubescent on its stems, abaxial leaf surface and involucre (vs sparsely arachnoid and sericeous pubescent), generally longer glandular trichomes (1.2 – 2.5 mm long vs 0.5 – 1.7 mm long), possessing solitary capitula (vs inflorescences densely sub-corymbiform, 3 – 5-headed, very rarely single-headed on dwarf specimens, e.g. *van der Hammen* 1317), an involucre of fewer phyllaries (15 vs 20) and fewer ray florets (16 – 18 vs 24). (Figs 1, 2 & 3), (Table 1).

DISTRIBUTION AND HABITAT. *Senecio scapioides* occurs in high grass páramos up to the lower limit of the superpáramo, at elevations of 3450 – 4053 m, throughout the north Cordillera Oriental of Colombia, including the páramo complexes of La Rusia, Iguaque-Merchan, Pisba and Tota-Bijagual-Mamapacha in the Department of Boyacá, and the complex of Chingaza in the Department of Cundinamarca, Colombia (Map 1). The species is a solitary terrestrial plant occurring as scattered individuals with low population density on the summits of mountains and on plateaus dominated by grass-stem rosette plant communities of *Calamagrostis effusa* (Kunth) Steud., *Chusquea tessellata* Munro (Poaceae) and *Espeletia barclayana* Cuatrec. (Compositae) (Fig. 3A).

SPECIMENS EXAMINED. COLOMBIA. Boyacá: Municipio de Arcabuco, Cordillera Oriental, Vereda Rupavita, Parque Natural Regional Paramo del Valle, 5°44'45"N, 73°22'7"W, 3696 m, 18 Nov. 2017 (fl.), *J. Aguilar-Cano* 4434 (holotype FMB, isotype K); Municipio de Belen, Complejo Guantiva-La Rusia, parte alta de la montaña, cerca al Boqueron El Consuelo, 6°4'8.3"N, 72°57'7.2"W, 4053 m, 23 Nov. 2017, *M. Diazgranados* 4443 (COL!, FMB!, K!); Municipio de Duitama, Santa Rosa de Viterbo-Duitama, paramo Pan de Azucar, near the road to El Carmen [5°54'1.56"N, 73°3'20.03"W], 3450 m, 30 Nov. 1998, *D. Stancik* 1490 (COL-422996!; FMB-36140!); Municipio de Mongua, Vereda Mongui, alrededores y parte alta del sector Laguna Negra. Parque Natural Regional Siscunsi-Oceta, 5°41'22.8186"N, 72°47'20.8032"W, 3832 m, 29 Nov. 2017, *M. Diazgranados* 4532 (COL!, FMB!, K!); Municipio de Pesca, Vereda Butaga, Páramo de La Cortadera, Mata Blanca, [5°31'21"N, 73°7'48"W], 3750 m, 16 Jan. 1982, *Matilde Bejarano* 92 (COL-242620!); Municipio de Toca, Páramo Cortadero, boundary with Páramo Siachoque. 5°30'N, 73°15'W, 3650 m, 14 Nov. 1998, *D. Stancik* 1380 (COL-421010!; FMB-37827!); Municipio Socotá, Páramo Pisba, Alto de Cardon. [6°3'45"N, 72°36'2"W], 3540 m, 11 Feb. 1999, *D. Stancik* 2317 (COL-430122!; FMB-38144!). **Cundinamarca:** Municipio de Fomeque, Páramo de Chingaza, cordilleras vecinas a la laguna y límite con el departamento del Meta [4°31'15"N, 73°42'55"W], 3200 – 3640 m, 1 Nov. 1966, *G. Huertas & L. Camargo* 6587 (COL-117236!, COL-205612!); Municipio de Fomeque, Parque Nacional de Chingaza, orilla del camino a San Juanito, pasando el boqueron [4°29'11"N, 73°46'33"W], 3610 m, 2 Dec. 1981, *P. Franco* 843 (COL-231859!).

CONSERVATION STATUS. *Senecio scapioides* is endemic to Colombia, occurring in the Andean páramo, throughout the northern Cordillera Oriental in the Departments of Boyacá and Cundinamarca. The estimated extent of occurrence (5989 km²) is almost within the threshold for Endangered in criterion B1 and its area

Fig. 1. *Senecio scapioides*. A habit; B disc floret; C disc floret corolla opened out showing stamen arrangement; D adaxial surface of apex of disc floret corolla lobe; E abaxial surface of apex of disc floret corolla lobe; F anther cylinder opened out; G stamen; H disc floret style; J apex of disc floret style arm showing the truncate apex with a tuft of irregularly distributed papillae; K ray floret; L ray floret style; M calycular bract; N phyllary; P apex of ray floret style arm with apices truncate covered by regularly distributed papillae; Q unicellular trichome from a calycular bract; R unicellular trichome from achene; S multicellular uniseriate trichome from phyllary. All from Aguilar-Cano 4434 (holotype FMB). DRAWN BY GUSTAVO SURLO.

Fig. 2. *Senecio scapioides*. A phyllary base; B phyllary apex; C ray floret; D ray floret limb; E detail of upper part of corolla tube and base of ray limb; F ray floret style; G disc floret corolla; H apex of mature disc floret. All from M. Diazgranados 4443 (K).

Fig. 3. *Senecio scapioides* at the type locality. **A** habitat; **B** habit; **C** basal part of stem and basal leaves; **D** top view of capitulum; **E** lateral view of capitulum. All from *Aguilar-Cano* 4434. PHOTOS: JOSÉ AGUILAR-CANO.

Table 1. Table of comparison of selected characters of *Senecio scapioides* and its closest relatives

Character	Species			
	<i>adglacialis</i>	<i>leucanthemoides</i>	<i>scapioides</i>	<i>funkii</i>
distribution	ARA, BOY	SAN	BOY, CUN	MAG, VEN
elevation (m)	3900 – 4400	3800	3200 – 4053	3345 – 4150
habitat	scree, sandy soil and rarely in grassland	no data	grassland and frailejónal*	among rocks
habit	perennial herbs	subshrubs or suffrutescent	perennial herbs	suffrutescent
total height (m)	0.4 – 0.7	0.6 – 1.0	0.20 – 0.45	0.5 – 1.0
rhizome diam. (mm)	4.0 – 8.3	3.0 – 6.8	1.5 – 3.5	no rhizome
trichome length (mm)	0.5 – 1.7	0.4 – 0.7	1.2 – 2.5	glabrous
basal stem branching	simple	branched	simple	branched
inflorescence stem indumentum	sparsely arachnoid and short-pubescent	sparsely arachnoid and short-pubescent, later glabrescent	densely arachnoid and sericeous	glabrous
basal leaves: petiole presence	absent	present	absent	present
basal leaves: petiole length (mm)	–	40 – 50	–	40 – 55
leaves: lamina size (length × width, mm)	50 – 129 × 5 – 12	25 – 65 × 9 – 16	22 – 45 (– 50) × 3 – 6	10 – 80 × 18 – 25
leaves: lamina shape	45 – 110 × 7 – 14	24 – 65 × 3.5 – 8	16 – 28 × 3 – 5	65 – 165 × 16 – 26
	elliptic-oblanccolate	elliptic	oblanccolate-oblong narrowly	narrowly elliptic
	oblong-lanceolate	narrowly-ovate	oblong-lanceolate or linear-lanceolate	narrowly elliptic
basal leaves: lamina base shape	narrowly-attenuate, dilated	narrowly-attenuate	abruptly attenuate, not dilated	decurrent
cauline	clasping, sub-auriculate	amplexicaul, auriculate	dilated but not auriculate	amplexicaul, auriculate
leaves: lamina indumentum	glabrescent	glabrescent	pubescent	glabrous
cauline (adaxial surface)	sparsely pubescent	sparsely pubescent, arachnoid	densely pubescent	glabrous
basal (abaxial surface)	glabrescent	glabrescent	arachnoid and pubescent	arachnoid
cauline (abaxial surface)	sparsely arachnoid and pubescent	sparsely pubescent, arachnoid	densely arachnoid and pubescent	arachnoid
inflorescence stem height (m)	0.3 – 0.5	0.3 – 0.4	0.15 – 0.37	0.10 – 0.15
inflorescence	scapiform, densely sub-corymbiform	scapiform, laxly corymbiform	scapiform, capitula solitary	scapiform, densely corymbiform, leafy throughout
number of capitula	3 – 5-headed (rarely solitary)	2 – 5-headed	heads solitary	3 – 4-headed
posture of mature capitula	nodding	erect	nodding	erect
phyllary indumentum	pubescent	puberulent	densely pubescent	glabrous
calycular bracts: number	10 – 12	numerous	10 – 12	16
calycular bracts: size (length × width, mm)	10 – 18 × 1.5 – 2.0	10 – 15 × 0.6 – 0.9	13 × 0.9 – 1.3	4.2 – 5.5 × 0.4 – 0.5
calycular bracts: shape	linear-lanceolate	linear-lanceolate	linear-lanceolate	linear
phyllaries: number	20	20	15	15 – 20
phyllaries: size (length × width, mm)	15 × 2 – 3	13 × 2 – 3	14 – 17 × 1.3 – 2.5	7.7 – 9.8 × 0.7 – 1.0

Table 1. (continued)

phyllaries: shape				
disk floret				
	florlets number	linear-lanceolate	oblong-lanceolate	linear-lanceolate
	corolla limb (mm)	170 – 180	numerous	numerous
	corolla tube length (mm)	6 – 6.5	6 – 7	3.4 – 4.1
	florlets number	3.0	3.0	2.1 – 3.1
ray floret	corolla limb shape	24	20 – 40	12 – 15
	corolla limb size (length × width, mm)	broadly elliptic	elliptic-oblong	elliptic-oblong
	corolla limb vein number	12 × 5 – 6	15 – 18 × 4.5	5.1 – 6.6 × 0.7 – 1.1
	corolla tube length (mm)	10	4	3
		3 – 4	5	3.6 – 4.3

**“vegetation dominated by *Espeletia Mutis* ex Bonpl. and other members of the subtribe Espeletiinae”

of occupancy is (348 km²) within the threshold for Endangered in criterion B2. The number and size of the populations is not well-known, and the species could not be accurately assessed based on its population size and decline. Nevertheless, the modern threats to habitat of livestock grazing and frequent fires, as well as climate warming and drying in the northern Andes (Vásquez *et al.* 2015; IPCC 2014), and more than a third of the Páramos de Boyacá now destroyed, mainly due to the introduction of grasslands and crops (Moreno *et al.* 2016), mean that the species habitat is in continuous decline. Therefore, following the criteria established by the IUCN (2013), *Senecio scapioides* is at present assessed as Endangered [EN B2 ab (iii)] mainly based on its distribution range. Research is urgently needed to establish the current population size of this new species, as well as to improve the understanding of plant diversity of Compositae in the Andean páramo.

PHENOLOGY. Flowering and fruiting simultaneously between November and January.

ETYMOLOGY. The specific epithet derives from the Latin *scapus* (scape) and *-oides* (like) and alludes to the scapiform inflorescence of the species.

DISCUSSION. A comparison of the material has shown that *Senecio scapioides* superficially matches that of *S. adglacialis* Cuatrec. and *S. leucanthemoides* Cuatrec., together forming a close morphological and geographical group within the senecionoid complex (subtribe Senecioninae s.str.) (Nordenstam *et al.* 2009) of specialised herbs or suffrutices exclusively adapted to the high páramos of the north Colombian Cordillera Oriental (Map 1). The three species are morphologically characterised by the presence of stems with creeping rhizomes, basal leaves forming a loose rosette, the inflorescence scapiform (Fig. 1A, 3B, C), yellow radiate capitula similar in size and shape (Fig. 3D, E), and staminodes present in the fertile ray florets (Fig. 2E).

Although most of the collections studied of the new species have been previously identified as *Senecio leucanthemoides* and/or *S. adglacialis*, this is understandable, because both species have yellow radiate capitula and grow in areas of páramo (Fig. 3). Both J. Cuatrecasas and S. Díaz-Piedrahita, renowned American synantherologists, saw material of the undescribed species but did not describe it.

Senecio adglacialis does not grow sympatrically with *S. scapioides*, since no material of the new species has been found from the superpáramo vegetation of the Sierra Nevada del Cocuy (Map 1). This Sierra has a high percentage of endemics within the Colombian Cordillera Oriental and several senecios are endemic to the Sierra (e.g. *S. cocuyanus* Cuatrec., *S. supremus* Cuatrec. and *S. pascuiandinus* Cuatrec.) (Cleef 1981; Miranda-Esquivel *et al.* 2002). Otherwise, *S. leucanthemoides* is only known from the type locality

Map 1. Geographical distribution of *Senecio scapioides* and similar species: *S. adglacialis*, *S. leucanthemoides* and *S. funckii*.

in the páramo of Almorzadero (Departamento de Santander) (Map 1), a humid shrubby páramo at 3800 m elevation, and represents the northernmost distribution for this group of rhizomatous, scapiform species with yellow radiate capitula. A table of comparison of selected characters amongst *Senecio scapioides* and these closest relatives is provided in Table 1.

The radiate capitula with the two types of yellow floret noted in the new species also resembles *Senecio funckii*, a species restricted to two disjunct localities in the Venezuelan Sierra Nevada de Merida and in the Colombian Sierra Nevada de Santa Marta (Ávila *et al.* in Bernal *et al.* 2015; Hokche *et al.* 2008) (Map 1). Nevertheless, the indumentum, habit and leaf type of *S. funckii* are distinctly different, being glabrate, suffrutescent with erect branches, no rhizome, an inflorescence leafy throughout, with flat leaves possessing doubly dentate margins, while *S. scapioides* and its morphological and geographical relatives, are perennial herbs with a basal rhizomatous stem, their basal leaves rosetiform, the inflorescences scapiform and the leaf margins always revolute, crenate, and completely covered with an arachnoid and sericeous indumentum (Fig. 1A; 2A, B; 3B, C).

Acknowledgements

This study was conducted under the memorandum of Collaboration between Instituto Humboldt and RBG Kew, “Building a Partnership of Systematic Excellence” supported by the Boyacá Bio programme and Newton fund (via BEIS = Department for Business, Energy & Industrial Strategy, UK). The authors would like to thank Gustavo Surlo, the Margaret Mee Fellowship Programme Artist Scholar 2018, and the 24th such scholar, for providing the black and white line drawings of this new species. Grateful thanks is also given to two anonymous reviewers whose comments have helped improve the original submission.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons

licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Avila, F., Funk, V. A., Diazgranados, M., Díaz-Piedrahíta, S. & Vargas, O. (contribs.). (2016). Asteraceae. In: R. Bernal, S. R. Gradstein & M. Celis (eds), *Catálogo de plantas y líquenes de Colombia*. Vol. I. Capítulos introductorios — Líquenes a Lythraceae, pp. 795 – 908. Instituto de Ciencias Naturales, Facultad de Ciencias, Sede Bogotá, Universidad Nacional de Colombia.
- Bernal, R., Gradstein, S. R. & Celis, M. (2015). *Catálogo de plantas y líquenes de Colombia*. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá. Available from: <http://catalogoplantasdecolombia.unal.edu.co>. (Data base accessed from 15 August 2016).
- _____, ____ & ____ (eds) (2016a). *Catálogo de plantas y líquenes de Colombia*. Vol. I: Capítulos introductorios — Líquenes a Lythraceae. Instituto de Ciencias Naturales, Facultad de Ciencias, Sede Bogotá, Universidad Nacional de Colombia.
- _____, ____ & ____ (eds) (2016b). *Catálogo de plantas y líquenes de Colombia*. Vol. II: Magnoliaceae a Zygophyllaceae — Especies introducidas y cultivadas. Instituto de Ciencias Naturales, Facultad de Ciencias, Sede Bogotá, Universidad Nacional de Colombia.
- Cleef, A. M. (1981). The vegetation of the páramos of the Colombian Cordillera Oriental. *Diss. Bot.* 61: 1 – 320. Cramer, Vaduz.
- Colombia BIO (2016). Colombia BIO. <https://www.colciencias.gov.co/portafolio/colombia-bio>. Site accessed 26 Dec. 2018 [Downloadable PDF *resena-colombibio-2016* explains the programme, in Spanish.]
- Hind, D. J. N. (2011). *An annotated preliminary checklist of the Compositae of Bolivia*. Vers. 2. [See www.kew.org/science/tropamerica/boliviacompositae for the web version and www.kew.org/science/tropamerica/boliviacompositae/checklist.pdf for the PDF file of the checklist].
- Hokche, O., Berry, P. E. & Huber, O. (2008). In: O. Hokche, P. E. Berry & O. Huber (eds), *Nuevo Catálogo de la Flora Vascular de Venezuela. Dicotiledoneas*, pp. 185 – 352. Fundación Instituto Botánico de Venezuela, Caracas.
- IPCC (2014). R. K. Pachauri & L. A. Meyer (eds), Climate Change (2014). *Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva.
- IUCN Standards and Petitions Subcommittee (2013). *Guidelines for using the IUCN Red List Categories and Criteria. Version 10.1*. Prepared by the Standards and Petitions Subcommittee. Available from: <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> (accessed 25 August 2018).
- Miranda-Esquivel, D., Rangel, J. & Roa, L. (2002). Endemismo en páramos colombianos con base en la distribución de espermatófitos y el análisis de parsimonia de endemismo (PAE). In: C. Ange-Jaramillo, C. Castaño-Urbe & C. L. Durán, *Memorias del Congreso Mundial de Páramos* 1: 253 – 266. Ministerio del Ambiente, Bogotá.
- Moreno, L. A., Andrade, G. I. & Ruíz-Contreras, L. F. (eds) (2016). Biodiversity 2016. *Status and Trends of Colombian Continental Biodiversity*. Research Institute of Biological Resources Alexander von Humboldt, Bogotá.
- Nordenstam, B., Pelsner, P. B., Kadereit, J. W. & Watson, L. E. (2009). Senecioneae. In: V. A. Funk, A. Susanna, T. F. Stuessy & R. J. Bayer (eds), *Systematics, Evolution, and Biogeography of Compositae*, pp. 503 – 525. International Association for Plant Taxonomy, Vienna.
- Vásquez, D. L. A., Balslev, H. & Sklenář, P. (2015). Human impact on tropical-alpine plant diversity in the northern Andes. *Biodivers. & Conservation* 24: 2673 – 2683.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.