Four new threatened species of *Rinorea* (Violaceae), treelets from the forests of Cameroon

Gaston Achoundong¹, Xander van der Burgt² & Martin Cheek²

Summary. Four species of *Rinorea* are described as new to science; all four species are endemic to evergreen rain forest in Cameroon. *Rinorea villiersii* Achound. and *R. amietii* Achound. are placed in *Rinorea* [unranked] *Crassiflorae* M.Brandt, while, *R. dewildei* Achound. and *R. faurei* Achound. fall in *Rinorea* [unranked] *Dentatae* Engl. The first species appears to be endemic to the Solé Forest Reserve northeast of Yabassi in Littoral Region. The second and the third species are found mainly in Littoral and South Regions, *R. amietii* extending to South West Region and *R. dewildei* extending to Central Region. The fourth species, *R. faurei* is an endemic of the Santchou Forest Reserve at the foot of Dschang Plateau in West Region. The taxonomic affinities of the four species are discussed, illustrations and a distribution map are provided, and their conservation status is assessed. All four species are threatened with extinction according to the 2012 IUCN categories and criteria.

Key Words. Butterflies, conservation, *Cymothoe*, staminal tube.

Introduction

In the course of revising the species of Violaceae of Africa, mainly in preparation for the account of the Violaceae for the *Flore du Cameroun*, the first author has, with various collaborators, already published 13 new species to science for this group (Achoundong & Onana 1998; Achoundong & Bos 1999; Achoundong & Bos 2001; Achoundong 2003; Achoundong & Cheek 2003; Achoundong & Cheek 2005; Achoundong & Bakker 2006). Other publications have used provisional names which are now used for the new species published here (Achoundong 1996; Achoundong 1997; Amiet & Achoundong 1996; Achoundong 2000; Bakker *et al.* 2006). In this paper, four such provisional species names are formally published to validate these names.

The most recent studies of the phylogeny and classification of African *Rinorea* Aubl. are set out by Wahlert (2010), Wahlert & Ballard (2012) and van Velzen *et al.* (2015). However, the classification of Brandt (1914) has still not been formally replaced. The first two new species described below, *R. villiersii* Achound. and *R. amietii* Achound. fall within *Rinorea* [unranked] *Crassiflorae* M.Brandt, while the second two species, *R. dewildei* Achound. and *R. faurei* Achound. fall in *Rinorea* [unranked] *Dentatae* Engl.

The genus *Rinorea* is pantropical, with 206 species currently accepted by Plants of the World Online, ([continuously updated](#)), accessed April 2021). Africa is the most species-diverse continent for *Rinorea* with 110

– 150 species (van Velzen *et al.* 2015). *Rinorea* species are mainly forest understorey shrubs or small trees. Morphologically, in continental Africa, they are characterised by having alternate, simple leaves, often with petioles of different lengths on the same stem and an usually long, curving apical bud (in the Neotropics and Madagascar, some species e.g. *Rinorea* sect. *Pubiflora* Wahlert & H.E.Ballard have opposite leaves). The flowers are often green, dull yellow, or shades of white and are zygomorphic. There are three sets of petals in *Rinorea*: an anterior petal (also known as the lower or ventral petal), two lateral petals and two posterior petals. These are likely homologous to the three sets of petals in other strongly zygomorphic genera of Violaceae, such as *Viola* L. (Wahlert 2010).

The anterior petal is larger than the other petals, and often modified, with taxonomically important, often diagnostic characters. The androecium has a staminal tube which is also zygomorphic: the anterior (lower or ventral) side is longer and entire, while on the dorsal side of many of the African species, the staminal tube is generally shorter and incised with a V-shaped cleft. The fruits are typical of the family: dry, trilocular, septicidal capsules with parietal placentation.

Rinorea are ecologically important and diverse in African forests, often with several sympatric species in one forest. For example, ten species were recorded in a few square kilometres of the Mefou Proposed National Park near Yaoundé (Achoundong in Cheek *et al.* 2011). Many species are range-restricted, found

¹ IRAD-Cameroon National Herbarium, BP 1601, Yaoundé, Cameroon.

² Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK. e-mail: m.cheek@kew.org

in such small areas that they are at risk of extinction from forest clearance. *Rinorea dewitii* Achound., *R. faustiana* Achound., *R. simoneae* Achound. and *R. thomasi* Achound. are all assessed as threatened in the *Red Data Book of the Flowering Plants of Cameroon* (Onana & Cheek 2011) and all but the first can be found on the IUCN Red List (iucnredlist.org) e.g., *R. thomasi* (Darbyshire & Cheek 2004a; Cheek 2017; Darbyshire & Cheek 2004b). In neighbouring Gabon, the recently published *R. calcicola* Velzen & Wieringa is also range-restricted and of conservation concern (van Velzen & Wieringa 2014). Cameroon has the highest species-diversity for the genus in tropical Africa with 53 species listed (Onana 2011), followed by Gabon, with 46 species (Sosef *et al.* 2006). However, the superficial similarity between species has made identification difficult for taxonomists, e.g., 194 specimens of *Rinorea* are listed as unidentified to species for Gabon in Sosef *et al.* (2006).

African *Rinorea* species are of great interest to entomologists, being important larval food plants of the butterfly genus *Cymothoe* (the gliders). Twenty-eight species of *Cymothoe* are known to feed on *Rinorea*, of which 18 are strictly monophagous, six are oligophagous and three feed on up to six species of *Rinorea* (Amiet & Achoundong 1996).

Materials & Methods

All specimens cited have been seen. Herbarium citations follow *Index Herbariorum* (Thiers, [continuously updated](#)). Specimens were studied online, on loan from or at BR, K, P, WAG and YA. We also searched *JSTOR Global Plants* (<https://plants.jstor.org/> accessed April 2021) for additional materials. Taxonomic authorities follow the *International Plant Names Index*, (IPNI [continuously updated](#)), and nomenclature follows Turland *et al.* (2018). The conservation assessments were made using Bachman *et al.* (2011) following the categories and criteria of IUCN (2012). Herbarium material was examined with a Leica Wild M8 dissecting binocular microscope fitted with an eyepiece graticule. Measurements were made from rehydrated material. The terms and format of the description follow the conventions of Achoundong & Cheek (2005). Post-facto georeferences for specimens without coordinates were obtained from Google Earth.

(https://www.google.com/intl/en_uk/earth/versions/).

Lamina 10 – 12 × 3.5 – 4.5 cm, glandular beneath; ovary glabrous ***Rinorea umbricola***
 Lamina 11 – 22 × 3 – 7 cm, not glandular beneath; ovary pubescent ***Rinorea villiersii***

Taxonomic Treatment

1. *Rinorea villiersii* Achound. sp. nov. Type: Cameroon, Littoral Region, Sole, north-west of Yabassi, fl. 20 May 1992, Achoundong 1928 (holotype YA; isotypes, P, WAG).

<http://www.ipni.org/urn:lsid:ipni.org:names:77220452-1>

Rinorea villiersii Achound. ined. (Achoundong 1996, 1997; Onana 2011: 151).

Rinorea solensis Achound. ined. (Achoundong 1996, 1997).

Evergreen shrub or treelet 1 – 2 (– 3) m tall, densely branched. Stems and leaves pubescent when young, glabrous at maturity. *Leaves* white when young, subcoriaceous, elliptic, 11 – 20 (– 22) × 3 – 5 (– 7) cm strongly long acuminate, base cuneate, margin serrate; lateral nerves 10 – 13 on each side of the midrib. Petiole 1.5 – 2 (– 3) cm long. Stipules triangular, 2 – 3 (– 4) × 0.5 – 1 (– 2) mm, mucronate, with two prominent longitudinal nerves, margins crenate, ciliate. *Inflorescence* terminal, corymbiform, 2.5 – 3.5 (– 5) × 1.5 – 2 (– 3) cm. Bracts 5 – 6, triangular, 3 – 4 (– 5) × 1 – 2 (– 3) mm, mucronate, ciliate, pubescent. *Flowers* zygomorphic, 3 – 4 (– 5) × 2 – 3 (– 4) mm; sepals 5, green, triangular, (1 – 2 (– 3) × 0.5 – 1 (– 1.5) mm ciliate. *Petals* 5, white, ovate, 4 – 5 (– 6) × 1 – 1.5 (– 2) mm; anterior (large) petal broader than the posterior (upper) and the lateral petals, constricted towards the apex, inner surface densely hairy towards the apex. Androecium 3 – 4 mm long; staminal tube 0.5 – 1 (– 1.5) mm long, margin entire, undulate, with a V-shaped incision in the dorsal face; filaments short, inserted on the inner face of staminal tube; staminal tube border hiding the lower part of anthers, anthers 0.5 – 1 (– 1.5) mm long; connective appendage red, ovate, up to 0.5 – 1 (– 1.5) mm long, thecal appendages filiform, bifid. Ovary fusiform, hairy, 1 – 1.5 (– 2) mm long; style erect, 1 – 1.5 (– 2) mm long. *Fruit* white, trilocular, fusiform 1 – 1.5 (– 2) × 1 – 1.5 cm, tuberculate. *Seeds* six per fruit, white, tetrahedral. Fig. 1.

RECOGNITION. *Rinorea villiersii* Achound. is similar to *R. umbricola* Engl., in habit, the long sepals, and the large androecium. It differs in the white juvenile leaves (vs green), the leaves long-acuminate with cuneate bases, pubescent, lacking gland dots on the lower surface (vs acuminate, cordate, glabrous, with gland dots on the lower surface), the ovary fusiform, pubescent (not ovoid, glabrous). The two species can be separated by the following key characters:

Fig. 1. *Rinorea villiersii*. A habit, flowering stem; B flower, side view; C sepal adaxial view; D sepal abaxial view; E anterior (large) petal, adaxial view; F one of the four smaller posterior and lateral petals, adaxial view; G staminal tube, side views. A & B from Achoundong 2212 (WAG); C – G from Achoundong 1928 (WAG). DRAWN BY J. M. (HANS) DE VRIES.

DISTRIBUTION. Cameroon. *Rinorea villiersii* is endemic to the Solé Forest, Solé-Dibong area, north west of Yabassi town, Littoral Region (Map 1).

SPECIMENS STUDIED. CAMEROON. Littoral Region, Solé, Yabassi area, fl., 15 March 1991, Amiet in *Achoundong* 1792 (P, YA); *ibid.*, fl. 20 May 1992, *Achoundong* 1928 (holotype YA; isotypes P, WAG); *ibid.*, fl. fr., May 1994, *Achoundong* 2212 (WAG, YA); Yabassi area, river bank in Solé Forest Reserve, fl., 9 May 2004, *Achoundong* 2328 (YA).

HABITAT. *Rinorea villiersii* is known from lowland evergreen rain forest in the Solé-Dibong area north-west of Yabassi. The relief in its range is undulating, with valleys and plains. All specimens were collected from the borders of a river, at altitudes from 20 to 100 m.

CONSERVATION STATUS. *Rinorea villiersii* is only known from the Solé Forest Reserve. Based on current evidence and the observations of the first author, we estimate the total extent of occurrence of *R. villiersii*, equating to the global area of occupation, as 20 km². This forest currently has good conservation status in that much of it is intact (Google Earth imagery viewed 11 April 2021) but there are no concrete conservation actions in place. The area has a low human population density. Unfortunately, it is only 14 km along a major road from the settlement of Loum which is a highly populated area with active farmers. They are expected in the future to invade the reserve for farming purposes, resulting in major threats to this species.

Botanical surveys and other plant studies for conservation management in forest areas north, west and east of Solé Forest Reserve have resulted in many thousands of specimens being collected and identified, but failed to find any additional specimens of *R. villiersii* (Cable & Cheek 1998; Cheek *et al.* 2000; Maisels *et al.* 2000; Chapman & Chapman 2001; Harvey *et al.* 2004; Cheek *et al.* 2004; Cheek *et al.* 2010; Harvey *et al.* 2010; Cheek *et al.* 2011). Although there are still poorly sampled locations with intact natural habitat in Cameroon, it is possible that *R. villiersii* is truly localised, range-restricted and threatened as are several other species of the genus (see introduction). *Rinorea villiersii* is therefore here assessed, on the basis of the single location, small known range-size and threats described above, as Critically Endangered, CR B1+2ab(iii).

ETYMOLOGY. The specific epithet commemorates the late Dr Jean François Villiers (1943 – 2001), taxonomic specialist in mimosoid Leguminosae (also known as Mimosaceae) of Africa and Madagascar, staff member of the Herbarium at MNHN, Paris and lecturer at the University of Yaoundé. He was the supervisor of the first author's doctoral thesis on

Rinorea. He did much to encourage botanists in Cameroon to study conservation.

VERNACULAR NAMES & USES. None are recorded.

2. *Rinorea amietii* Achound. sp. nov. Type: Cameroon, Littoral Region, Marienberg, March 1992, fl. *Achoundong* 1853 (holotype P; isotypes WAG, YA).

<http://www.ipni.org/urn:lsid:ipni.org:names:77220453-1>

Rinorea amietii ined. *Achoundong* (1996: 544); Amiet & *Achoundong* (1996: 465); *Achoundong* (1997); Onana (2011: 150).

Shrub or small tree 5 – 7 (– 8) m high. Diameter of the trunk at 1.3 m above the ground up to 10 cm. Stems and leaves ferruginously hairy, most densely pubescent when young. Leaves subcoriaceous, elliptic, 9 – 17 (– 20) × 3 – 7 (– 9) cm, broadly acuminate, base cuneate, margin serrate, lateral nerves 12 – 13 on each side of the midrib. Petiole 5 – 3 (– 4) cm long. Stipules triangular, 2 – 5 × 1 – 1.5 mm, mucronate, densely pubescent. *Inflorescences* terminal or subterminal, pubescent, corymbiform, 1.5 – 5 (– 6) × 1.5 – 2.5 (– 4) cm, bracts triangular 3 – 5 (– 6) × 1 – 1.5 (– 2) mm, pubescent, densely so on the midrib, apiculate. *Flower* zygomorphic, 4 – 5 (– 6) × 3 – 4 mm. Sepals 5, bright yellow, oblong 1.5 – 2.5 (– 3) × 0.5 – 1.5 (– 2) mm, pubescent. Petals 5, yellow, ovate to elliptic, 2 – 4 (– 5) × 1 – 1.5 (– 2) mm, narrowed at the base, apex rounded, ciliate. Anterior (lower) petal enlarged at the middle, narrowed at the extremities, more strongly near the apex; hairy at the apex of the inner face, lateral and posterior petals oblong, smaller. *Androecium* 2.5 – 3.5 (– 4) mm long; staminal tube up to 1 – 1.5 (– 2) mm high; margin free, undulate, with a V-shaped incision in dorsal face; filaments all inserted on the inner face of the staminal tube; anthers 0.5 – 1.5 (– 2) mm long; connective appendage red, ovate 1 – 1.5 (– 2) mm long, thecal appendages bifid, filiform. Ovary bottle-shaped, 0.5 – 1 (– 1.5) mm long, glabrous; style erect, 1.2 – 2 (– 2.5) mm long, enlarged at the summit, with longitudinal furrows. *Fruit* ovoid, 2 × 1.5 cm, surface white, slightly tuberculate. *Seeds* six, black, angular, dorsal faces convex, inner faces planar. Fig. 2.

RECOGNITION. In its habit, its ferruginous leaves and its very short apical stem bud, *Rinorea amietii* is morphologically (but not phylogenetically) close to *R. subsessilis* M.Brandt, but differs in the leaves being pubescent and non-glandular below (vs glabrous and glandular), the petal midrib pubescent (not glabrous), and in additional features in the key below. The two species are separated by the following characters:

Fig. 2. *Rinorea amietii*. **A** habit, sterile leafy shoot; **B** habit showing position of terminal inflorescence; **C** detail of **B** showing portion of inflorescence; **D** habit, stem with inflorescence and buds; **E** – **G** views of staminal tube. **A** – **C** & **E** – **G** from Achoundong 1744 (YA); **D** from Achoundong 1853 (WAG). **A** – **C** & **E** – **G** DRAWN BY JEAN MICHEL ONANA, **D** BY J. M. (HANS) DE VRIES.

Lamina glabrous, glandular below, fruit green, smooth, inflorescence glabrous, 3 – 15 cm long, sepals glabrous, ovate, apex rounded not mucronate, petal midrib glabrous **R. subsessilis**
 Lamina pubescent, not glandular below, fruit white, verrucose, inflorescence pubescent, 2 – 5 cm long, sepals densely pubescent, triangular to trapezoid, acute at the summit, mucronate, petal midrib pubescent. **R. amietii**

DISTRIBUTION. Cameroon. *Rinorea amietii* is endemic to coastal (littoral) Cameroon in the South West Region, Littoral Region and South Region (Map 1).

SPECIMENS STUDIED. CAMEROON. South West Region, Ekumbe Mufako, fl., May 1994, *Ndam* 1198 (K, SCA, YA); Ekumbe Mufako, fl., May 1994, *Sonké* 1217 (K, SCA, YA); Ekumbe Mufako, fl., May 1994, *Thomas* 10121 (K, SCA, YA); **Littoral Region**, Dizangue area, Ossa Lake, fl. 5 April 1984, *Achoundong* 944 (P, YA); km 30, Edéa-Douala road, fl. 13 Jan. 1990, *Achoundong* 1550 (P, YA); Mangonbe border, fl. 2 March 1990, *Achoundong* 1627 (K, P, YA); 30 km on Edéa-Douala road, fl., fr., 28 July 1990, *Achoundong* 1744 (WAG, YA); Bonépoupa, 30 km on Edéa-Douala road, fl. 22 May 1991, *Achoundong* 1824 (P, YA); km 30 Edéa-Bonépoupa road, fl. 22 May 1991, fl., fr. *Achoundong* 1836 (P, YA); Mangonbé border, Edéa-Songloulou road, 22 May 1991, fl. *Achoundong* 1839 (YA); **Littoral Region**, Marienberg, March 1992, fl. *Achoundong* 1853 (holotype P; isotype, K); region de Mouanko, fl. March 1992, *Achoundong* 1864 (YA); Edéa-Douala road 19 Feb. 2001, *Achoundong* 2106 (YA); 20 km from Douala on Edéa road, Carrière Ducam Duclair, 20 Dec. 2002, *Achoundong* 2198 (YA); 25 km NW of Edéa, SW of “Carrière Ducam Duclair”, forest at field of Marcel Melingui Engama 3°55.01'N, 10°03.30'E, fr. 5 May 2006, *van Velzen* 50 (WAG041636); **South Region** Kribi area, Bidou I, fl. March 1993, *Achoundong* 2011 (YA); Bipindi area, Nsola, fl., March 1993, *Achoundong* 2020 *bis* (YA); Bipindi area, Nsola, fl., June 1994, *Achoundong* 2204 (YA); 20 km from Kribi on Lolodorf road, fr. 7 Aug. 1969, *Bos* 5154 (K, WAG); 20 km from Kribi, 3 km N of Lolodorf road, fr. 18 July 1969, *Bos* 5101 (K, WAG); Bipindi, fl., 1896, *Zenker* 1078 (K); Bipindi, fl. 1896, *Zenker* 1079 (K, P); Bipindi, fl. 1904, *Zenker* 2680 (BM, K).

HABITAT. *Rinorea amietii* is known from littoral lowland evergreen rain forests. The altitude ranges from 0 to 300 m.

CONSERVATION STATUS. The littoral lowland evergreen rain forests of Cameroon are rich in *Rinorea* species. Many of the species are narrow endemics and so are vulnerable to extinction from forest clearance. Littoral forest habitat is highly threatened due to its ability to host industrial agriculture and because it is a target for logging activity. Many industrial farms are already located there, cultivating crops such as banana (*Musa* spp.), palm oil (*Elaeis guineensis*), and rubber (*Hevea brasiliensis*). Due to its proximity to the coastal ports and the density of roads, littoral forest habitat is in demand for clearance for the expansion of these industries.

Currently eight locations are recorded for *Rinorea amietii* based on the 23 specimens studied,

following extensive surveys in Cameroon detailed above under *R. villiersii*. The extent of occurrence is estimated as 9,545 km², and the area of occupation as 60 km². Given the threats above, we assess *R. amietii* as Vulnerable VU B1+2ab(iii).

ETYMOLOGY. The specific epithet commemorates Professor Jean-Louis Amiet, lecturer at University of Yaoundé. He studied the relationship between butterflies and plants, especially that between *Cymothoe* (the gliders with about 75 species in Africa) and the genus *Rinorea* (Amiet 1997; Amiet 2000), and inspired the first author in his life's taxonomic research on the genus *Rinorea* in Africa (Amiet & Achoundong 1996). He is commemorated by several other species such as *Afrothismia amietii* Cheek (Thismiaceae, Cheek 2007). Amiet, an entomologist, came to understand in the 1980s that the *Cymothoe* butterflies he studied were often monophagous, the larvae of some species eating only one species of *Rinorea* (see Introduction). However, at that point, these different species of *Rinorea* were not recognised by taxonomists because of the superficial similarity of many other species. To address this difficulty he recruited the first author!

NOTES. Recent data from the chloroplast marker trnL-F and the nuclear ribosomal ITS, indicate that *Rinorea amietii* is related to *R. longisepala* Engl., *R. ledermannii* Engl. and *R. umbricola* Engl. (Robin van Velzen pers. comm. to Achoundong, June 2021) supporting placement in *Rinorea* [unranked] *Crassiflorae*.

3. *Rinorea dewildei* Achound., sp. nov. Type: Cameroon, Central Region, 35 km NW d' Eséka, fl. 23 March 1964, *W. J. J. O. de Wilde* 2218 (holotype WAG; isotypes K001381925, P, YA).

<http://www.ipni.org/urn:lsid:ipni.org:names:77220455-1>

Rinorea dewildei ined. Achoundong (1996: 544), Amiet & Achoundong (1996: 310); Onana (2011: 150).

Shrub or treelet up to 5 m high. Young twigs puberulent. Leaves papery, lamina elliptic, 7 – 18 × 2 – 7 cm, apex broadly acuminate, base cuneate to obtuse, margin serrate, lateral nerves 6 – 8 pairs, 3 – 4 of which are more prominent than the others. Stipules triangular 6 × 2 mm, mucronate, abaxial midrib pronounced, pubescent, margin ciliolate. Petiole 5 – 15 mm long, pubescent. *Inflorescence* terminal, glabrous, corymbose, 3.5 – 4 cm. long; bracts triangular, up to 1 × 1.5 mm,

ciliolate. *Flowers* yellow, 4×2 mm; sepals yellow, triangular, 2.5×1.2 mm, ciliolate. Petals yellow, anterior (lower petal) ovate, 4×1 mm, ciliolate; 3-nerved; posterior and lateral petals oblong, 3×1 mm, apex truncate-retuse, 1-nerved. Androecium 2 – 3 mm long; staminal tube urceolate, 1.5 mm long; margin free, undulate, with a dorsal slit; staminal filaments inserted on the inner face of the staminal tube; filament of the dorsal stamen inserted near the distal edge of the staminal tube, on the dorsal slit; anthers 0.5 mm long; connective appendage red, ovate 0.5 – 1 mm long; thecal appendage bifid. Ovary ovoid, glabrous, 1 mm long; style

2 mm long, expanded at the summit. *Fruit* white, ovoid, tuberculate, $1.5 (-2) \times 1 (-1.5)$ cm. *Seeds* tetrahedral, 0.5×0.5 cm. Fig. 3.

RECOGNITION. In *Rinorea dewildei* the lamina is distinctive in that 3 – 4 of the lateral nerves are more prominent than the others. *Rinorea cerasifolia* M.Brandt shows the same character. *Rinorea dewildei* has petioles 0.5 – 1.5 cm long, sepals twice as long as broad (vs to 5 cm long, sepals as long as broad in *R. cerasifolia*). The species are further distinguished as follows:

Petiole up to 5 cm long, tertiary venation of lamina sparse and inconspicuous; sepals ovate, as long as broad; all staminal filaments inserted on the inner face of the staminal tube.	Rinorea cerasifolia
Petiole 0.5 – 1.5 cm long; tertiary venation of lamina dense and prominent; sepals triangular, twice as long as broad; one of the staminal filaments inserted on the slit of staminal tube margin, the others on the inner face	Rinorea dewildei

DISTRIBUTION. Cameroon. *Rinorea dewildei* is found in the Central Region, Littoral Region and South Region (Map 1).

SPECIMENS STUDIED. CAMEROON. Littoral Region, border of Nyong river at Dehane, fl., 1 March 1990, *Achoundong* 1623 (K, P, YA); *ibid.*, *Achoundong* 2105 (YA); “Carrière Ducam Duclair”, 20 km N of Edéa-Douala road, fl., 20 Feb. 2002, *Achoundong* 2197 (YA); 12 km Edéa road, fl., 23 March 1994, *Achoundong* 2339 ED (P, YA); Edéa area, fl., May 1994 *Achoundong* 17 ED (YA); **Central Region,** Makak, littoral forest, 100 km W of Yaoundé, young fr. Oct. 1938, *Jacques-Felix* 2288, (P); 35 km NW of Eséka, 23 March 1964, *W. J. J. O. de Wilde* 2218 (holotype WAG; isotypes K, P, YA). 6 km NE of Edéa, forest at Eding Island, $3^{\circ}48.97'N$, $10^{\circ}08.17'E$, imm. fr., 7 May 2006, *van Velzen* 43 (WAG0416447). **South Region.** Bibabimvoto, Dipikar Island, in the Campo area along transect T2; fr., 19 July 2000, *Tchouto* T2X195 (WAG); Mabiogo, Dipikar island, in the Campo area along transect T1, fr., 20 Sept. 2000, *Tchouto* T1X144 (WAG); Bimbabivoto, Forest along transect T2 in Campo Maan area, fr., 18 July 2000, *Tchouto* 2951 (WAG).

HABITAT. *Rinorea dewildei* occurs in littoral lowland evergreen rainforest. The altitude ranges from 0 to 700 m. This species appears common in many areas of Cameroonian littoral forest but despite this has surprisingly few collections.

CONSERVATION STATUS. According to Tchouto, *Rinorea dewildei* is common in the Campo-Ma'an National Park. Consequently, the species is well protected within the park. However, at locations outside the park, such as at Makak, Edéa and Eséka, the habitat of this species is being cleared for agriculture and urbanisation.

Currently, after extensive surveys in Cameroon (see under *Rinorea villiersii*), six locations are recorded for

R. dewildei (see specimens above), the extent of occurrence is estimated as 12,430 km², and the area of occupation as 36 km². Given the threats above, we assess *R. dewildei* as VU B1+2ab(iii).

ETYMOLOGY. The specific epithet commemorates Dr W. J. J. O. de Wilde. He and his wife, Brigitte de Wilde-Duyfjes collected the type specimen. Together, from their base at N'Kolbisson in Yaoundé in the 1960s, they collected numerous excellent botanical herbarium specimens in large sets through many parts of Cameroon, including the littoral areas where *Rinorea dewildei* still grows. At this time their home institution, Herbarium Vadenense (WAG), then at Wageningen, Netherlands, was highly active in conducting botanical surveys in poorly studied but species-rich areas of tropical Africa, such as Cameroon and Gabon. Sixty years later, we are still naming species new to science resulting from their efforts, such as this.

NOTES. One anomalous specimen from Gabon (Moyen-Ogooué, Camp Mboumi base, river bank, fr., fl. 20 Aug. 1999, *Issembe* 191 (K, LBV, WAG), shares many similarities with *Rinorea dewildei*, and was initially identified as this species, but is here excluded. It may represent a further undescribed new species to science.

4. *Rinorea faurei* Achound. sp. nov. Type: Cameroon, West Region, Santchou Forest Reserve, fl. *Achoundong* 1936 (holotype YA, isotypes P, WAG).

<http://www.ipni.org/urn:lsid:ipni.org:names:77220456-1>

Rinorea faurei ined. *Achoundong* (1996: 544), *Achoundong* (1997: 255); Onana (2011: 150).

Shrub or treelet 1.5 – 2.5 (– 5) m high, stems and leaves glabrous. *Leaves* sub-coriaceous, elliptic to slightly

Fig. 3. *Rinorea dewildei*. **A** habit, flowering stem; **B** sepal; **C** petal; **D** pistil; **E** & **F** staminal tube, side views. All from *W. J. J. O. de Wilde & B. E. E. de Wilde-Duyfjes* 2218 (WAG). DRAWN BY J. M. (HANS) DE VRIES.

ovate, 12 – 21 × 4.5 – 6 (– 9.5) cm, acuminate, base cuneate to obtuse, margin slightly serrate to subentire; lateral nerves 8 – 12 on each side of the midrib. Petiole 1 – 2.5 (– 5) cm long. Stipules triangular, 6 × 2 mm, mucronate, midrib abaxially pubescent, margin ciliolate. *Inflorescences* terminal or subterminal, thyriform, glabrous c. 4 × 1.5 cm; bracts triangular, 1 × 1.5 mm, apiculate, abaxial midrib and margin ciliolate. *Flowers* white, zygomorphic, 4.5 – 6 mm long. Sepals 5, green, triangular to elliptic 1 – 1.3 × 1 – 1.5 mm. Petals white to yellow, elliptic, flat, 1.5 – 3 × 1 – 1.5 mm, apex rounded, margin ciliate. Anterior (lower) petal broader than the posterior (upper) and lateral petals, slightly constricted towards the apex, inner surface hairy near the apex. Androecium 3 mm long, staminal tube 0.3 – 1 mm long, margin free, with a V-shaped dorsal incision; filaments inserted on the inner face of the tube; anthers of the ventral side (face), fully exerted from the tube with free space

between the tube summit and the lower part of the anthers; anthers 5, each 1 mm long, connective appendage red, ovate, 1 – 1.3 mm long, apex rounded, base decurrent on the upper part of the anthers. Ovary 3-locular, yellow, pyriform, 7 × 1 mm, style straight, cylindrical, white, 2 mm long, slightly constricted at the apex, stigma capitate. *Fruit* white, broader than long, 2 × 3 cm, strongly 3-lobed. *Seeds* six, white, acutely angled, dorsal face convex, ventral face flat c. 5 × 5 mm. Fig. 4.

RECOGNITION. In its habit and its thyriform inflorescence *Rinorea faurei* is close to two other *Rinorea* species of the forest understory: *R. sinuata* Chipp and *R. thomasi*, differing from the first in having the flowers dispersed in an elongated, lax cyme (vs a dome-shaped, compact cyme), differing from the second in the leaves elliptic, ovate or obovate and long-acuminate (vs oblong and shortly acuminate):

1. Lamina oblong, shortly acuminate	<i>Rinorea thomasi</i>
1. Lamina elliptic or ovate to obovate, long-acuminate	2
2. Inflorescence a dome-shaped, compact cyme	<i>Rinorea sinuata</i>
2. Inflorescence an elongated, lax cyme, (flowers dispersed)	<i>Rinorea faurei</i>

DISTRIBUTION. Cameroon. *Rinorea faurei* is endemic to the Santchou area in West Region (Map 1).

SPECIMENS STUDIED. CAMEROON. West Region. Santchou Forest Reserve, fl., 17 March 1991, *Achoundong* 1786 (P, WAG, YA); *ibid.*, fl., fr. 19 May 1992, *Achoundong* 1925 (YA); *ibid.*, *Achoundong* 1936 (holotype YA; isotypes P, WAG).

HABITAT. *Rinorea faurei* occurs in understory of evergreen rain forest in the Santchou Plain at the foot of the Dschang Plateau. The estimated elevation above sea level is 770 m. The area of forest is more than 7000 ha. Santchou Forest Reserve is also the home of *R. dentata* Kuntze, *R. oblongifolia* C.Marquand, *R. yaundensis* Engl. and *R. batesii* Chipp (*Achoundong pers. obs.* 1991 – 1992).

CONSERVATION STATUS. Santchou Forest Reserve has official protection status. However, lack of concrete action on the ground means a strong risk of the reserve being invaded by farmers since the Santchou Plain is intensively cultivated.

Currently, after extensive surveys in Cameroon (see under *Rinorea villiersii* above), only a single location in the sense of IUCN (2012) is recorded for *R. faurei* (see specimens above), the extent of occurrence and the area of occupation is estimated as 70 km². Given the threats above, we assess *R. faurei* as Critically Endangered, CR B1+2ab(iii).

ETYMOLOGY. The specific epithet honours Jean-Jacques Faure, former Head of the Forestry High School in Cameroon. His contribution to the protec-

tion of the environment of Central Africa has been important.

Discussion

More collections of *Rinorea* in Cameroon are needed. The more widespread species such as *R. dewildei* (see note there) have surprisingly few collections, even though the first author has noted they can be quite common within their range. It appears that many collectors avoid targeting the genus in their botanical inventory work, perhaps doubting that one species differs from another and seeking to avoid duplicating collections of the same taxon. The first author, in contrast, targeted the genus in his field surveys, accounting for his collections being the majority of those cited in this paper. This demonstrates how under-collected these Cameroon littoral forests are generally and confirms that more herbarium specimens should be collected to determine which species are present in these forests before they are lost.

About 2000 species of vascular plant have been described as new to science each year for the last decade or more. Until species are known to science, they cannot be assessed for their conservation status and the possibility of protecting them is reduced (Cheek *et al.* 2020). To maximise the survival prospects of range-restricted species there is an urgent need not only to document them formally in compliance with the requirements of

Fig. 4. *Rinorea faurei*. A habit, flowering stem; B flower, side view; C sepal; D petals, large (left) and small (right); E & F staminal tube and pistil, side views; G staminal tube and pistil, opened to show inner surface. All from Achoundong 1936 (WAG). DRAWN BY J. M. (HANS) DE VRIES.

Map 1. Global distributions of *Rinorea villiersii*, *R. amietii*, *R. dewildei* and *R. faurei*. Prepared with www.simplemappr.net

the relevant nomenclatural code (Turland *et al.* 2018), but also to formally assess the species for their extinction risk, applying the criteria of a recognised system, of which the IUCN Red List of Threatened Species is the most widely accepted (Bachman *et al.* 2019). Despite rapid increases over recent years in numbers of plant species represented by assessments on the Red List, the vast majority of plant species still lack such assessments (Nic Lughadha *et al.* 2020).

Increasing representation of plants on the Red List is imperative but documenting threatened species in the level of detail required for inclusion on the Red List is resource intensive (Juffe-Bignoli *et al.* 2016) and securing the required peer-review for completed assessments to be accepted for the Red List can be slow and problematic (Bachman *et al.* 2019). Alternative approaches to evaluating extinction risk, including automated or semi-automated methods involving digitally available information and artificial intelligence, have been applied to plants in recent years (e.g., Darrah *et al.* 2017; Nic Lughadha *et al.* 2019; Zizka *et al.* 2020), often with demonstrable success in correctly predicting the threat status of species previously assessed for the Red List. However, caution is needed in considering such automated assessment approaches for adoption as conservation

tools, not only for a variety of technical reasons (Walker *et al.* 2020) but also because they lack the power that an IUCN Red List assessment has to help change the fate of a species.

As a global standard, the IUCN Red List not only provides stable and evidence-based methods for biodiversity scientists to assess and document the extinction risk of individual species, thus facilitating comparison of the extinction risk of taxonomic groups and areas worldwide (e.g., Nic Lughadha *et al.* 2020; Mair *et al.* 2021), it also supports the safeguarding and sustainability frameworks used by businesses and their major lenders (Bennun *et al.* 2018; Juffe-Bignoli *et al.* 2016). For example, clients of the International Finance Corporation (World Bank Group) are required to use the Red List to inform project risks and to refrain from activities leading to a net reduction in populations of species assessed on the Red List as Endangered (EN) or Critically Endangered (CR), over a reasonable time-scale. Species whose likely extinction has been avoided or reduced as a result of these rules include *Stylochaeton pilosum* Bogner, *Marsdenia exellii* C.Norman, *Raphionacme caerulea* E.A.Bruce (all EN), and *Tarenna hutchinsonii* Bremek. (CR) (Couch *et al.* 2014; Couch *et al.* 2019).

National governments and leaders also recognise the importance of species assessed as threatened by the Red List, as demonstrated recently in Cameroon when, in part due to the high number of plant species on the Red List (Lovell 2020), a logging concession was revoked for the Ebo forest (Kew Science News 2020).

Thus while the Conservation Status statements included in the species treatments in the present paper suffice to highlight the threatened status of these species to the taxonomic botany community, completion and review of full assessments must follow so that they can be published on the IUCN Red List. Their assessments on the Red List will facilitate their inclusion in larger scale studies, including future global extinction risk estimates and conservation prioritisation exercises, but, most importantly, it will enable not only scientists, but also local communities, NGOs and national authorities to promote and take action to safeguard them.

Documented extinctions of plant species are increasing (Humphreys *et al.* 2019) and recent estimates suggest that as many as two fifths of the world's plant species are now threatened with extinction (Nic Lughadha *et al.* 2020). In Cameroon, *Oxygyne triandra* Schltr. and *Afrothismia pachyantha* Schltr. of South West Region, Cameroon are now known to be globally extinct (Cheek & Williams 1999; Cheek *et al.* 2018a; Cheek *et al.* 2019). In some cases, Cameroon species appear to have become extinct even before they are known to science, such as *Vepris bali* Cheek (Cheek *et al.* 2018b).

Most of the 815 Cameroonian species in the Red Data Book for the plants of Cameroon are threatened with extinction due to habitat clearance or degradation, especially of forest for small-holder and plantation agriculture following logging (Onana & Cheek 2011). Efforts are now being made to delimit the highest priority areas in Cameroon for plant conservation as Tropical Important Plant Areas (TIPAs) using the revised IPA criteria set out in Darbyshire *et al.* (2017). This is intended to help avoid the global extinction of additional endemic species such as the *Rinorea* species published in this paper which will be included in the future proposed TIPAs.

Acknowledgements

The first author thanks IRAD-National Herbarium of Cameroon (YA) for support in his retirement that has enabled him to continue and finalise his taxonomic research on Violaceae of Cameroon for the *Flore du Cameroun*. He also thanks colleagues at Wageningen for retrieving and transmitting the excellent figures that illustrate this paper, mainly drawn by J. M. (Hans) de Vries, with valuable contributions by Jean Michel Onana. He thanks Robin van Velzen for data on the

molecular phylogenetic placement of taxa. This paper was completed as part of the Cameroon TIPAs (Tropical Important Plant Areas) project at RBG, Kew, which is supported by Players of Peoples Postcode Lottery. Formal Red Listing of these four species, once they are formally published, will be supported by the John S. Cohen Foundation.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Achoundong, G. (1996). Les *Rinorea* comme indicateurs des grands types forestiers du Cameroun. In: L. J. G. van der Maesen, X. M. van der Burgt & J. M. Van Medenbach-de Rooy (eds), *The Biodiversity of African Plants*, pp. 536 – 544. Kluwer Academic Publishers, Dordrecht. https://doi.org/10.1007/978-94-009-0285-5_69
- _____. (1997). *Rinorea* du Cameroun, *Systématique, Biologie, Ecologie, Phytogéographie*. Thèse Université de Yaoundé I, Yaoundé.
- _____. (2000). Les *Rinorea* et l'étude des refuges forestiers en Afrique. In: M. Servant & S. Servant-Vildary (eds), *Dynamique à long terme des écosystèmes forestiers intertropicaux*, pp. 19 – 29. Mémoire UNESCO, Paris.
- _____. (2003). Novitates Gabonenses 45. Une nouvelle espèce de *Rinorea* (Violaceae) du Gabon. *Adansonia* n.s. 25: 211 – 215.
- _____. & Bakker, F. T. (2006). Deux nouvelles espèces de *Rinorea*, série *Illicifoliae* (Violaceae) du Cameroun. *Adansonia*, sér. 3, 28 (1): 129 – 136.
- _____. & Bos, J. J. (1999). Novitates Gabonenses: 37. Espèces nouvelles de *Rinorea* (Violaceae) du Gabon. *Adansonia* 21 (1): 125 – 131.
- _____. & _____. (2001). Deux espèces nouvelles de *Rinorea* (Violaceae) du Congo et du Gabon. *Adansonia* 23 (1): 155 – 159.

- ____ & Cheek, M. (2003). Two new species of *Rinorea* (Violaceae) from western Cameroon. *Kew Bull.* 58: 957 – 964. <https://doi.org/10.2307/4111209>
- ____ & ____ (2005). Two further new species of *Rinorea* (Violaceae) from Cameroon. *Kew Bull.* 60 (4): 581 – 586.
- ____ & Onana, J. (1998). *Allexis zygomorpha* (Violaceae): a new species from the Littoral Forest of Cameroon. *Kew Bull.* 53 (4): 1009 – 1010. <https://doi.org/10.2307/4118897>
- Amiet, J.-L. (1997). Spécialisation trophique et premières états chez les *Cymothoe*: implications taxonomiques (Lepidoptera, Nymphalidae). *Bull. Soc. Entomol. France* 102: 15 – 29.
- ____ (2000). Les premiers états des *Cymothoe*: Morphologie et intérêt phylogénique (Lepidoptera, Nymphalidae). *Bull. Soc. Entomol. France* 106: 349 – 390.
- ____ & Achoundong, G. (1996). Un exemple de spéciation trophique chez les Lepidoptères: les *Cymothoe* camerounaises inféodées au *Rinorea* (Violacées) (Lepidoptera, Nymphalidae). *Bull. Soc. Entomol. France* 105 (5): 449 – 466.
- Bachman, S. P., Field, R., Reader, T., Raimondo, D., Donaldson, J., Schatz, G. E. & Lughadha, E. N. (2019). Progress, challenges and opportunities for Red Listing. *Biol. Conservation* 234: 45 – 55. <https://doi.org/10.1016/j.biocon.2019.03.002>
- ____, Moat, J., Hill, A., de la Torre, J. & Scott, B. (2011). Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys*. 150: 117 – 126. <https://doi.org/10.3897/zookeys.150.2109>
- Bakker, F. T., van Gemerden, B. S. & Achoundong, G. (2006). Molecular systematics of African *Rinorea* Aublet. (Violaceae). In: S. A. Ghazanfar & H. J. Beentje (eds), *Taxonomy and ecology of African plants, their conservation and sustainable use*, pp. 33 – 44. Royal Botanic Gardens, Kew.
- Bennun, L., Regan, E. C., Bird, J., van Bochove, J. W., Katariya, V., Livingstone, S., Mitchell, R., Savy, C., Starkey, M., Temple, H. & Pilgrim, J. D. (2018). The value of the IUCN Red List for business decision-making. *Conservation Lett.* 11 (1): p.e12353. <https://doi.org/10.1111/conl.12353>
- Brandt, M. (1914). Übersicht über die afrikanischen Arten der Gattung *Rinorea* Aubl. *Bot. Jahrb. Syst.* 51: 405 – 418.
- Cable, S. & Cheek, M. (1998). *The Plants of Mt Cameroon, a Conservation Checklist*. Royal Botanic Gardens, Kew.
- Chapman, J. & Chapman, H. (2001). *The Forests of Taraba and Adamawa States, Nigeria an Ecological Account and Plant Species Checklist*. University of Canterbury, Christchurch.
- Cheek, M. (2007). *Afrothismia amietii* (Burmanniaceae), a new species from Cameroon. *Kew Bull.* 61: 605 – 607. <http://www.jstor.org/stable/20443306>
- ____ (2017). *Rinorea simoneae*. *The IUCN Red List of Threatened Species* 2017: e.T110087540A110087542. <https://doi.org/10.2305/IUCN.UK.2017-3.RLTS.T110087540A110087542.en>
- ____, Etuge, M. & Williams, S. (2019). *Afrothismia kupensis* sp. nov. (Thismiaceae), Critically Endangered, with observations on its pollination and notes on the endemics of Mt Kupe, Cameroon. *Blumea* 64 (1): 158 – 164. <https://doi.org/10.3767/blumea.2019.64.02.06>
- ____, Gosline, G. & Onana, J.-M. (2018b). *Vepris bali* (Rutaceae), a new critically endangered (possibly extinct) cloud forest tree species from Bali Ngemba, Cameroon. *Willdenowia* 48: 285 – 292. doi: <https://doi.org/10.3372/wi.48.48207>
- ____, Harvey Y. & Onana J.-M. (2010). *The Plants of Dom, Bamenda Highlands, Cameroon, A Conservation Checklist*. Royal Botanic Gardens, Kew.
- ____, ____ & ____ (2011). *The Plants of Mefou Proposed National Park, Yaoundé, Cameroon, A Conservation Checklist*. Royal Botanic Gardens, Kew.
- ____, Nic Lughadha, E., Kirk, P., Lindon, H., Carretero, J., Looney, B., Douglas, B., Haelewaters, D., Gaya, E., Llewellyn, T., Ainsworth, M., Gafforov, Y., Hyde, K., Crous, P., Hughes, M., Walker, B. E., Forzza, R. C., Wong, K. M. & Niskanen, T. (2020). New scientific discoveries: plants and fungi. *Plants, People Planet* 2: 371 – 388. <https://doi.org/10.1002/ppp3.10148>
- ____, Onana, J.-M. & Pollard, B. J. (2000). *The Plants of Mount Oku and the Ijim Ridge, Cameroon, a Conservation Checklist*. Kew, Royal Botanic Gardens.
- ____, Pollard, B. J., Darbyshire, I., Onana, J.-M. & Wild, C. (2004). *The Plants of Kupe, Mwanenguba and the Bakossi Mountains, Cameroon: A Conservation Checklist*. Royal Botanic Gardens, Kew.
- ____, Tsukaya, H., Rudall, P. J. & Suetsugu, K. (2018a). Taxonomic monograph of *Oxygyne* (Thismiaceae), rare achlorophyllous mycoheterotrophs with strongly disjunct distribution. *PeerJ*, 6, e4828. <https://doi.org/10.7717/peerj.4828>
- ____ & Williams, S. (1999). A Review of African Saprophytic Flowering Plants. In: J. R. Timberlake & S. Kativu (eds), *African Plants. Biodiversity, Taxonomy & Uses. Proceedings of the 15th AETFAT Congress at Harare, Zimbabwe*, pp. 39 – 49. Royal Botanic Gardens, Kew.
- Couch, C., Molmou, D., Camara, B., Cheek, M., Merklinger, F., Davies, L., Harvey, Y., Lopez Poveda, L. & Redstone, S. (2014). Conservation of threatened Guinean inselberg species. In: XXth AETFAT Congress, South Africa, 2014 [Abstract 96]. *Scripta Bot. Belg.* (52: 96).
- ____, Cheek, M., Haba, P., Molmou, D., Williams, J., Magassouba, S., Doumbouya, S. & Diallo, M. Y. (2019). Threatened Habitats and Tropical Impor-

- tant Plant Areas (TIPAs) of Guinea, West Africa. Royal Botanic Gardens, Kew.
- Darbyshire, I. & Cheek, M. (2004a). *Rinorea faustiana*. *The IUCN Red List of Threatened Species* 2004: e.T46185A11034817. <https://doi.org/10.2305/IUCN.UK.2004.RLTS.T46185A11034817.en>.
- _____, & _____ (2004b). *Rinorea thomasi*. *The IUCN Red List of Threatened Species* 2004: e.T46186A11034916. <https://doi.org/10.2305/IUCN.UK.2004.RLTS.T46186A11034916.en>.
- _____, Anderson, S., Asatryan, A., Byfield, A., Cheek, M., Clubbe, C., Ghrabi, Z., Harris, T., Heatubun, C. D., Kalema, J., Magassouba, S., McCarthy, B., Milliken, W., Montmollin, B. de, Nic Lughadha, E., Onana, J. M., Saidou, D., Sarbu, A., Shrestha, K. & Radford, E. A. (2017). Important Plant Areas: revised selection criteria for a global approach to plant conservation. *Biodivers. Conservation* 26: 1767 – 1800. <https://doi.org/10.1007/s10531-017-1336-6>
- Darrah, S. E., Bland, L. M., Bachman, S. P., Clubbe, C. P. & Trias-Blasi, A. (2017). Using coarse-scale species distribution data to predict extinction risk in plants. *Diversity and Distributions* 23: 435 – 447. <https://doi.org/10.1111/ddi.12532>
- Harvey, Y., Pollard, B. J., Darbyshire, I., Onana, J.-M. & Cheek, M. (2004). *The Plants of Bali Ngemba Forest Reserve, Cameroon. A Conservation Checklist*. Royal Botanic Gardens, Kew.
- _____, Tchiengue, B. & Cheek, M. (2010). *The plants of the Lebiam Highlands, a conservation checklist*. Royal Botanic Gardens, Kew.
- Humphreys, A. M., Govaerts, R., Ficinski, S. Z., Lughadha, E. N. & Vorontsova, M. S. (2019). Global dataset shows geography and life form predict modern plant extinction and rediscovery. *Nature Ecol. Evol.* 3.7: 1043 – 1047. <https://doi.org/10.1038/s41559-019-0906-2>
- IPNI (continuously updated). *The International Plant Names Index*. Available from: <http://ipni.org/> [Accessed March 2021].
- IUCN (2012). *IUCN red list categories: Version 3.1*. IUCN Species Survival Commission, Gland and Cambridge.
- Juffe-Bignoli, D., Brooks, T. M., Butchart, S. H., Jenkins, R. B., Boe, K., Hoffmann, M., Angulo, A., Bachman, S., Böhm, M., Brummitt, N. & Carpenter, K. E. (2016). Assessing the cost of global biodiversity and conservation knowledge. *PLoS One* 11(8) <https://doi.org/10.1371/journal.pone.0160640>
- Kew Science News (2020). Ebo Forest Logging Plans Suspended. <https://www.kew.org/read-and-watch/ebo-forest-logging-suspended> [Accessed 5 May 2021].
- Lovell, R. (2020). Timber! The threat to Cameroon's Ebo Forest. <https://www.kew.org/read-and-watch/ebo-forest-cameroon> [Accessed 5 May 2021].
- Mair, L., Bennun, L. A., Brooks, T. M., Butchart, S. H., Bolam, F. C., Burgess, N. D., Ekstrom, J. M., Milner-Gulland, E. J., Hoffmann, M., Ma, K. & Macfarlane, N. B. (2021). A metric for spatially explicit contributions to science-based species targets. *Nature Ecol. Evol.* 5: 836 – 844. <https://doi.org/10.1038/s41559-021-01432-0>
- Maisels, F. M., Cheek, M. & Wild, C. (2000). Rare plants on Mount Oku summit, Cameroon. *Oryx* 34: 136 – 140. <https://doi.org/10.1017/s0030605300031057>.
- Nic Lughadha, E., Walker, B. E., Canteiro, C., Chadburn, H., Davis, A. P., Hargreaves, S., Lucas, E. J., Schuiteman, A., Williams, E., Bachman, S. P. & Baines, D. (2019). The use and misuse of herbarium specimens in evaluating plant extinction risks. *Philos. Trans., Ser. B*, 374 (1763), p.20170402. <https://doi.org/10.1098/rstb.2017.0402>
- _____, Bachman, S. P., Leão, T. C., Forest, F., Halley, J. M., Moat, J., Acedo, C., Bacon, K. L., Brewer, R. F., Gâteblé, G. & Gonçalves, S. C. (2020). Extinction risk and threats to plants and fungi. *Plants, People, Planet* 2 (5): 389 – 408. <https://nph.onlinelibrary.wiley.com/doi/10.1002/ppp3.10146>
- Onana, J.-M. (2011). *The vascular plants of Cameroon, a taxonomic checklist with IUCN Assessments*. Royal Botanic Gardens, Kew.
- _____, & Cheek, M. (2011). *Red data book of the flowering plants of Cameroon, IUCN global assessments*. Royal Botanic Gardens, Kew.
- Plants of the World Online (continuously updated). Facilitated by the Royal Botanic Gardens, Kew. http://www.plantsoftheworldonline.org/?f=accepted_names&q=Vepris (downloaded 1 May 2021)
- Sosef, M. S. M., Wieringa, J. J., Jongkind, C. C. H., Achoundong, G., Azizet Issembé, Y., Bedigian, D., Van Den Berg, R. G., Breteler, F. J., Cheek, M. & Degreef, J. (2006). Checklist of Gabonese Vascular Plants. *Scripta Bot. Belg.* 35. National Botanic Garden of Belgium.
- Thiers, B. (continuously updated). *Index Herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's Virtual Herbarium. Available from: <http://sweetgum.nybg.org/ih/> [Accessed March 2021].
- Turland, N. J., Wiersema, J. H., Barrie, F. R., Greuter, W., Hawksworth, D. L., Herendeen, P. S., Knapp, S., Kusber, W.-H., Li, D.-Z., Marhold, K., May, T. W., McNeill, J., Monro, A. M., Prado, J., Price, M. J. & Smith, G. F. (eds) (2018). International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Veg.* 159. Koeltz Botanical Books, Glashütten. <https://doi.org/10.12705/Code.2018>
- van Velzen, R. & Wieringa, J. J. (2014). *Rinorea calcicola* (Violaceae), an endangered new species from south-eastern Gabon. *Phytotaxa* 167 (3): 267 – 275. <https://doi.org/10.11646/phytotaxa.167.3.5>

- _____, Wahlert, G. A., Sosef, M. S., Onstein, R. E. & Bakker, F. T. (2015). Phylogenetics of African *Rinorea* (Violaceae): Elucidating Infrageneric Relationships using Plastid and Nuclear DNA Sequences. *Syst. Bot.* 40 (1): 174 – 184. <https://doi.org/10.1600/036364415X686486>
- Wahlert, G. A. (2010). *Phylogeny, biogeography, and a taxonomic revision of Rinorea (Violaceae) from Madagascar and the Comoro Islands*. Ph.D. Thesis. Ohio University, Athens.
- _____, & Ballard, H. E. (2012). A phylogeny of *Rinorea* (Violaceae) inferred from plastid DNA sequences with an emphasis on the African and Malagasy species. *Syst. Bot.* 37: 964 – 973. <https://doi.org/10.1600/036364412X656392>
- Walker, B. E., Leão, T. C., Bachman, S. P., Bolam, F. C. & Nic Lughadha, E. (2020). Caution needed when predicting species threat status for conservation prioritization on a global scale. *Frontiers Pl. Sci.* 11: 520. <https://doi.org/10.3389/fpls.2020.00520>
- Zizka, A., Silvestro, D., Vitt, P. & Knight, T. M. (2020). Automated conservation assessment of the orchid family with deep learning. *Conservation Biol.* <https://doi.org/10.1111/cobi.13616>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.