Correction to: Niche Partitioning among Snook (Pisces: Centropomidae) in Rivers of Southeastern Florida and Implications for Species Range Limits

Philip W. Stevens¹ · Jynessa Dutka-Gianelli¹ · Eric J. Nagid² · Alexis A. Trotter¹ · Kevin G. Johnson³ · Travis Tuten² · James A. Whittington⁴

Published online: 18 December 2019
© Coastal and Estuarine Research Federation 2019

Correction to: Estuaries and Coasts.

<https://doi.org/10.1007/s12237-019-00650-x>

In the original version of the article there was an error during the production of Fig. 3. The original article has been corrected and the corrected figure is as follows:

The online version of the original article can be found at <https://doi.org/10.1007/s12237-019-00650-x>

✉ Philip W. Stevens
philip.stevens@myfwc.com

¹ Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute, St. Petersburg, FL, USA

² Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute, Gainesville, FL, USA

³ Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute, Eustis, FL, USA

⁴ Florida Fish and Wildlife Conservation Commission, Tequesta Field Laboratory, Fish and Wildlife Research Institute, Tequesta, FL, USA

Fig. 3 Electrofishing captures (number per transect) of large-bodied (> 175 mm TL) common snook (triangles) and fat snook (circles) in the St Sebastian and Loxahatchee rivers. Also shown are capture locations of

snook using large haul seines along open estuarine shorelines of the Indian River Lagoon