CORRECTION

Correction to: Porcine Picornavirus 3C Protease Degrades PRDX6 to Impair PRDX6-mediated Antiviral Function

Congcong Wang¹ · Huanhuan Feng¹ · Xiangle Zhang¹ · Kangli Li¹ · Fan Yang¹ · Weijun Cao¹ · Huisheng Liu¹ · Lili Gao¹ · Zhaoning Xue¹ · Xiangtao Liu¹ · Zixiang Zhu¹ · Haixue Zheng¹

Published online: 12 November 2021
© Wuhan Institute of Virology, CAS 2021

Correction to: Virologica Sinica (2021) 36:948–957
<https://doi.org/10.1007/s12250-021-00352-4>

Due to our negligence, the original version of this article, published online on March 15, 2021, contained a mistake in Figure 2E (The Knockdown band of Western blotting was provided incorrectly). The correct Fig. 2E is given

below. We apologize for this error and state that this does not change the scientific conclusions of the article in any way.

Congcong Wang and Huanhuan Feng contributed equally to this work.

The original article can be found online at <https://doi.org/10.1007/s12250-021-00352-4>.

✉ Haixue Zheng
haixuezheng@163.com

✉ Zixiang Zhu
zhuzixiang@126.com

¹ State Key Laboratory of Veterinary Etiological Biology, OIE/ National Foot and Mouth Diseases Reference Laboratory, Key Laboratory of Animal Virology of Ministry of Agriculture, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou 730046, China

Fig. 2 PRDX6 inhibited FMDV replication. **A–C** PK-15 cells were transfected with 0, 0.5, 1 or 2 μ g of Myc-PRDX6 expressing plasmids for 24 h. The cells were then infected with FMDV (MOI = 0.5) for 12 h. The expression of viral proteins (**A**) and viral RNA (**B**) were detected by Western blotting and qPCR respectively. The virus yields were measured by TCID₅₀ assay (**C**). **D** EC-4 cells were transfected with 0, 1 or 2 μ g of Myc-PRDX6 expressing plasmids for 24 h. The cells were then mock-infected or infected with FMDV for 12 h. The expression of viral proteins and viral RNA were detected by Western blotting and qPCR respectively. **E** PK-15 cells were transfected with

120 nM of nontargeting control NC siRNA or PRDX6 siRNA (siRNA-329, siRNA-645 siRNA-729) for 36 h, the knockdown efficiency of each siRNA was then evaluated by qPCR and Western blotting analysis respectively. **F–H** PK-15 cells were transfected with 120 nM of NC siRNA or PRDX6 siRNA (siRNA-329) for 36 h, the cells were then infected with FMDV (MOI = 0.5) for 0, 4, 8 or 12 h. The expression of viral proteins (**F**) and viral RNA (**G**) were detected by Western blotting and qPCR respectively. The virus yields were measured by TCID₅₀ assay (**H**).