CORRECTION

Correction to: Different apoptotic effects of saxifragifolin C in human breast cancer cells

Kyung-Ho Kim¹ · Ji-Yun Kim¹ · Jong-Hwan Kwak¹ · Byung Oh Kim² ·
Suhkneung Pyo¹

Published online: 22 November 2018
© The Pharmaceutical Society of Korea 2018

Correction to: Arch. Pharm. Res. (2016) 39:577–589
<https://doi.org/10.1007/s12272-016-0729-5>

The authors regret that incorrect western band of Bax (MDA-MB-231) in Fig. 6a (right panel) was mistakenly uploaded in the original publication. The correct Fig. 6a is

shown below. This correction does not change the conclusions of this manuscript. The authors would like to apologize for any inconvenience caused.

The original article can be found online at <https://doi.org/10.1007/s12272-016-0729-5>.

✉ Suhkneung Pyo
snpyo@skku.edu

¹ School of Pharmacy, Sungkyunkwan University, Suwon, Kyunggi-do 440-746, Republic of Korea

² School of Food Science & Biotechnology, College of Agriculture & Life Sciences, Kyungpook National University, Daegu, Republic of Korea

Fig. 6 The role of ER α in saxifragifolin C-induced apoptosis. MCF-7 cells and MDA-MB-231 cells were transfected with vector control, vector inserted specific knockdown of ER α (shER α), or overexpression of ER α (HA-ER α) and then treated with Saxi C for 12 h. Total cells lysates were subjected to Western blot using ER α , Bcl₂, Bcl-xL, or Bax antibodies (a). β -Actin was used as an equal loading control. The levels ER α , Bcl₂, Bcl-xL, or Bax of expression are in arbitrary units, and data are normalized to respective amount of β -actin protein. Breast cancer cells were transfected with vector control, shER α (b), or HA-ER α (c) and then treated with Saxi C for 12 h. Saxi C-induced apoptosis was determined by Annexin V/PI staining. The Arabic number in each corner indicates the proportion of each quadrant