

## Erratum to: Effects of redox-active interlayer anions on the oxygen evolution reactivity of NiFe-layered double hydroxide nanosheets

Daojin Zhou<sup>1</sup>, Zhao Cai<sup>1</sup>, Yongmin Bi<sup>1</sup>, Weiliang Tian<sup>1,2</sup>, Ma Luo<sup>1</sup>, Qian Zhang<sup>1</sup>, Qian Zhang<sup>1</sup>, Qixian Xie<sup>1</sup>, Jindi Wang<sup>1</sup>, Yaping Li<sup>1</sup>, Yun Kuang<sup>1</sup>, Xue Duan<sup>1</sup>, Michal Bajdich<sup>3</sup>, Samira Siahrostami<sup>4</sup> (✉), and Xiaoming Sun<sup>1</sup> (✉)

<sup>1</sup> State Key Laboratory of Chemical Resource Engineering, College of Energy, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing 100029, China

<sup>2</sup> Key Laboratory of Chemical Engineering in South Xinjiang, College of Life Science, Tarim University, Alar 843300, China

<sup>3</sup> SUNCAT Center for Interface Science and Catalysis, SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA

<sup>4</sup> SUNCAT Center for Interface Science and Catalysis, Department of Chemical Engineering, Stanford University, Stanford, California 94305, USA


© Tsinghua University Press and Springer-Verlag GmbH Germany, part of Springer Nature 2019.

### Erratum to

Nano Research 2018, 11(3): 1358–1368

https://doi.org/10.1007/s12274-017-1750-9

The labels in Fig. 8 in the original version of this article were unfortunately misplaced. The corrected figure is as follow.



**Figure 8** Binding energy and Bader charge of (a) Ni and (b) Fe sites as a function of the experimentally measured onset potential in as-prepared LDHs (●: Binding energy; ▲: Bader charge).

The online version of the original article can be found at  
https://doi.org/10.1007/s12274-017-1750-9