Correction to: A comparison of clinicopathological characteristics and long-term survival outcomes between symptomatic and screen-detected breast cancer in Japanese women

Hitoshi Inari¹ · Satoru Shimizu¹ · Nobuyasu Suganuma¹ · Tatsuya Yoshida¹ · Hirotaka Nakayama² · Takashi Yamanaka¹ · Ayumi Yamanaka² · Yasushi Rino² · Munetaka Masuda²

Published online: 10 January 2018
© The Japanese Breast Cancer Society 2018

Correction to: Breast Cancer (2017) 24:98–103
<https://doi.org/10.1007/s12282-016-0678-3>

In the original publication of this article, Table 1 was published incorrectly with rows out of order under “Distant metastasis (*n*, %)”.

The correct Table 1 is given in the following page.

The original article can be found online at <https://doi.org/10.1007/s12282-016-0678-3>.

✉ Hitoshi Inari
inari-hitoshi@nifty.com

¹ Department of Breast and Endocrine Surgery, Kanagawa Cancer Center, 232 Nakao, Asahi-ku, Yokohama, Japan

² Department of Surgery, Yokohama City University, 39 Fukuura, Kanazawa-ku, Yokohama, Japan

Table 1 Demographic and clinicopathological characteristics by method of detection

Characteristics	Sympto- matic cancers (<i>n</i> = 977)	Screen-detected cancers (<i>n</i> = 182)	<i>P</i> value
Menopausal status (<i>n</i> , %)			
Pre	407 (42)	70 (38)	0.461
Post	570 (58)	112 (62)	
Age (<i>n</i> , %)			
< 50	341 (35)	60 (33)	0.339
≥ 50	636 (65)	122 (67)	
Distant metastasis (<i>n</i> , %)			
No	951 (97)	181 (99.4)	0.106
Yes	26 (3)	1 (0.6)	
Operation (<i>n</i> , %)			
Mastectomy	483 (49)	45 (25)	< 0.01
Partial resection	494 (51)	137 (75)	
Histology (<i>n</i> , %)			
Invasive carcinoma	919 (94)	146 (80)	< 0.01
Non-invasive carci- noma	58 (6)	36 (20)	
Invasive ductal carcinoma	781 (79)	123 (67)	< 0.01
Special types	138 (15)	23 (13)	
Others	58 (6)	36 (20)	
Tumor size (<i>n</i> , %)			
≤ 20 mm	398 (40)	124 (68)	< 0.01
≥ 20 mm	579 (60)	57 (31.5)	
Unknown	0	1 (0.5)	
Lymph node status (<i>n</i> , %)			
Negative	546 (56)	143 (79)	< 0.01
Positive	406 (42)	34 (19)	
Unknown	25 (2)	5 (2)	
Estrogen receptor (<i>n</i> , %)			
Negative	281 (29)	27 (15)	< 0.01
Positive	609 (62)	133 (73)	
Unknown	87 (9)	22 (12)	
Progesterone receptor (<i>n</i> , %)			
Negative	429 (44)	50 (27)	< 0.01
Positive	460 (47)	110 (60)	
Unknown	88 (9)	22 (13)	
Adjuvant chemotherapy (<i>n</i> , %)			
No	291 (30)	100 (55)	< 0.01
Yes	601 (61)	54 (30)	
Unknown	85 (9)	54 (30)	
Adjuvant hormone therapy (<i>n</i> , %)			
No	316 (32)	54 (29)	0.645
Yes	621 (63)	122 (67)	
Unknown	40 (5)	6 (4)	