



# Correction: Programmed death ligand 1-positive immune cells in primary tumor or metastatic axillary lymph nodes can predict prognosis of triple-negative breast cancer even when present at < 1% in the tumor region

Nobumoto Tomioka<sup>1</sup> · Kanako C. Hatanaka<sup>2</sup> · Dai Okuyama<sup>3</sup> · Ken-ichi Watanabe<sup>1</sup> · Mitsugu Yamamoto<sup>1</sup> · Hideki Maeda<sup>1</sup> · Hanae Tachikawa<sup>1</sup> · Sayuri Kuwahara<sup>1</sup> · Ai Shimizu<sup>4</sup> · Hiroaki Suzuki<sup>3</sup> · Yutaka Hatanaka<sup>2,5</sup> · Masato Takahashi<sup>6</sup>

Published online: 27 April 2023  
© The Author(s) 2023

## Correction: Breast Cancer

<https://doi.org/10.1007/s12282-023-01442-9>

In the original publication of the article, in table 2, the second column heading “1%” should be aligned to the value “& all>1%(25)” in third row. The correct Table 2 is given in this correction.

---

The original article can be found online at <https://doi.org/10.1007/s12282-023-01442-9>.

---

✉ Nobumoto Tomioka  
[nmhtomsfhcc@yahoo.co.jp](mailto:nmhtomsfhcc@yahoo.co.jp)

<sup>1</sup> Department of Breast Surgery, National Hospital Organization (NHO) Hokkaido Cancer Center, 4-2 Kikusui, Shiroishi-ku, Sapporo 003-0804, Japan

<sup>2</sup> Clinical Research and Medical Innovation Center of Development of Advanced Diagnostics, Hokkaido University Hospital, Sapporo, Japan

<sup>3</sup> Department of Clinical Pathology, National Hospital Organization (NHO) Hokkaido Cancer Center, Sapporo, Japan

<sup>4</sup> Department of Surgical Pathology, Hokkaido University Hospital, Sapporo, Japan

<sup>5</sup> Research Division of Genome Companion Diagnostics, Hokkaido University Hospital, Sapporo, Japan

<sup>6</sup> Department of Breast Surgery, Hokkaido University Hospital, Sapporo, Japan

**Table 2** DFS and OS of patients stratified according to the cut-off of 70 PD-L1(SP142)+ immune cells

70 cells		1%	DFS ( <i>p</i> )	HR / 95%CI	OS ( <i>p</i> )	HR / 95%CI
<70 cells (15 matched) vs	≥70 cells but <1% (15 matched)		0.004	0.140 0.030 – 0.657	0.020	0.195 0.042 – 0.904
All <70 cells (26) vs	All ≥70 cells but <1% (20)		<0.001	0.058 0.008 – 0.436	0.012	0.231 0.066 – 0.805
All <70 cells (26) vs	All ≥70 cells but <1% (20)	& all ≥1% (25)	<0.001	0.177 0.073 – 0.432	0.002	0.280 0.117 – 0.669
	All <1% (46) vs	All ≥1% (25)	0.138	0.503 0.199 – 1.270	0.154	0.491 0.181 – 1.332
	TILs <30% (55) vs	TILs ≥30% (16)	0.1111	0.506 0.216 – 1.189	0.004	0.308 0.132 – 0.719

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will

need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.