



# Awareness, Perception, and Practices Towards Blood Donation Among Undergraduate Health Science Students of India During the COVID-19 Pandemic

Nitin Joseph<sup>1,2</sup> · Sakshi Khaitan<sup>2,3</sup>

Received: 12 December 2021 / Accepted: 28 April 2022 / Published online: 19 May 2022  
© The Author(s), under exclusive licence to Indian Society of Hematology and Blood Transfusion 2022

**Abstract** To assess the awareness, perception, and practices of health science students towards blood donation during the COVID-19 pandemic. This cross-sectional study was done among the undergraduate medical, dental, physiotherapy, and audiology, speech and learning pathology students in May 2021. A self-administered questionnaire designed using Google Doc was used for data collection. Out of the 461 participants, only 171(37.1%) knew that Coronavirus was not transmitted through blood transfusion. Only 125(27.1%) participants knew that a minimum of 14 days is required before a donor who tested positive for COVID-19 can donate blood. As many as 339(73.5%) participants expressed their willingness to donate blood during the current pandemic. Having donated blood in the past ( $p = 0.001$ ), having vaccinated with COVID-19 vaccines ( $p = 0.029$ ), having taken both the vaccine doses ( $p = 0.0499$ ), and absence of anaemia ( $p = 0.0159$ ) were associated with willingness to donate blood during the pandemic. Only 83(18%) participants had donated blood after the onset of the pandemic. Out of the rest 378, 106(28%) participants did not donate blood due to the fear of getting infected with Coronavirus. Absence of chronic co-morbidities ( $p = 0.0288$ ) was associated with the history of having donated blood after the onset of COVID-19 pandemic among the participants. Awareness

of participants regarding certain key issues related to blood donation and COVID-19 were found lacking. Counselling services to alleviate fears associated with blood donation and awareness sessions to remove misconceptions are required among students to improve blood donation practices.

**Keywords** Blood donation · Health science students · Awareness · Perception · Practice

## Introduction

The ongoing COVID-19 pandemic has disrupted health care and community operations worldwide. It has led to a shortage of essential medicines, equipment for personal protection, hospital beds, and health care personnel [1]. However, another lifesaving component that has not been adequately projected is the shortage of blood donations leading to a global shortage in the blood supply [1, 2].

The drop in voluntary blood donors has been reported in several European and Asia studies following the COVID-19 outbreak [3–6]. Number of stakeholders involved in managing blood centres and blood transfusion services have raised concerns regarding maintaining the safety of restrained gathering and social distancing during blood donation activities [7].

A number of other measures must be practised to ensure safety in the blood donation procedure during the current circumstances. The staff, donors, and visitors need to wear facemasks throughout, and utmost care needs to be taken to screen donors to ensure that people with any symptoms suggestive of illness are prevented from donating blood [7–9]. According to the Occupational Safety and Health Administration guidelines, staff at the blood donation

✉ Nitin Joseph  
drnitinjoseph@gmail.com

Sakshi Khaitan  
khaitansakshi1999@gmail.com

<sup>1</sup> Department of Community Medicine, Kasturba Medical College, Mangalore, India

<sup>2</sup> Manipal Academy of Higher Education, Manipal, India

<sup>3</sup> Kasturba Medical College, Mangalore, India

centres need to also wear gloves and change gloves and wash hands between contacts from one donor to another. After each donor has vacated the station, disinfection procedures must be practiced consistently before setting the station for the new donor. All the equipment required for the blood donation process needs to be sterilized, and most of these should preferably be of a single-use type. [9] A gap of at least 6 feet needs to be ensured in the donors' waiting area and between blood donation cots during the bleeding procedure [7, 10].

All the donors need to ensure that a gap of at least 14 days is to be maintained after a positive COVID-19 test result [11]. Blood and its components collected from the donor within 14 days of disease onset need to be recalled [12].

Hence, several safety precautions must be ensured at blood donation centres to avoid COVID-19 transmission. The anxiety associated with the fear of getting infected with Coronavirus due to these concerns has been reported to be a major deterrent to the altruistic behaviour of blood donation [3, 4]. Added to this, the curbs in movement following a nationwide lockdown and the social distancing advised for preventing COVID-19 spread are being interpreted by the public as to not congregate for blood donation opportunities [1, 7, 13].

On the other hand, the pandemic has created a situation wherein adequate blood reserves are required for its control. Neutralizing antibodies in convalescent plasma donated by patients who recently recovered from this disease has been found to reduce the viral load and improve the prognosis among severe COVID-19 patients [14].

Thus with day to day rise in the number of active patients with COVID-19, the requirement of blood supplies and donated plasma for case management has become a global concern [11, 15]. The shortage of blood has affected the management of patients with other illnesses too. It has threatened the life of patients dependent on blood transfusion as a life-saving measure. This comprises patients with Thalassemia, ones who are critically ill, and accident victims [7]. Shortage of blood has also resulted in the postponement of elective surgeries [16].

Therefore, maintaining adequate stock of blood supply is a paramount concern during the ongoing pandemic. Reliable blood reserves can be ensured if there is regular blood donation practice among the eligible population. There have been reports of misconceptions about the transmission of COVID-19 among blood donors that have affected the blood donation rate [1]. There have also been concerns about minimal opportunities to donate, and concerns like feeling unfit to donate blood among university students in prior studies [17, 18]. Therefore, the current focus should be on building a sustainable platform for blood donors by improving awareness of its importance

and correcting misconceptions. This should be equally complemented with an effective program for voluntary blood donors [19].

Health professional students form a potential pool of eligible blood donors [19]. Therefore, it is essential to study their motivation toward blood donation and how it has been impacted during the phase of this widespread illness. Therefore, this study was done to assess the awareness, perception, and practices of health science students towards blood donation during the ongoing COVID-19 pandemic.

## Materials and methods

This cross-sectional study was done among undergraduate Medical, Dental, Physiotherapy, and Audiology, Speech and Language Pathology students of a Private University in Mangalore city of South India. The institutional ethics committee approved the study protocol and the study questionnaire. The Deans of the respective institutions gave permission. The study was conducted in May 2021 during the peak of the second COVID-19 wave, when several social restrictions and lockdown measures were in full swing.

In a study in Kangra, India [20], among medical students, 22.9% had donated blood in the past. Based on this finding, using the formula  $Z\alpha^2pq/d^2$ , the sample size comes to 323 at 95% confidence interval and 80% power. Adding 10% to incorporate non-response rate, the final sample size was taken as 355 participants. Only participants aged 18 years or above were included in this study. Non-consenting participants were excluded from this study. A self-administered questionnaire designed using Google Doc was used for data collection. It was content validated with the help of faculty members from the Department of Medical Education. It was further pilot tested among five students from each of the disciplines. After a few minor modifications following the pilot study, the final questionnaire was circulated among the students using WhatsApp.

The study information sheet and the consent form were at the questionnaire's start. Participants not consenting had the option to decline participation, leading to the submission of an empty questionnaire.

The questionnaire was semi-structured and was prepared using a review of related literature available from online sources. The first section of the questionnaire inquired about the basic demographic details of the participants. Other information such as underlying co-morbidities, history of allergic disorders, medication history, whether suffered from COVID-19 in the past and history of having

taken single or both doses of COVID-19 vaccines were also enquired.

Each of the other three sections enquired about awareness, perception, and practices towards blood donation among the participants.

Deferral from blood donation was taken as at least 14 days for those who tested positive for COVID-19 [21]. The minimum time gap between blood donations was taken as 3 months [22]. The period up to which individuals who received a COVID-19 vaccine can donate convalescent plasma was taken as 6 months [21]. Height and weight were self-reported by the participants. Body mass index (BMI) was categorized as per the Asia–Pacific classification [23].

Submitted responses were extracted from Google forms to MS Excel and were then transferred to IBM SPSS for Windows version 25.0, Armonk, New York for data analysis. The Chi-square test and Fisher's Exact test were used for testing association. The significance level was set at 5 percent.

The Cronbach's alpha value of the reliability of the questionnaire was found to be 0.868, indicating good internal consistency.

## Results

Out of the total 461 participants, 364(79%) were MBBS, 79(17.1%) were BDS, 10(2.2%) were BPT, and 8(1.7%) were BASLP students. The mean age and standard deviation of the participants were  $21.9 \pm 1.6$  years. Majority of them were females [279(60.5%)] and were native of urban areas [381(82.6%)]. 356(77.2%) participants were from nuclear, 70(15.2%) were from joint, and 35(7.6%) were from three generation families.

Only 171(37.1%) participants knew that the Coronavirus is not transmissible through blood transfusion. Only 125(27.1%) participants knew that a minimum of 14 days is required before a donor who tested positive for COVID-19 can donate blood (Table 1). Awareness regarding good practices to be followed by donors prior to donating blood such as taking sufficient rest, avoidance of alcohol consumption, avoidance of smoking, drinking plenty of water, taking iron supplements, and taking multi-vitamin supplements was known to 379(82.2%), 385(83.5%), 365(79.2%), 409(88.7%), 209(45.4%) and 189(41%) participants respectively.

As many as 359(77.9%) participants felt that the lack of blood donation by people during the current COVID-19 pandemic is leading to the shortage of blood for the patients. As many as 339(73.5%) participants expressed their willingness to donate blood during the current pandemic. The majority [271(79.9%)] thought so to help

severely ill COVID-19 patients requiring blood components. 70(15.2%) were not sure, and the rest 52(11.3%) participants were not willing to donate blood during the current circumstances. Out of these 52 participants, 35(67.3%) felt so due to the feeling that social distancing measures are not adequately practiced at the blood donation centres (Table 2).

As many as 100(21.7%) participants had donated blood at any time in the past. The most common reasons for not donating blood among the rest 361 participants were age below 18 years [39(10.8%)], and lack of opportunity to donate blood [35(9.7%)]. 378 (82%) participants had not donated blood after the onset of the COVID-19 pandemic. Among them, 106 (28%) had not donated blood due to the fear of getting infected with Coronavirus (Table 3).

The other narrative comments given by participants were: awareness regarding the importance of blood donation needs to be propagated among people, particularly during the COVID-19 pandemic (by 4), blood donors after every blood donation need to be appreciated with non-monetary rewards (by 1), full aseptic precautions at hospitals need to be ensured during the COVID-19 pandemic for the safety of the blood donors (by 1), all voluntary blood donors need to register their names at the blood banks so that the donors can be informed as and when the need of blood arises (by 1), and people need to be made aware that donating blood is a safe procedure with no adverse effects provided the donors fulfill the eligibility criteria for blood donation (by 1).

Chronic co-morbidities were present among 40(8.7%) participants. Majority of them had allergic disorders (among 23), followed by anaemia (among 7), followed by asthma (among 6), and epilepsy (among 3) in addition to other co-morbidities. 101(21.9%) participants reported history of allergy to any substance. Blood thinners and anti-cancer drugs were taken by 2 participants each.

410(88.9%) participants were vaccinated for COVID-19. The most common vaccines taken by them were Covishield [373(91%)] and Covaxin [34(8.4%)]. Moderna, Pfizer, and Sinopharm was taken by 1(0.2%) participant each.

Having donated blood in the past, having been vaccinated with COVID-19 vaccines, having taken both the vaccine doses, and absence of anaemia were associated with willingness to donate blood during the current pandemic among the participants (Table 4).

Male participants and those without chronic co-morbidities donated blood significantly more in the past in comparison to others. Absence of chronic co-morbidities was associated with a history of having donated blood after the onset of the COVID-19 pandemic among the participants (Table 5).

**Table 1** Awareness of participants regarding issues concerning blood donation during the COVID-19 pandemic (n = 461)

Awareness regarding	Number	Percentage
<i>Coronavirus is transmissible by blood transfusion</i>		
No	171	37.1
<i>Minimum duration of postponement of blood donation for a donor who tested positive for COVID-19</i>		
14 days	125	27.1
<i>Deferral time for donors who were vaccinated with an inactivated viral vector vaccine or other RNA based vaccines</i>		
There is no deferral time	21	4.5
<i>Individuals who received a COVID-19 vaccine can donate convalescent plasma</i>		
Yes, they can	182	39.5
<i>If yes, within how many months? (n = 182)</i>		
6 months	13	7.1
<i>COVID-19 patients after complete recovery have plasma antibodies that can help other patients with life-threatening COVID-19</i>		
Yes	306	66.4
<i>Awareness regarding the frequency of blood donation</i>		
Once in 3 months	20	4.3
<i>Blood donors need to be screened for COVID-19</i>		
No need	36	7.8

**Table 2** Perception towards blood donation among the participants during the COVID-19 pandemic (n = 461)

Perception towards	Number	Percentage
<i>Lack of blood donation by people during the current COVID-19 pandemic is leading to a shortage of blood for patients</i>		
Yes	359	77.9
<i>Willingness to donate blood during the current pandemic if the need arises</i>		
Yes	339	73.5
<i>Reasons stated for expressing willingness towards blood donation during the current pandemic (n = 339)*</i>		
To help severely ill COVID-19 patients requiring blood components	271	79.9
Being physically fit with good blood count	199	58.7
To meet blood requirements at various health facilities	124	36.6
Inspired by other blood donors who are friends or relatives	55	16.2
<i>Reasons stated for not expressing willingness towards blood donation during the current pandemic (n = 52)*</i>		
Lack of social distancing measures at blood donation centres	35	67.3
Blood donation makes one feel physically weak	32	61.5
Fear of acquiring infection with COVID-19	24	46.1
Lack of trust in aseptic measures practiced at the blood donation centres	22	42.3
<i>Blood donation should be promoted among people during the prevailing COVID-19 pandemic</i>		
Yes	376	81.6
<i>Willingness to promote blood donation drive among family members and friends</i>		
Yes	364	79.0

\*Multiple responses

## Discussion

The need to improve awareness regarding blood donation among health science students features among the list of recommendations framed by the World Health

Organization [17]. They being a young population, constitute an eligible group for donating blood.

Just over one-third of the participants in this study knew that Coronavirus is not transmissible through blood and blood products. This indicates inadequate awareness

**Table 3** Blood donation practices among the participants (n = 461)

Practices	Number	Percentage
<i>Donated blood in the past</i>		
Yes	100	21.7
<i>Reasons for not having donated blood so far (n = 361)*</i>		
Was below 18 years of age and hence was not eligible	39	10.8
Lack of opportunities	35	9.7
Due to anaemic status in the past	34	9.4
Due to underweight status	12	3.3
<i>History of donating blood after the onset of COVID-19 pandemic</i>		
Yes	83	18.0
<i>Reason for not donating blood after the onset of COVID-19 pandemic (n = 378)*</i>		
Due to fear of getting infected with Coronavirus	106	28.0
Lack of opportunities	22	5.8
Due to lockdown	8	2.1

\*Multiple responses

**Table 4** Factors associated with willingness to donate blood during the current COVID-19 pandemic among the participants

Factors	Willingness to donate blood during the current pandemic		Total	Chi-square value, <i>p</i> value
	Yes(%)	No/Not sure(%)		
<i>Coronavirus is transmissible by blood transfusion</i>				
No	133(77.8)	38(22.2)	171	$X^2 = 2.514, p = 0.113$
Yes/Not sure	206(71.0)	84(29.0)	290	
<i>Having donated blood in the past</i>				
Yes	87(87)	13(13)	100	$X^2 = 11.896, p = 0.001$
No	252(69.8)	109(30.2)	361	
<i>Current status of anaemia</i>				
Yes	2(28.6)	5(71.4)	7	$p = 0.0159$
No	337(74.2)	117(25.8)	454	
<i>COVID-19 vaccine</i>				
Taken	308(75.1)	102(24.9)	410	$X^2 = 4.791, p = 0.029$
Not taken	31(60.8)	20(39.2)	51	
Total	339	122	461	
<i>Number of doses of COVID-19 vaccines taken so far</i>				
One	86(68.8)	39(31.2)	125	$X^2 = 3.846, p = 0.0499$
Two	222(77.9)	63(22.1)	285	
Total	311	102	410	

among students regarding the transmission of COVID-19. Awareness was however not associated with their perception towards willingness to donate blood. Even then, it becomes essential to correct this misconception among the health science students so that they provide correct information to others. Communication promotes a positive blood donation attitude and helps sustain the blood donor program by encouraging the first donation and repeat donations thereafter [24].

About three in four participants in this study expressed their willingness to donate blood during the ongoing pandemic. The majority wanted to donate blood to help COVID-19 patients with severe illnesses who require blood components. Similar observations were made in a study done in USA [19] where 91% of medical students expressed their willingness to donate blood, despite the pandemic. This reflects strong altruism among young medical students to donate blood in times of medical emergencies. In another study done in Sudan, majority of

**Table 5** Factors associated with the history of having donated blood in the past/after the onset of COVID-19 among the participants

Factors	Donated blood in the past		Total	Chi-square value, <i>p</i> value
	Yes(%)	No(%)		
<i>Gender</i>				
Males	58(31.9)	124(68.1)	182	$X^2 = 18.333, p < 0.001$
Females	42(15.1)	237(84.9)	279	
<i>Presence of chronic co-morbidities</i>				
Yes	3(7.5)	37(92.5)	40	$p = 0.0254$
No	97(23.0)	324(77.0)	421	
Total	100	361	461	
Factors	Donated blood after the onset of the COVID-19 pandemic		Total	<i>p</i> value
	Yes(%)	No(%)		
<i>Presence of chronic co-morbidities</i>				
Yes	2(5.0)	38(95.0)	40	$p = 0.0288$
No	81(19.2)	340(80.8)	421	
Total	83	378	461	

participants (63.4%) were willing to donate blood during the pandemic if they got regular invitations. Other reasons stated for agreeing to blood donation in the same study were: control of the pandemic (52.4%), provision of better transportation facilities (29.6%), and provision of incentives for donating blood (6.5%) by the participants [25]. Provision of donor transportation facilities, switching sites for blood collections, and scheduling appointments for blood donations feature among the suggested measures by the WHO to ensure safety during blood donation [12].

As many as 11.3% of participants in this study were not willing to donate blood during the current pandemic. The most common reason being concern regarding inadequate social distancing measures at blood donation centres. Such concerns need to be alleviated among students by reassuring them of the various safety precautions being implemented at the blood donation centres.

In the present study, factors associated with willingness to donate blood during the current pandemic among the participants were: history of having donated blood in the past, absence of anaemia, having taken the COVID-19 vaccine, and having taken both COVID-19 vaccine doses. This was different from the findings of the study done in USA, where having donated blood in the past was not associated with the present willingness to donate blood during the current pandemic [19].

In relation to the practice of donating blood, as many as 82% participants in the present study did not do so after the onset of the COVID-19 pandemic. The most common reason for this being the fear of getting infected with Coronavirus, followed by the lack of opportunities to donate blood. This meant that counselling services and

more opportunities need to be given by the blood banks to improve the participation of medical students in blood donation activities during the current pandemic. In other studies, reasons for lapse from blood donation during the ongoing pandemic were: to avoid Coronavirus infection (60.6%) [25], time constraints (40.2%) [25], transportation problems (32.5%) [25], inconvenience to travel to the blood donation centre (6.1%) [25], medical ineligibility (6.5%) [25], inconvenient timings of the blood bank centre (4.2%) [25], concern about their personal safety (3%) [19], and fear of needles or blood (1.2%) [25]. The various donation-related fears might adversely affect the recruitment and retention of donors. Mass media can play an important role here by removing misconceptions and reaffirming the safety of the blood donation procedure among the donors to maintain an adequate blood supply [1]. Blood collection centres also need to ensure minimum donor exposure while procuring blood [1]. The presence of mobile blood collection units might be a solution to address transportation problems causing barriers to blood donation [25].

To enhance the involvement of medical students as donors, in campus blood donation drives need to be organized by following all COVID-19 appropriate preventive measures. This will ensure accessibility among donors despite the restricted mobility imposed by the lockdown [1]. A registry needs to be prepared with a list of all interested students, and it needs to be shared with the local blood banks. This will ensure adequate opportunities to donate blood [19]. Added to this, regular notifications need to be given to students through social media, WhatsApp, and emails about upcoming blood donation events [19].

Social media networks were reported to be an effective medium for sharing blood donation requests by Abbasi et al. [26].

In studies done in Saudi Arabia [6] and China [4], 39.5% and 67% reductions were seen in the number of blood donors after the COVID-19 outbreak. In the European study, half of the blood donors declared that they had donated less than usual frequency after the pandemic onset. Participants in the European study who anticipated high risk of infection and those who were adherent to COVID-19 appropriate behaviour like social distancing measures were significantly less likely to donate blood. History of more blood donation activities in the recent 2 years, young age and participants employed in educational or health sectors were more likely to donate blood during the current crisis [27]. In the present study, the absence of chronic comorbidities among the participants was associated with the history of having donated blood after the onset of the COVID-19 pandemic.

## Conclusion

Awareness of participants about certain key issues related to blood donation and COVID-19 was lacking. However, three-fourths of them expressed their willingness to donate blood. More than 80% of the participants had not donated blood after the onset of the pandemic. This was mainly because of the fear of getting infected with Coronavirus. Hence, counselling services and awareness sessions are required among students to improve their blood donation practices. The factors identified to be associated with willingness and reluctance to donate blood in this study are relevant in the context of this pandemic. These need to be addressed to improve blood donation in the setting.

## Limitations

There is a possibility of not disclosing information related to the perception and practices towards blood donation among the participants in this study.

**Acknowledgements** The authors of this study would like to thank the students who enthusiastically participated in this study.

**Author's Contribution** NJ: guarantor of this research work, concept, design, literature search, data collection tool preparation, data analysis, interpretation of data, manuscript preparation, revising the work critically for important intellectual content. SK: literature search, data collection, manuscript editing, revising the work critically for important intellectual content. This manuscript has been read and approved by all the authors.

**Funding** No source of funding was acquired for this research study.

## Declarations

**Conflict of interest** The authors declare that they have no competing interests.

**Ethics Approval** This was obtained from the institutional ethics committee with reference number IEC KMC MLR 05–2021/179 dated 20<sup>th</sup> May 2021 and was in accordance with the 1964 Helsinki declaration.

**Informed Consent** Written informed consent to participate was taken from the participants.

**Data Protection, Confidentiality and Privacy** The participants were informed that the data generated as a part of this study would be used for research purposes only. Complete anonymity of the information provided was ensured to them.

**Consent to Publish** The participants were also informed that the data generated as a part of this study would be used for publication.

## References

1. Kumar S, Azim D, Nasim S, Hashmi SH (2020) Dwindling blood reserves: An ominous downside of COVID-19 pandemic. *Transfus Apher Sci* 59:102818
2. Pan American Health Organization. PAHO warns of potential blood shortages during the COVID-19 pandemic. <https://www.paho.org/en/news/10-4-2020-paho-warns-potential-blood-shortage-during-covid-19-pandemic> Updated 10 Apr 2020; Accessed 6 May 2021
3. Leung JNS, Lee C (2020) Impact of the COVID-19—a regional blood centre's perspective. *ISBT Sci Ser* 15:362–364
4. Wang Y, Han W, Pan L, Wang C, Liu Y, Hu W, Zhou H, Zheng X (2020) Impact of COVID-19 on blood centres in Zhejiang province China. *Vox Sang* 115:502–506
5. The Local. Blood reserves in Germany run critically short due to corona crisis. <https://www.thelocal.de/20200520/blood-reserves-run-critically-short-due-to-corona-epidemic/>. Accessed 14 May 2021
6. Yahia AIO (2020) Management of blood supply and demand during the COVID-19 pandemic in King Abdullah Hospital, Bisha. *Saudi Arabia Transfus Apher Sci* 59:102836
7. National Blood Transfusion Council. Ministry of Health and Family Welfare, Government of India. National guide to blood transfusion services in India in light of COVID-19 pandemic. <https://www.mohfw.gov.in/pdf/NBTCGUIDANCEFORCOVID19.pdf> Updated 25 Mar 2020; Accessed 16 Feb 2022
8. Memorial Blood Centers. Coronavirus and blood donation. <https://www.mbc.org/donate-blood/coronavirus-blood-donation/> Updated 18 Aug 2021; Accessed 15 Feb 2022
9. Stanford Blood Center. FAQ on COVID-19 and blood donation. <https://stanfordbloodcenter.org/faq-on-novel-coronavirus-and-blood-donation/> Updated 2 Feb 2022; Accessed 20 Feb 2022
10. Centers for Disease Control and Prevention. Guidance for Blood and Plasma Facilities. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/blood-and-plasma-collection.html> Updated 29 Apr 2020; Accessed 20 Feb 2022
11. New York Blood Center. COVID-19 and blood donation. <https://www.nybc.org/donate-blood/covid-19-and-blood-donation-copy/#:~:text=DONORS%20PRESENTING%3A,-Regardless%20of%20vaccination&text=mouth%20and%20nose-,Must%20be%2014%20days%20symptom%20free%20if%20>

- recovered%20from%20COVID,to%20exposure%20risk%20C%20including%20travel Accessed 28 Feb 2022
12. World Health Organization. Maintaining a safe and adequate blood supply during the pandemic outbreak of coronavirus disease (COVID-19). <https://apps.who.int/iris/bitstream/handle/10665/331523/WHO-2019-nCoV-BloodSupply-2020.1-eng.pdf?ua=1> Updated 20 Mar 2020; Accessed 15 Feb 2022
  13. Zorlu F. World facing blood shortages due to COVID-19 pandemic. <https://www.aa.com.tr/en/latest-on-coronavirus-outbreak/world-facing-blood-shortages-due-to-covid-19-pandemic/1802664> Cited 13 Apr 2020; Accessed 18 May 2021
  14. Duan K, Liu B, Li C, Zhang H, Yu T, Qu J et al (2020) Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proc Natl Acad Sci USA* 117:9490–9496
  15. The American National Red Cross. American Red Cross Faces Severe Blood Shortage As Coronavirus Outbreak Threatens Availability of Nation's Supply. <https://www.redcross.org/about-us/news-and-events/press-release/2020/american-red-cross-faces-severe-blood-shortage-as-coronavirus-outbreak-threatens-availability-of-nations-supply.html> Updated 17 May 2020; Accessed 2 May 2021
  16. Miller H. Blood Donations Needed During Coronavirus Pandemic. CNBC; 2020 <https://www.cnbc.com/2020/03/23/blood-donations-needed-during-coronavirus-pandemic.html> Updated 23 Mar 2020; Accessed 18 May 2021
  17. Alsalmi MA, Almalki HM, Alghamdi AA, Aljasir BA (2019) Knowledge, attitude and practice of blood donation among health professions students in Saudi Arabia; A cross-sectional study. *J Family Med Prim Care* 8:2322–2327
  18. Raghuvanshi B, Pehlajani NK, Sinha MK (2016) Voluntary Blood Donation among Students - A Cross-Sectional Study on Knowledge and Practice vs. Attitude. *J Clin Diagn Res.*;10:EC18-EC22
  19. Khatun R, Otaibi BW, Ssentongo A, Hazelton JP, Cooper A (2021) Medical Student Attitudes Toward Blood Donation in Times of Increased Need. *Am Surg*, p 31348211011083
  20. Chauhan R, Kumar R, Thakur S (2018) A study to assess the knowledge, attitude, and practices about blood donation among medical students of a medical college in North India. *J Family Med Prim Care* 7:693–697
  21. American Red Cross. What to know about the Corona virus and Blood donation. <https://www.redcrossblood.org/donate-blood/dlp/coronavirus-covid-19-and-blood-donation.html> Accessed 26 Apr 2021
  22. Indian Red Cross Society. Blood Centre. <https://indianredcross.org/ircs/program/BloodBank#:~:text=The%20minimum%20time%20advised%20between,regain%20the%20normal%20haemoglobin%20count.&text=The%20age%20of%20the%20donor,less%20than%2012.5%20g%2Fdl>. Accessed 26 Apr 2021
  23. Lim JU, Lee JH, Kim JS, Hwang YI, Kim TH, Lim SY et al (2017) Comparison of World Health Organization and Asia-Pacific body mass index classifications in COPD patients. *Int J Chron Obstruct Pulmon Dis* 12:2465–2475
  24. Lownik E, Riley E, Konstenius T, Riley W, McCullough J (2012) Knowledge, attitudes and practices surveys of blood donation in developing countries. *Vox Sang* 103:64–74
  25. Sayedahmed AMS, Ali KAM, Ali SBS, Ahmed HSM, Shrif FSM, Ali NAA (2020) Coronavirus disease (COVID-19) and decrease in blood donation: A cross-sectional study from Sudan. *ISBT Sci Ser* 15:381–385
  26. Abbasi RA, Maqbool O, Mushtaq M, Aljohani NR, Daud A, Alowibdi JS et al (2018) Saving lives using social media: Analysis of the role of twitter for personal blood donation requests and dissemination. *Telemat Inform* 35:892–912
  27. Chandler T, Neumann-Bohme S, Sabat I, Barros PP, Brouwer W, van Exel J, et al (2021) Blood donation in times of crisis: Early insight into the impact of COVID-19 on blood donors and their motivation to donate across European countries. *Vox Sang*. <https://doi.org/10.1111/vox.13103>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.