



LETTER

# Comment on: “Retrospective Claims Analysis Indirectly Comparing Medication Adherence and Persistence Between Intravenous Biologics and Oral Small-Molecule Therapies in Inflammatory Bowel Diseases”

John Woolcott · Joseph C. Cappelleri · Puza Sharma · Irene Modesto

Received: December 4, 2019 / Published online: March 15, 2020  
© The Author(s) 2020

## Key Summary Points

The authors of Moran et al. used mutually exclusive IBD and RA infliximab treatment data to bridge differences across the unique inflammatory diseases, concluding that, after adjustment, adherence was higher with infusions than oral medications.

The analysis by Moran et al. does not take into account a number of important factors:

1. No discussion of reasons for discontinuation of treatment, nor the important differences between RA and IBD patient populations, was included in the manuscript.

2. Previous studies have shown wide variability in the concept of adherence, as well as its measurement, which could affect the conclusions of a study such as that presented by Moran et al.

3. Real-world data comparing tofacitinib with common biologics for the treatment of RA reported persistence and adherence of tofacitinib were at least comparable to that of the biologics.

We believe that there are pitfalls associated with the indirect method applied by Moran et al. and their results should be interpreted with caution.

Dear Editor,

We read with great interest the article by Moran et al. entitled “Retrospective Claims Analysis Indirectly Comparing Medication Adherence and Persistence Between Intravenous Biologics and Oral Small-Molecule Therapies in Inflammatory Bowel Diseases” [1].

In this retrospective cohort analysis of a claims database of adult patients diagnosed with either inflammatory bowel disease (IBD) or rheumatoid arthritis (RA), the authors investigate adherence and persistence with respect to vedolizumab in IBD, tofacitinib in RA, or

---

**Enhanced Digital Features** To view enhanced digital features for this article go to <https://doi.org/10.6084/m9.figshare.11836269>.

---

J. Woolcott  
Pfizer Inc, Collegeville, PA, USA

J. C. Cappelleri  
Pfizer Inc, Groton, CT, USA

P. Sharma · I. Modesto (✉)  
Pfizer Inc, New York, NY, USA  
e-mail: Irene.Modesto@pfizer.com

infliximab in RA or IBD treatment. Using mutually exclusive IBD and RA infliximab treatment data to bridge differences across the unique inflammatory diseases, the authors conclude that, after adjustment, adherence was higher with infusions than oral medications [1]. These results are in contrast to findings from previous well-conducted studies [2, 3]. Furthermore, the analysis by Moran et al. does not take into account a number of important factors, and we suggest relies on questionable methodology. These limitations cast doubt on the validity of their findings and overall conclusions, which we believe should be brought to the attention of the authors and your readers.

Although the authors of the paper noted that “there are several reasons for discontinuation that pertain to each disease” as a limitation of the study [1], no discussion of reasons nor the important differences between RA and IBD patient populations was included in the manuscript, such as age of the patients, presence of comorbidities, and number of concomitant therapies. Indirect comparisons utilizing observational studies, such as that described in Moran et al., are uncommon, since the heterogeneity of patient populations in the real world make such comparisons difficult. The study also failed to recognize tofacitinib dosing differences between the two diseases, both in terms of dose strength and overall posology. In accordance with US prescribing information, the recommended tofacitinib dose for RA is 5 mg twice daily (BID) or 11 mg once daily, whereas for ulcerative colitis (UC), the recommended dose is 10 mg BID for induction (8 weeks, continue for a maximum of 16 weeks if needed) followed by 5 mg BID or 10 mg BID for maintenance (use of 10 mg BID beyond induction should be limited and used for the shortest duration) [4]. Furthermore, for tumor necrosis factor inhibitors, including infliximab, real-world data have shown that changes in dose and dose schedules are more common for patients with IBD vs those with RA [5, 6], highlighting the complexity involved in comparing the same therapies across different disease populations. It is also noteworthy that tofacitinib is indicated for UC, in contrast to vedolizumab and infliximab, which have indications for both Crohn’s disease

and UC, reiterating the inappropriateness of these comparisons, which included patients with UC and also patients with Crohn’s disease.

A systematic review of 24 studies of RA, spondyloarthritis, and psoriatic arthritis examined adherence to biologic therapies and concluded that there was wide variability in the concept of adherence as well as in its measurement [7]. The choice of methods used might therefore be expected to affect the conclusions of a study such as that presented in Moran et al. Of note, although two methods were used to evaluate adherence, significant differences between vedolizumab/IBD and tofacitinib/RA were observed only for one of them after the adjustment method was applied [1].

Finally, published data on the concept of persistence and adherence with tofacitinib have demonstrated 2- and 5-year estimated drug survival rates of 75.5% and 49.4%, respectively, in a clinical trial setting [2], while real-world data comparing tofacitinib with common biologics (adalimumab, etanercept, and abatacept) for the treatment of RA reported persistence, and adherence of tofacitinib was at least comparable to that of the biologics [3].

As stated in the article by Moran et al., their results are not generalizable and need to be confirmed in tofacitinib-treated IBD patients. In the absence of direct study in tofacitinib-treated patients with UC, we believe that there are pitfalls associated with the indirect method applied by Moran et al. (as noted by the authors themselves) that amount to an uneven evaluation and comparison with potential for bias, and therefore the results should be interpreted with caution. This should be brought to the attention of your readership and prescribers.

Sincerely,

John Woolcott, PhD; Joseph C. Cappelleri MS, MPH, PhD; Puza Sharma, MD, MPH, PhD; Irene Modesto, MD, PhD

## ACKNOWLEDGMENTS

**Funding.** J Woolcott, JC Cappelleri, P Sharma, and I Modesto are employees and

shareholders of Pfizer Inc. Medical writing support was funded by Pfizer Inc. No Rapid Service or Open Access Fee was received by the journal for the publication of this letter.

**Medical Writing Assistance.** Medical writing support, under the guidance of the authors, was provided by Chris Guise, PhD, CMC Connect, McCann Health Medical Communications and was funded by Pfizer Inc, New York, NY, USA, in accordance with Good Publication Practice (GPP3) guidelines (Ann Intern Med 2015;163:461-4).

**Authorship.** All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published.

**Disclosures.** J Woolcott, JC Cappelleri, P Sharma, and I Modesto are employees and shareholders of Pfizer Inc.

**Compliance with Ethics Guidelines.** This letter to the editor is based on a previously conducted study involving the use of anonymized electronic healthcare records. This study was not performed by the authors.

**Peer Review.** Please note, contrary to the journal's standard single-blind peer-review process, as a letter this article underwent review by a member of the journal's Editorial Board.

**Data Availability.** Data sharing is not applicable to this article as no datasets were generated or analyzed for this letter.

**Open Access.** This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are

included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

1. Moran K, Null K, Huang Z, Lissos T, Kane S. Retrospective claims analysis indirectly comparing medication adherence and persistence between intravenous biologics and oral small-molecule therapies in inflammatory bowel diseases. *Adv Ther.* 2019;36:2260–72.
2. Pope J, Keystone E, Jamal S, et al. Persistence of tofacitinib in the treatment of rheumatoid arthritis in open-label, long-term extension studies up to 9.5 years. *ACR Open Rheumatol.* 2019;1:73–82.
3. Harnett J, Gerber R, Gruben D, Koenig AS, Chen C. Evaluation of real-world experience with tofacitinib compared with adalimumab, etanercept, and abatacept in RA patients with 1 previous biologic DMARD: data from a U.S. administrative claims database. *J Manag Care Spec Pharmacy.* 2016;22:1457–71.
4. Pfizer Inc. XELJANZ prescribing information. 2019. <https://labeling.pfizer.com/ShowLabeling.aspx?id=959#S5.1>. Accessed Aug 20 2019.
5. Gemayel NC, Rizzello E, Atanasov P, Wirth D, Borsi A. Dose escalation and switching of biologics in ulcerative colitis: a systematic literature review in real-world evidence. *Curr Med Res Opin.* 2019;35:1911–23.
6. Moots RJ, Haraoui B, Matucci-Cerinic M, et al. Differences in biologic dose-escalation, non-biologic and steroid intensification among three anti-TNF agents: evidence from clinical practice. *Clin Exp Rheumatol.* 2011;29:26–34.
7. López-González R, León L, Loza E, Redondo M, Garcia de Yébenes M, Carmona L. Adherence to biologic therapies and associated factors in rheumatoid arthritis, spondyloarthritis and psoriatic arthritis: a systematic literature review. *Clin Exp Rheumatol.* 2015;33:559–69.