CORRECTION

Correction to: Sulodexide in the Treatment of Chronic Venous Insufficiency: Results of the All-Russian Multicenter ACVEDUCT Program

Andrey V. Chupin · Sergey E. Katorkin · Ivan I. Katelnitsky ·
Oksana V. Katelnitskaya · Igor I. Prostov · Alexey S. Petrikov ·
Alexander P. Koshevoi · Larisa F. Lyudkova

Published online: August 30, 2020
© Springer Healthcare Ltd., part of Springer Nature 2020

Correction to: Adv Ther (2020) 37:2071–2082
<https://doi.org/10.1007/s12325-020-01270-9>

In the original article, the “Duration, Types of Therapy and Dynamics of Improvements” topic published incorrectly.

The correct version is “The number of doctor visits for patients according to the protocol ranged from 3 to 5 (mean $\pm$ SD 3.6 ± 0.8 visits). The mean $\pm$ SD duration of sulodexide therapy

was 46.8 ± 13.1 days. The duration of therapy increased with CEAP class, from 29.9 ± 14.5 days in patients with C1 to 48.9 ± 13.8 days in those with C6. The majority of patients (63%; $n = 1417$) visited the doctor three times and used parenteral and oral formulations of sulodexide (85%; $n = 1203$) during the treatment period. Of the remaining 214 patients, 170 (71%) only received sulodexide orally and 44 (29%) only received the drug parenterally”.

The original article can be found online at <https://doi.org/10.1007/s12325-020-01270-9>.

A. V. Chupin (✉)
Federal Scientific Clinical Centre of the Federal
Medical and Biological Agency of Russia, Moscow,
Russia
e-mail: achupin@rambler.ru

S. E. Katorkin
Samara State Medical University of the RF Ministry
of Public Health, Samara, Russia

I. I. Katelnitsky · O. V. Katelnitskaya · I. I. Prostov
Rostov State Medical University of the RF Ministry
of Public Health, Rostov, Russia

A. S. Petrikov
Altai State Medical University of the RF Ministry of
Public Health, Barnaul, Russia

A. P. Koshevoi
Clinic “SibMedCentre”, Tomsk, Russia

L. F. Lyudkova
Krasnoyarsk Interregional Polyclinic No 1,
Krasnoyarsk, Russia