CORRECTION

Correction to: Treatment of Open-Angle Glaucoma and Ocular Hypertension with Preservative-Free Tafluprost/Timolol Fixed-Dose Combination Therapy: UK and Ireland Results from the VISIONARY Study

Ejaz Ansari · Jasna Pavicic-Astalos · Filis Ayan · Anthony J. King ·
Matthew Kinsella · Eugene Ng · Anca Nita on behalf of The VISIONARY Group

Published online: November 30, 2021
© Springer Healthcare Ltd., part of Springer Nature 2021

Correction to: Adv Ther (2021) 38:2990–3002
<https://doi.org/10.1007/s12325-021-01725-7>

On the original article, there is an error in the safety data: where the manuscript reports that there were 23/49 (46.9%) treatment-related AEs, it should in fact say there were 26/49 (53.1%) treatment-related AEs.

The original version has been corrected.

The original article can be found online at <https://doi.org/10.1007/s12325-021-01725-7>.

E. Ansari (✉)
Maidstone and Tunbridge Wells NHS Trust,
University of Kent at Canterbury, Kent, UK
e-mail: e.ansari@nhs.net

J. Pavicic-Astalos · E. Ng
Institute of Eye Surgery, UPMC Whitfield,
Waterford, Ireland

F. Ayan
Santen UK Ltd, St Albans, UK

A. J. King
Nottingham University Hospitals NHS Trust,
Nottingham, UK

M. Kinsella
Buckinghamshire Healthcare NHS Trust, Stoke
Mandeville Hospital, Aylesbury, Buckinghamshire,
UK

A. Nita
Royal Victoria Infirmary, The Newcastle Upon Tyne
Hospitals NHS Foundation Trust, Newcastle upon
Tyne, UK