

Erratum to: 2-D Duffing Oscillator: Elliptic Functions from a Dynamical Systems Point of View

Francisco Javier Molero · Martín Lara ·
Sebastián Ferrer · Francisco Céspedes

Published online: 6 March 2013
© Springer Basel 2013

Erratum to: Qual Theory Dyn Syst
DOI 10.1007/s12346-012-0081-1

In the original publication of the article, Figs. 3 and 5 are incorrect. The correct versions of Figs. 3 and 5 are given below:

The online version of the original article can be found under doi:[10.1007/s12346-012-0081-1](https://doi.org/10.1007/s12346-012-0081-1).

F. J. Molero (✉) · M. Lara · S. Ferrer · F. Céspedes
Departamento de Matemática Aplicada, Universidad de Murcia, Campus de Espinardo,
30071 Espinardo, Murcia, Spain
e-mail: franciscojavier.molero@um.es

M. Lara
e-mail: mlara0@gmail.com

S. Ferrer
e-mail: sferrer@um.es

F. Céspedes
e-mail: franciscojuan.cespedes@um.es




Fig. 3 Different types of 3D potentials represented from (29) over the parameter plane. Regions with bounded motions can be appreciated. All figures are “revolution plots” where $x^2 + y^2 \equiv r^2$




Fig. 5 Effective potential and reduced radial flows. The value of the integral is $G = 0.1$. Details of the region 4 are given in Fig. 7