

ERRATUM

Erratum to: Antiplatelet therapy with CABG: chaos in the Netherlands

F. W. A. Verheugt¹

Published online: 9 August 2017
© The Author(s) 2017. This article is an open access publication.

Erratum to:

Neth Heart J 2017

DOI 10.1007/s12471-017-1022-z

In the PDF version of the article, Tab. 1 was incorrectly displayed. The corrected version is shown below.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Table 1 Recommendation with level of evidence of timing and duration of antiplatelet therapy peri-CABG

Timing	Agent	Stable CAD without stent(s)	CABG Indication	
			Stable CAD with recent stent(s)	ACS with or without stent(s)
Preoperative	Aspirin	Continue (IA) ^a	Continue (IA)	Continue (IA)
	P2Y12	n.a.	Stop 5–7 days (IIA)	Stop 5–7 days (IIA) ^b
Postoperative	Aspirin	Lifelong (IA)	Lifelong (IA)	Lifelong (IA)
	P2Y12	n.a.	Day 1 restart (IIA) ^c	Day 1 restart (IIB) ^d

n.a. not applicable

^aUnless expected perioperative high bleeding risk

^bUnless continued clinical instability

^cDuration depending on stent type and patient characteristics

^dDuration at least 1 year after index ACS

The online version of the original article can be found under doi:
[10.1007/s12471-017-1022-z](https://doi.org/10.1007/s12471-017-1022-z).

✉ F. W. A. Verheugt
f.w.a.verheugt@olvg.nl

¹ Department of Cardiology, Onze Lieve Vrouwe Gasthuis, Amsterdam, The Netherlands

