Correction to: Beryl occurrences in the granitic complex of Guerioune, Laouni, Southeastern Algeria: electron microprobe, infrared spectroscopy, and fluid inclusions data

Ahmed Hamis^{1,2} · Mokrane Kesraoui² · Abdelhak Boutaleb² · Yves Fuchs³ · Christian Marignac⁴

Published online: 22 March 2021

© Saudi Society for Geosciences 2021

Correction to: Arabian Journal of Geosciences (2021) 14:357

<https://doi.org/10.1007/s12517-021-06675-6>

The original version of this paper was published with error. Figures 4 and 5 were interchanged. Also, Fig. 12 caption is incomplete. Given in this article are the correct figures and captions.

The original article has been corrected.

Fig. 4 Plots for the studied granites using geochemical diagrams of **a** alumina saturation index diagram of Rollinson (1993), **b** R1-R2 rock classification diagram of De La Roche et al., (1980), **c** Streckeisen & Le Maitre (1979)

The online version of the original article can be found at <https://doi.org/10.1007/s12517-021-06675-6>

✉ Ahmed Hamis
tizihamis@gmail.com

¹ Mouloud Mammeri University of Tizi Ouzou, F.B A S, Tizi Ouzou, Algeria

² Metallogeny and Magmatism Laboratory, (USTHB), Algiers, Algeria

³ Laboratoire Géomatériaux et Environnement, EA 4508, Université Paris Est Marne La Vallée, Champs-sur-Marne, France

⁴ Institut National Polytechnique de Lorraine, UMR 7566 du CNRS, Nancy, France

Fig. 5 **a** Outcrop of a pegmatite underlying a quartz vein (1-vein of quartz, 2-beryl bearing pegmatite), **b** Beryl in pockets hosted in pink biotite-granite, **c** Beryl in pocket in porphyritic biotite-granite, **d** Beryl surrounded by zinnwaldite in a miarolitic pegmatite in pink biotite-granite

Fig. 12 Fluid inclusion types: monophase liquid (**a**), monophase vapor (**b**), two-phase liquid-vapor (**c**), three-phase liquid-vapor-solid (**d**), necking-down (**e**)