



# Correction: Body size estimation method for seasonally growing farmed yellowtail *Seriola quinqueradiata* in an aquaculture net cage using a stereo camera

Kazuyoshi Komeyama<sup>1</sup> · Atsushi Ikegami<sup>2,3</sup> · Kichinosuke Fukuda<sup>4</sup> · Azusa Ishida<sup>3</sup> · Yuto Sasaki<sup>3</sup> · Hitoshi Maeno<sup>2</sup> · Shigeru Asaumi<sup>2</sup> · Takashi Uchida<sup>5</sup> · Yusei Katahira<sup>5</sup> · Akio Seki<sup>5</sup> · Tetsuo Oka<sup>5</sup> · Yasuhiko Shiina<sup>5</sup> · Yuki Takahashi<sup>1</sup>

© The Author(s), under exclusive licence to Japanese Society of Fisheries Science 2024

**Correction: Fisheries Science (2023) 90:227–237**  
<https://doi.org/10.1007/s12562-023-01736-z>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In this article the author name Kazuyoshi Komeyama was incorrectly written as Kazusyoshi Komeyama.

The original article has been corrected.

---

The original article can be found online at <https://doi.org/10.1007/s12562-023-01736-z>.

---

✉ Kazuyoshi Komeyama  
komeyama@fish.hokudai.ac.jp

<sup>1</sup> Faculty of Fisheries Sciences, Hokkaido University, 3-1-1, Minato-cho, Hakodate, Hokkaido 041-8611, Japan

<sup>2</sup> FURUNO Electric Co., Ltd., 9-52 Ashihara-cho, Nishinomiya, Hyogo 662-8580, Japan

<sup>3</sup> Graduate School of Fisheries Sciences, Hokkaido University, Minato-cho, Hakodate, Hokkaido 041-8611, Japan

<sup>4</sup> School of Fisheries Sciences, Hokkaido University, Minato-cho, Hakodate, Hokkaido 041-8611, Japan

<sup>5</sup> Maruha Nichiro Aquaculture Technology Development Centre Co., Ltd, 837, Kushi Bonotsu, Minamisatsuma, Kagoshima 898-0211, Japan