

Erratum to: Note on *Wahlgreniella nervata* (Gillette, 1908) (Hemiptera: Aphididae): a new pest of Damask rose in Turkey

Shalva Barjadze · İsmail Karaca · Bülent Yaşar ·
Nana Gratiashvili

Published online: 3 June 2011
© Springer Science+Business Media B.V. 2011

Erratum to: Phytoparasitica DOI 10.1007/s12600-011-0161-9

After submission of our manuscript, we found that *Wahlgreniella nervata* had been recorded from several locations in Turkey, but not from Isparta Province, earlier than our study (Çiraklı *et al.* 2008; Eser *et al.* 2009; Kaygin *et al.* 2009). However, Damask rose is a new host for this aphid species, as indicated in our manuscript.

References

- Çiraklı, A., Görür, G., & Işık, M. (2008). The determination of the aphid (Hemiptera: Aphididae) species of Central Denizli. *Selçuk Üniversitesi Ziraat Fakültesi Dergisi*, 22, 12–18 [in Turkish].
- Eser, S., Görür, G., Tepecik, İ., & Akyıldırım, H. (2009). Aphid (Hemiptera: Aphidoidea) species of the Urla district of İzmir region. *Journal of Applied Biological Sciences*, 3, 89–92.
- Kaygin, A. T., Görür, G., & Sade, F. C. (2009). Aphid (Hemiptera: Aphididae) species determined on herbaceous and shrub plants in Bartın Province in Western Blacksea Region of Turkey. *African Journal of Biotechnology*, 8, 2893–2897.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12600-011-0161-9>.

S. Barjadze (✉) · N. Gratiashvili
Entomology and Biocontrol Research Centre,
Ilia State University,
0179 Tbilisi, Georgia
e-mail: shalva1980@yahoo.com

İ. Karaca · B. Yaşar
Department of Plant Protection, Faculty of Agriculture,
Süleyman Demirel University,
32260 Isparta, Turkey