Correction to: Predation rate linked to life table of *Chrysoperla carnea* (Stephen) (Neuroptera: Chrysopidae) fed on small walnut aphid (*Chromaphis juglandicola*) (Kalt.) (Hemiptera: Aphididae): with population and predation projections

Mehmet Yılmaz · Evin Polat Akköprü

Published online: 24 July 2021
© Springer Nature B.V. 2021

Correction to: Phytoparasitica
<https://doi.org/10.1007/s12600-020-00834-3>

The original version of this article unfortunately contained a mistake in the Table 2 and in the body text, the lambda symbol (λ) can be seen in html format of Table 2 but was not presented in the pdf.

The corrected Table 2 is shown below.

Table 2 Reproduction and population parameters of *C. carnea* fed with *C. juglandicola*

Parameters	<i>C. carnea</i>
Fecundity (eggs/female)	257.7 ± 11.78
(APOP) (d)	8.00 ± 0.6746
(TPOP) (d)	33.00 ± 0.650
Oviposition days	43.37 ± 1.640
$r (d^{-1})$	0.072 ± 0.007
$\lambda (d^{-1})$	1.075 ± 0.007
R_0 (offspring)	62.48 ± 19.42
T (d)	57.16 ± 1.152

Standard errors were estimated by using the bootstraps technique with 100,000 resampling

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s12600-020-00834-3>

M. Yılmaz · E. P. Akköprü (✉)
Department of Plant Protection, Faculty of Agriculture, University of Van Yüzüncü Yıl, 65080 Van, Turkey
e-mail: polatevin@gmail.com.tr