EDITORIAL
ORAL FRAILITY
J.E. MORLEY

Corresponding author: John E. Morley, MB, BCh, Division of Geriatric Medicine, Saint Louis University, SLUCare Academic Pavilion, Section 2500, 1008 S. Spring Ave., 2nd Floor, St. Louis, MO 63110, USA, Email: john.morley@health.slu.edu

Key words: Oral health, eating assessment tool, oral frailty, nutrition.

The concepts of physical frailty and sarcopenia are now accepted as important geriatric syndromes (1-4). To enhance the recognition of these syndromes rapid screening tests, i.e., FRAIL and SARC-F, have been developed (5-12). Recently, it has been suggested that “oral frailty” should be considered a geriatric syndrome and regularly screened for in older persons (13, 14). “Oral frailty” can be defined as difficulty in chewing associated with age related changes in swallowing (presbyphagia) (15, 16). The Eating Assessment Tool – 10 (EAT-10) has been demonstrated to be an excellent screening tool for Oral Frailty (17, 18).

Table 1

D-E-N-T-A-L screening questionnaire (altered slightly from (42)) (3 or more points suggests oral frailty)

Dysphagia (3 points)/Dry mouth (1 point)
Eating difficulty (1 point)
No recent dental care (1 point)
Tooth or mouth pain (2 points)
Alterations or change in food selection (1 point)
Lesions, sores or lumps in mouth (2 points)

Older persons with sarcopenia have been commonly shown to have swallowing disorders (19-21). Decreased tongue pressure is common in older persons and is associated with dysphagia and poor nutrition (22-24). Persons with poor oral health are more likely to have aspiration malnutrition, sarcopenia and physical frailty (25, 26). It needs to be recognized that while sarcopenic dysphagia can cause undernutrition it can also be caused by severe malnutrition such as cachexia (27). Poor grip strength, walking speed and head lifting strength, all of which are indicative of sarcopenia has been associated with dysphagia (28, 29). Similarly tongue strength is associated with grip strength (30). Using ultrasound, older persons with sarcopenic dysphagia had smaller muscle mass (31). Utilizing high resolution manometry sarcopenic patients had lower mesopharyngeal contractility associated with upper esophageal sphincter integrity (32). Cognitive frailty is not uncommon with aging (33, 34) and dementia

associated with low calf circumference is associated with poor swallowing function (35). The bi-directional association between sarcopenia and presbyphagia is shown in Figure 1.

Figure 1
The Causes of “Oral Frailty”

A consensus paper on the diagnosis of dysphagia being related to sarcopenia has been published (36). The criteria required generalized sarcopenia in a person with dysphagia and the demonstration of a decline in oropharyngeal muscle mass. Other causes for sarcopenia need to be excluded.

The treatment of oral frailty requires appropriate treatment for problems with teeth and dentures and treatment of xerostomia if present. In addition, treatment requires caloric support with a supplement containing between 1.0 – 1.5 g/k/day high quality protein (37). For those with sarcopenia resistance training of the tongue and other oropharyngeal muscles needs to be carried out by a speech therapist (38). In older persons with aspiration pneumonia this rehabilitation needs to be introduced early in hospitalization to reduce mortality (39). Upright positioning in bed and early activity are essential components of the management.

Oral frailty is an important geriatric syndrome (40). Its major causes are poor dental status and sarcopenia. Treatment including resistance exercise of muscles involved in swallowing, needs to be instituted as soon as it is recognized. Screening for oral frailty should include screening with the EAT-10 questionnaire (41) or the D-E-N-T-A-L Questionnaire (Table 1) (42).

ORAL FRAILTY

Disclosures: The author declares there are no conflicts

References

- Dent E, Morley JE, Cruz-Jentoft AJ, et al. Physical frailty: ICFSR International Clinical Practice Guidelines for identification and management. *J Nutr Health Aging* 2019;34:771-787.
- Dent E, Morley JE, Cruz-Jentoft AJ, et al. International Clinical Practice Guidelines for Sarcopenia (ICFSR): Screening, diagnosis and management. *J Nutr Health Aging* 2018;22:1148-1161.
- Walston J, Bandeen-Roche K, Buta B, et al. Moving frailty toward clinical practice; NIA intramural frailty science symposium summary. *J Am Geriatr Soc* 2019;67:1559-1564.
- Bauer J, Morley JE, Schols AMWJ, et al. Sarcopenia: A time for action. An SCWD position paper. *J Cachexia Sarcopenia Muscle* 2019;10:956-961.
- Woo J, Yu R, Wong M, et al. Frailty screening in the community using the FRAIL scale. *J Am Med Dir Assoc* 2015;16:412-419.
- Woo J, Leung J, Morley JE. Comparison of frailty indicators based on clinical phenotype and the multiple deficit approach in predicting mortality and physical limitation. *J Am Geriatr Soc* 2012;60:1478-1486.
- Malmstrom TK, Miller DK, Morley JE. A comparison of four frailty models. *J Am Geriatr Soc* 2014;62:721-726.
- Bahat G, Yilmaz O, Kilic C, et al. Performance of SARC-F in regard to sarcopenia definitions, muscle mass and functional measures. *J Nutr Health Aging* 2018;22:898-903.
- Malmstrom TK, Miller DK, Simonsick EM, Ferrucci L, Morley JE. SARC-F: A symptom score to predict persons with sarcopenia at risk for poor functional outcomes. *J Cachexia Sarcopenia Muscle* 2016;7:28-36.
- Kurita N, Wakita T, Kamitani T, et al. SARC-F validation and SARC-F+EBM derivation in musculoskeletal disease: The SPSS-OK study. *J Nutr Health Aging* 2019;23:732-738. DOI: 10.1007/s12603-019-1222.
- Sanchez-Rodriguez D, Marco E, Davalos-Yerovi V, Lopez-Escobar J, et al. Translation and validation of the Spanish version of the SARC-F questionnaire to assess sarcopenia in older people. *J Nutr Health Aging* 2019;23:518-524.
- Bahat G, Oren MM, Yilmaz O, Kilic C, Aydin K, Karan MA, Bahat G, et al. Comparing SARC-F with SARC-CalF to screen sarcopenia in community living older adults. *J Nutr Health Aging* 2018;22:1148-1161. Doi: 10.1007/s12603-018-1139-9.
- Woo J, Tong C, Yu R. Chewing difficulty should be included as a geriatric syndrome. *Nutrients* 2018;10:1997. Doi: 10.3390/nu10121997.
- Watanabe Y, Okada K, Kondo M, et al. Oral health for achieving longevity. *Geriatr Gerontol Int* 2020;Apr 19. Doi: 10.1111/ggi.13921 [Epub ahead of print].
- Wakabayashi J. Frailty Aging
- Semba RD, Blaum CS, Bartali B, et al. Denture use, malnutrition, frailty, and mortality among older women living in the community. *J Nutr Health Aging* 2006;10:161-167.
- Takeuchi N, Sawada N, Ekuni D, Morita M. Oral diadochokinesis is related to decline in swallowing function among community-dwelling Japanese elderly: A cross-sectional study. *Aging Clin Exp Res* 2020;Apr 18. Doi: 10.1007/s40520-020-01547-7 [Epub ahead of print].
- Nishida T, Yamabe K, Ide Y, Honda S. Utility of the eating assessment tool-10 (EAT-10) in evaluating self-reported dysphagia associated with oral frailty in Japanese community-dwelling older people. *J Nutr Health Aging* 2020;24:3-8.
- Azzolino D, Damanti S, Bertagnoli L, Lucchi T, Cesari M. Sarcopenia and swallowing disorders in older people. *Aging Clin Exp Res* 2019;31:799-805.
- Kuroda Y, Kuroda R. Relationship between thinness and swallowing function in Japanese older adults: Implications for sarcopenic dysphagia. *J Am Geriatr Soc* 2012;60:785-1786.
- Yoshida M, Kikutani T, Tsuga K, et al. Decreased tongue pressure reflects symptom of dysphagia. *Dysphagia* 2006;21:61-65.
- Maeda K, Akagi J. Decreased tongue pressure is associated with sarcopenia and sarcopenia dysphagia in the elderly. *Dysphagia* 2015;30:80-87.
- Izumi M, Sonoki K, Ohta Y, et al. Impact of tongue pressure and peak expiratory flow rate on nutritional status of older residents of nursing homes in Japan: A cross-sectional study. *J Nutr Health Aging* 2020;24:512-517.
- Izumi M, Sonoki K, Ohta Y, et al. Impact of tongue pressure and peak expiratory flow rate on nutritional status of older residents of nursing homes in Japan: A cross-sectional study. *J Nutr Health Aging* 2020;24:512-517.
- Takagi D, Hirano H, Watanabe Y, et al. Relationship between skeletal muscle mass and swallowing function in patients with Alzheimer's disease. *Geriatr Gerontology Int* 2017;17:402-409.
- Zhang Y, Ge M, Zhao L, et al. Association between number of teeth, denture use and frailty: Findings from the West China health and aging trend study. *J Nutr Health Aging* 2020;24:423-428.
- Hironaka S, Kugimiya Y, Watanabe Y, et al. Association between oral, social, and physical frailty in community-dwelling older adults. *Arch Gerontol Geriatr* 2020;89:104-105.
- Wakabayashi H, Sashika H, Matsushima M. Head lifting strength is associated with dysphagia and malnutrition in frail older adults. *Geriatr Gerontol Int*. 2015;15:410-416.
- Wakasugi Y, Tohara H, Machida N, et al. Can grip strength and/or walking speed be simple indicators of the deterioration in tongue pressure and jaw opening force in older individuals? *Gerodontology* 2017;34:455-459.
- Sakai K, Nakayama E, Tohara H, et al. Diagnostic accuracy of lip force and tongue strength for sarcopenic dysphagia in older inpatients: A cross-sectional observational study. *Clin Nutr* 2019;38:303-309.
- Ogawa N, Mori T, Fujishima I, et al. Ultrasonography to measure swallowing muscle mass and quality in older patients with sarcopenic dysphagia. *J Am Med Dir Assoc* 2018;19:516-522.
- Kaneida K, Fujishima I, Wakabayashi H, et al. Relationship between tongue pressure and pharyngeal function assessed using high-resolution manometry in older dysphagia patients with sarcopenia: A pilot study. *Dysphagia* 2020; Mar 5 doi: 10.1007/s00455-020-10095-1 [Epub ahead of print].
- Kwan RYC, Leung AYM, Yee A, Lau LT, Xu XY, Dai DLK. Cognitive frailty and its association with nutrition and depression in community-dwelling older people. *J Nutr Health Aging* 2019;23:943-948.
- Moon JH, Huh JS, Won CW, Kim HJ. Is polypharmacy associated with cognitive frailty in the elderly? Results from the Korean frailty and aging cohort study. *J Nutr Health Aging* 2019;23:958-965.
- Nishida T, Yamabe K, Honda S. Dysphagia is associated with oral, physical, cognitive and psychological frailty in Japanese community-dwelling elderly persons. *Gerodontology* 2020;37:185-190.
- Fujishima K, Fujii-Kurachi M, Arai H, et al. Sarcopenia and dysphagia: Position paper by four professional organizations. *Geriatr Gerontol Int* 2019;19:91-97.
- Bauer J, Biolo G, Cederholm T, et al. Evidence-based recommendations for optimal dietary protein intake in older people: A position paper from the PROT-AGE study group. *J Am Med Dir Assoc* 2013;14:542-559.
- Park J-S, Lee S-H, Jung S-H, et al. Tongue strengthening exercise is effective in improving the oropharyngeal muscles associated with swallowing in community-dwelling older adults in South Korea: A randomized trial. *Medicine (Baltimore)* 2019;98(40):e17304.
- Shiraishi A, Yoshimura Yoshihiro, Wakabayashi H, Tsuji Y, Shimazu S, Jeon S. Impaired Oral health status on admission is associated with poor clinical outcomes in post-acute inpatients: A prospective cohort study. *Clin Nutr* 2019;38:2677-2683.
- Payne M, Morley JE. Editorial: Dysphagia, dementia and frailty. *J Nutr Health Aging* 2018;22:562-565.
- Speyer R, Cordier R, Kertscher B, Heijnen BJ. Psychometric properties of questionnaires on functional health status in oropharyngeal dysphagia: A systematic literature review. *Biomed Res Int* 2014;458678. Doi: 10.1155/2014/458678. Epub 2014 Apr 29.
- Bush LA, Horenkamp, Morley JE, Spiro 3rd A. D-E-N-T-A-L: A rapid self-administered screening instrument to promote referrals for further evaluation in older adults. *J Am Geriatr Soc* 1996;44:979-981.